Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 88 (2015)

Heft: 3-4

Artikel: Assessing the stability of stonefly (Plecoptera) biodiversity in the Swiss

National Park

Autor: Knipsel, Sandra / Lubini, Verena

DOI: https://doi.org/10.5169/seals-583848

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

MITTEILUNGEN DER SCHWEIZERISCHEN ENTOMOLOGISCHEN GESELLSCHAFT BULLETIN DE LA SOCIETE ENTOMOLOGIQUE SUISSE

88: 257–271, 2015 doi:10.5281/zenodo.33951

Assessing the stability of stonefly (Plecoptera) biodiversity in the Swiss National Park

Sandra Knispel¹ & Verena Lubini²

- ¹ Bureau Akuatik, écologie aquatique et appliquée, route d'Essertines 3, CH-1416 Pailly; sandra.knispel@akuatik.ch
- ² Büro für Gewässerökologie, Eichhalde 14, CH-8053 Zürich

An inventory of stonefly species in the alpine Swiss National Park was conducted during 2011 and 2012 over 33 sites in various stream types, from springs and brooklets to larger rivers. 34 species were identified, representing 60% of the central eastern Alps fauna. Most species were widespread alpine species, 5 were alpine endemics and 3 species were threatened in Switzerland. Cold stenotherm and rheophilic species dominated, but eurytherm and limnophilic species appeared in streams with high habitat diversity. Historical stonefly data, collected between 1934 and 1964, enabled a long-term comparison. The stability of the stonefly fauna over more than half a century is remarkable. The low impacted stream morphology and flow regime, as well as diversity of stream types and substrate are important biodiversity drivers. In the present study one species was missing, but has always been rare in the Park. Three species are new for the Park. One species expands to higher locations and this could be related to climate changes. The present species data basis is particularly valuable for assessment of further impacts, as some changes are only perceptible at the species level.

Keywords: Plecoptera, freshwater ecology, alpine streams, biodiversity, species, protected area

INTRODUCTION

The Swiss National Park (SNP) includes a complex hydrological network in a multitude of small catchment areas including springs, brooklets, brooks, torrents, rivers, ponds and lakes. These aquatic environments shelter a characteristic fauna including the insect order Plecoptera (Stoneflies). Aquatic insects in this region have been studied by Adolf Nadig between 1934 and 1942 (Nadig 1942), and by Jacques Aubert from 1949 to 1964 (Aubert 1965). Nadig specialized in spring fauna, whereas Aubert concentrated on the insect order Plecoptera. He collected in the SNP as well as in the neighbouring regions, and also studied the specimens collected by Nadig. The results were presented in a volume of the series dedicated to the research activities in the SNP (Aubert 1965). Stoneflies in the SNP were poorly studied during the following decades (less than 10 new records in the national database CSCF, as of October 2013). Nevertheless, macroinvertebrates were still collected during long term monitoring programs and stonefly records have recently been added to the national database. A second inventory of the stonefly species was conducted in 2011 and 2012, in order to assess today's species composition and distribution. Comparison with historical data is therefore possible and sets the basis for a long term monitoring.




Fig. 1: Map of sites studied in 2011-2012 (o) and by Aubert (1965) (●).

MATERIAL & METHODS

Study area

The SNP is located at the very south-east corner of Switzerland in the canton of Graubünden (Fig. 1). Created in 1914, it is the oldest national park in the Alps and central Europe, classified by IUCN as a category 1 nature reserve. It covers an area of 170 km² at a mean altitude of 2320 m asl. 31 % of the SNP is covered by forest, 17 % by grassland, the rest has no vegetation cover and glaciers are absent.

Characteristics of the catchment area

The hydrological network of the SNP belongs to the Inn catchment and therefore its waters drain via the Danube River into the Black Sea. Streams have a natural alpine nival runoff regime, except the River Spöl, which is regulated by the Livigno dam since 1962, and the River Clemgia, which is regulated since 1967. The Ofenpass road crosses the Park and some streams are very locally impacted (bridges). Beyond that the freshwater system is morphologically pristine. Most streams are classified as alpine, situated over 1800 m except the Spöl valley, the

River Clemgia and the lower section of the River Fuorn (beneath II Fuorn). The maximum elevation of nearly all catchment basins is above 3000 m asl. Water temperatures rarely exceed 10 °C (Eisenhut 2013). The flow regime is very dynamic, with low flow in winter, higher and fluctuating flows during summer, which is influenced by snowmelt and heavy thunderstorms. Rivers have a wide and rocky streambed. Springs are rheocrenes, except one limnocrene, originating from a pool at II Fuorn. Because of mostly permeable soils, ponds and lakes are the exception, and the Macun-lakes belong to the SNP since 2000.

Tab. 1: Sampling sites and sampling periods.

Stream, location	coord. X	coord. Y	altitude (m)			12			
						5.20	2	7	2
				11	11	15.	201	201	201
				20	20	or	3.6.	7.	.9.
				1-4.6.201	5-8.9.201	2-3.4 or 15.5.2012	5-18.6.2012	24-31.7.2012	13-16.9.2012
Ova dal Fuorn, Il Fuorn	811848	171847	1781	X	X	X		- 7	_
God sur Il Fuorn, brook	811603	171869	1777	X	X	X			
Champlönch, brook	810000	172760	2010	X	X				X
Ova da Val Ftur, Val Ftur	810717	172511	1871	X	X				
Ova da Stabelchod	714425	171525	1916	X	X	X			
Ova da Val Chavagl	812830	171530	1857	X	X				
Ova dals Pluogls	815740	170615	1940	X	X				
Alp La Schera, brook	811028	169370	2087	X	X				
Val Ftur, brook	810740	172470	1888	X	X				
Clemgia, Pradatsch	819055	179221	1666	X	X				
Spring, Spöl right bank	809885	169378	1754				X	X	X
Spring, Plan da l'Acqua Suot	809416	169065	1752				X	X	X
Ova da l'Acqua, Plan da l'Acqua Suot	809450	169115	1745				X	X	X
Spring, Spöl left bank	809619	169070	1712				X	X	X
Ova da Cluozza, Chamanna	804860	171536	1810				X	X	X
Spring near path, Val Cluozza	804604	171842	1820				X	X	X
Spring, opposite to Val dal Fuorn	812581	171598	1841				X	X	X
Spring, God dal Fuorn (GFQ1)	812178	171678	1822				X	X	X
Ova da Trupchun, Alp Trupchun	802059	163915	2024				X	X	X
God Trupchun, brook	801351	164225	1991				X	X	X
Ova da Trupchun, Alp Purcher	800250	164885	1868				X	X	X
Spring, Alp Purcher	800156	164901	1855				X	X	X
Spring Dschembrina, Val Trupchun	801157	164451	1975				X	X	X
Spring, Ova dals Pluogls	816117	170555	1969					X	X
Ova dal Fuorn, under O. Stabelchod	814320	171195	1891			X			
Ova dal Fuorn, Punt La Drossa	810560	170755	1695			X			
Ova da Val Ftur, lower Val Ftur	811271	171607	1764			X			
Macun, Aua da Zeznina	806193	178766	2594					X	
Macun, N Lai Sura	805748	178832	2631					X	
Macun, Lais d'Immez	805805	178661	2623					X	
Ova da Cluozza, Plaun Sassa	805532	169254	1935				X		
Punt Periv, brook	810023	168814	1707				X		
Macun, springbrook	805579	178503	2662					X	

We considered 5 types of freshwater habitats:

- mid-sized streams, flowing on large valley floors: rivers
- lateral 2nd order tributaries with variable flow over the year and steep slope
 (> 7%): torrents
- small 1st order streams flowing in forested or open areas (former pastures),
 gentle slope (< 3 %): brooks
- springs and their spring brooklets: spring brooklets
- standing water bodies: ponds and lakes

Sites and sampling methods

In 2011 and 2012, 23 primary sites were sampled two or three times each (Tab. 1) for three groups of aquatic insects, Ephemeroptera, Plecoptera and Trichoptera. These sites are representative of stream types and different valleys in the SNP, and had already been sampled by Aubert or Nadig (Fig. 1). Some secondary sites (Tab. 1) were sampled to look for early emerging species or using the opportunity to access and sample particular systems (e.g. Macun). Sampling methods were combined. Larvae were collected in all aquatic habitats using a kick-net and qualitative sampling by hand. Adults were collected in the riparian vegetation, using a sweep net or a beating sheet. Each site was prospected for 30 to 45 minutes depending on the stream size. Insects were preserved in 80 % ethanol and the probe was labelled with date, name of the site and coordinates (Garmin etrex). Sampling periods were chosen according to the phenology of stonefly species: April-May (remaining snow cover in 2012), June (beginning of vegetation period), end of July, September. Stoneflies were identified to species level (Lubini et al. 2012a), the specimens preserved and labelled as recommended (Stucki 2012) and deposited in the collections of the Musée Cantonal de Zoologie in Lausanne (Switzerland). Data are recorded in the national database (CSCF, Neuchâtel).

Other data sources

Stoneflies from two long term monitoring programs focusing on benthic fauna were controlled or identified: monitoring of hydropower effects in the Spöl river basin (Mürle & Ortlepp 2012), biodiversity monitoring in ponds at the Macun Lakes region (Indermühle & Oertli 2007). Data from the national database (CSCF, Neuchâtel) was provided, including historical data from Nadig (1942) and Aubert (1965) who sampled 90 stream sites in the SNP region, 40 of which were located inside the SNP limits.

Data analyses

The species were classified considering their ecological preferences for current and temperature using the system of Moog (1995, 2002) completed by Graf *et al.* (2009). Temperature preference was not documented for *Taeniopteryx kuehtreiberi*, and current preference was not documented for *Leuctra rosinae* and *Protonemura algovia*. For *Rhabdiopteryx harperi* and *Leuctra muranyii*, the preferences of *R. alpina* and *L. braueri*, respectively, were used because the species have ecological similarities concerning zonation and habitat.

Similarity (%) of the species composition between different streams was calculated using Jaccard-index (Jaccard 1908). Values ≥ 64 % are considered as simi-

Tab. 2: Species list: Species in **bold** are new for the SNP; — A: Categories as in Red Data List of Switzerland (Lubini *et al.* 2012b); — B: Conservation priorities as in the Swiss list of national priority species (OFEV 2011); — C: Responsibility as in the Swiss list of national priority species (OFEV 2011); — in grey, species not found since 1951.

Notes from 1 to 11: leg. or det. when not by the authors, and species names before nomenclature changes and corrections vs Aubert (1965); — 1: syn. *Perlodes intricata* (Pictet 1841); — 2: first records leg. Ortlepp 2008, syn. *Perla maxima* (Scopoli 1763); — 3: re-identification of *Chloroperla tripunctata* (Scopoli 1763); — 4: syn. *Chloroperla montana* (Pictet 1841); — 5: confused with *Rhabdiopteryx alpina*; — 6: leg. Ortlepp, det. Mürle; — 7: re-identification of *Protonemura nimborella* Moselyi 1930; — 8: leg. Mürle, det. Mürle; — 9: leg. Ortlepp, det. Knispel; — 10: re-identification of *Leuctra braueri* Kempny 1898; — 11: leg. Aubert, det. Aubert.

			Threatening category ^A	Alpine endemite (E)	National priority ^B	National responsibility ^c	Last records in the SNP
Perlodidae	Dictyogenus alpinum	(Pictet, 1841)					2012
	Dictyogenus fontium	Ris, 1896	NT				2012
	Isoperla rivulorum	(Pictet, 1841)					2012
	Perlodes intricatus ¹	(Pictet 1841)					2013
Perlidae	Perla grandis ²	Rambur 1842					2012
Chloroperlidae	Chloroperla susemicheli 3	Zwick 1967					2012
	Siphonoperla montana 4	(Pictet 1841)	NT				2012
Taeniopterygidae	Rhabdiopteryx alpina	Kühtreiber, 1934	NT		4	2	2012
	Rhabdiopteryx harperi ⁵	Vinçon & Murányi, 2008	VU	Е	3	2	2012
	Rhabdiopteryx neglecta	(Albarda, 1889)					2012
	Taeniopteryx kuehtreiberi 6	Aubert, 1950					2013
Nemouridae	Amphinemura standfussi	(Ris, 1902)	NT				2012
	Protonemura algovia ⁷	Mendl 1968	VU	Е	3	2	2012
	Protonemura brevistyla	(Ris, 1902)					2012
	Protonemura lateralis	(Pictet, 1836)					2012
	Protonemura nimborum	(Ris, 1902)					2012
	Protonemura nitida	(Pictet, 1835)					2012
	Nemoura cinerea	(Retzius, 1783)					2011
	Nemoura mortoni	Ris, 1902					2012
	Nemoura obtusa	Ris, 1902	NT		4	2	2012
	Nemoura sinuata	Ris, 1902	NT		4	2	2012
	Nemoura undulata	Ris, 1902	CR	Е	1	2	2012
	Nemurella pictetii	Klapalek, 1900					2013
Capniidae	Capnia nigra	(Pictet, 1833)					2012
	Capnia vidua	Klapalek, 1904	NT				2012
Leuctridae	Leuctra alpina	Kühtreiber, 1934					2012
	Leuctra armata	Kempny, 1899	NT		4	2	2012
	Leuctra helvetica 8	Aubert, 1956	EN	Е	2	2	2014
	Leuctra inermis	Kempny, 1899					2012
	Leuctra major	Brinck, 1949					2012
	Leuctra moselyi 9	Morton, 1929					2012
	Leuctra muranyii 10	Vinçon & Graf 2011		Е			2012
	Leuctra rauscheri	Aubert, 1957	NT				2012
	Leuctra rosinae	Kempny, 1900					2012
	Leuctra teriolensis 11	Kempny, 1900					1951

Tab. 3: Species composition and abundances in the SNP streams studied since 2011. Presence of adult stages (A), larval stages (L) including exuviae. — *sampled by Mürle & Orlepp (2012) during the ongoing Spöl basin monitoring; — **species sampled in a stream type where it can obviously not originate from; not taken into account in the Jaccard index and the stream type diversity.

	Sites	Spól*	Clemgia, Pradatsch	Ova dal Fuorn	Ova Cluozza	Ova Trupchun	Ova Ftur	Ova da Val Chavagl	Ova da l'Acqua, Plan da l'Acqua Suot	Ova da Stabelchod	God Trupchun, brook
	Freshwater habitat type (number of sites when grouped)	river (4)	river	river (4)	river (2)	river (2)	torrent (2)	torrent	torrent	torrent	torrent
Perlodidae	Dictyogenus alpinum		5L	1A 8L	10L	3L	2L	8L	6L	1L	1L
	Dictyogenus fontium Isoperla rivulorum Perlodes intricatus	2L	8A	1A** 10L 9L	2L 2A 1L	1L		1L 1L	1A 2L 1L	2L	
Perlidae	Perla grandis			2L							
Chloroperlidae	Chloroperla susemicheli	1A 1L	1L		1L				1A 19L		
	Siphonoperla montana		3L	3L	2A 7L	1L		10L	2L	1L	
Taeniopterygidae	Rhabdiopteryx alpina				1A						
	Rhabdiopteryx harperi			12L			6A 14L			8A 13L	
	Rhabdiopteryx neglecta			2A 10L			8L				1A
	Taeniopteryx kuehtreiberi	1L*		2L*							
Nemouridae	Amphinemura standfussi										
	Protonemura algovia			1L	1L						
	Protonemura brevistyla			1A 2L	20L	8L		1L	19L	11L	8L
	Protonemura lateralis		2L	2L					2L		1A 1L
	Protonemura nimborum			15L			2L			1A	2A
	Protonemura nitida	4A	1A 5L	3L							
	Nemoura cinerea										
	Nemoura mortoni		9L	7L	14L			1L	3L	5L	5L
	Nemoura obtusa										
	Nemoura sinuata									17	
	Nemoura undulata		1A**						101	1L	
Capniidae	Nemurella pictetii Capnia nigra	3A3L	1A** 1A						12L		
Capinidae	Capnia nigra Capnia vidua	3A 3L	IA	14A 2L							
Leuctridae	Leuctra alpina		1A	14A 2L							
	Leuctra armata		IA								
	Leuctra inermis	101 A	4A	5A	2A			1A	1A	2L	
	Leuctra helvetica	10171	7/1	6A *	D/1			1/1	1/1	212	
	Leuctra major	4A		011	9A				1A		
	Leuctra major	6A									
	Leuctra muranyii			1A**							
1	Leuctra rauscheri	2A	2A	7A		1A	1A				
	Leuctra rosinae	16A	2A	4A	14A 15L	2A		2A	1A	4A 1L	1A

lar, and values between 50 and 63% as a weak similarity (Smukalla & Friedrich 1994). Similarity was calculated for streams sampled with comparable effort (3 sampling periods). For streams sampled at different sites, a species list for the whole stream is given in Tab. 3 (e.g. Ova dal Fuorn, Ova da Val Ftur). Species in the adult stage were sometimes sampled in a stream type where they can obviously not originate from. These occurrences were pointed out in Tab. 3 and not taken into account in stream type diversity (Tab. 4) and Jaccard-index (Tab. 5).

RESULTS

Biodiversity and stream type

Today the SNP shelters 34 stonefly species (Tab. 2), which represent nearly 60 % of all 57 species listed in the "central eastern Alps" biogeographic zone (Lubini *et*

God sur II Fuorn, brook	Val Ftur, brook	Ova dals Pluogis	Champlönch, brook	Alp La Schera, brook	Spring near path, Val Cluozza	Spring, God dal Fuorn (GFQ1)	Spring, Alp Purcher	Spring, Plan da l'Acqua Suot	Spring, Spöl right bank	Spring, Spöl left bank	Spring Dschembrina, Val Trupchun	Spring, Ova dals Pluogls	Spring, opposite to Val dal Fuorn	Macun
brook	brook	brook	brook	brook	spring brooklet	spring brooklet	spring brooklet	spring brooklet	spring brooklet	∏spring brooklet	spring brooklet	spring brooklet	spring brooklet	lake
4L 12L 1A	2L 1L	1A 13L 3L	1L 5L	1A3L 2L	4A 5L	1A	1L 10L 2L			IL		12A 5L	3L	
								1A 4L					2A	1A**
1A	4A	9 1 OI	15A 3L	2A	04.51	1A** 2A						1.4	1T	
1L 1A 3L 1L 6L	1L	8A 9L 6A 6L	3L 2L	1A 5L	1A SL 1A	2A	4A 9L		2L	1A 7L		1A 6A 9L	1L 1A 16L	10A 11L
1A 7L 4L 1A	4L 1L 1L	3A 18L	5L	3A 2L 2A 10L	1A 1L 1A	1L	1A3L	5L	1A	2A 1L	1A 6L	1L	2L 1L 8A 5L	
1A		4L				5A 27L	12A 19L	9L	4A			15A 14L		
1A	1L	3A	14A	60A 2L 2A				2A**	6A 3A**	1A 15A**		2A	1A 2A**	
4L 9L 1A	1A 8L	24A 4A 4L	1L	20A 4L	32L	10A 22L	47A 2A		5A 2L	21A 9L	1A		11A	

al. 2012a, Gonseth et al. 2001). The fauna is composed mainly of widespread and characteristic alpine species, with additional presence of five endemic alpine species. Nemoura undulata, found in a spring brooklet and in the Ova Stabelchod (torrent), is a species limited to the central Alps, known from the Engadin (Inn valley), as well as Leuctra muranyii, Protonemura algovia and Rhabdiopteryx harperi. The geographic distribution of Protonemura algovia covers the entire Swiss Alps and was found in the SNP in the rivers Fuorn and Cluozza. Leuctra muranyii, previously confused with L. braueri, is a crenophilic species collected in several springs and spring brooklets in the SNP. In Europe its distribution is limited to the south-eastern Alps, from Italy to the Austrian upper Inn valley (Vinçon & Graf 2011) and, in Switzerland, is limited to the SNP, to the Val Müstair, lower Engadin and Puschlav valleys. Leuctra helvetica is an extensive alpine and epirhithral species, originally described from the Engadin (Val Sesvenna). In Switzerland it is limited to the

		Spring brooklets			
	Rivers (N=5)	(N=9)	Brooks (N=5)	Torrents (N=5)	Lakes (N=2)
Total species	25	20	18	17	1
Arithmet. mean	12 ± 5	6 ± 2	10 ± 3	8 ± 3	
Maxima/Minima	19/6	10/2	16/8	12/5	

Engadin (Inn valley) and the Val Colla on the southern slope of the Alps. Detailed species composition and their abundances in the streams are given in Tab. 3. One third of the species are threatened or nearly threatened in Switzerland (Lubini *et al.* 2012b). *Nemoura undulata* is critically endangered (category CR), *Leuctra helvetica* is endangered (category EN), *Protonemura algovia* and *Rhabdiopteryx harperi* vulnerable (category VU). *Nemoura undulata* has a very high national priority (OFEV 2011).

The highest species diversity is found in rivers (Tab. 4) with some characteristic species: Capnia nigra, C. vidua, Perla grandis, Protonemura algovia (Tab. 3). Few species live in torrents where waterflow is reduced after the snowmelt and which even dry out in summer, like some of the streams at Alp Trupchun. Torrents and rivers have species in common, which were not found in other stream types in larval stages: Rhabdiopteryx harperi, R. neglecta and Siphonoperla montana. Chloroperla susemicheli and Dictyogenus alpinum were rare in brooklets. Lakes only shelter larvae of Nemurella pictetii. Numerous species live in springs and brooks.

Comparing communities between stream types shows that very few streams have a similar species composition, which might be related to distinct ecological characteristics, although stream physiography might be similar. Highest similarity was shown for a small group of brooks (Tab. 5). The brook at Alp La Schera had a similar stonefly fauna to that at Champlönch and the forested brook in Val Ftur (\geq 64 % of similarity) because the species composition included strongly creno-

Tab. 5: Similarity (%) of the species composition between different rivers, torrents and brooks (Jaccard 1908), 2011-2012 observations. Values \geq 64 % (in bold) are considered as similar (Smukalla & Friedrich, 1994).

				riv	vers			torre	ents				bro	oks
		Clemgia	Ova dal Fuorn	Ova Cluozza	Ova Trupchun	Ova Ftur	Ova da Val Chavagl	Ova da l'Acqua	Ova da Stabelchod	God Trupchun	God sur il Fuorn	Val Ftur	Ova dals Pluogls	Champlönch
	Ova dal Fuorn	45												
rivers	Ova Cluozza	41	41											
	Ova Trupchun	38	29	38										
	Ova Ftur	7	21	7	11									
nts	Ova da Val Chavagl	43	42	67	56	9								
torrents	Ova da l'Acqua	50	41	7 1	38	7	67							
2	Ov da Stabelchod	29	45	47	33	27	64	47						
	God Trupchun	27	37	40	30	38	36	36	42					
	God sur il Fuorn	33	40	27	22	5	33	40	30	28				
	Val Ftur	17	12	17	7	0	13	17	6	14	47			
S	Ova dals Pluogls	31	33	31	36	0	42	50	27	33	47	29		
brooks	Champlönch	18	13	11	8	0	14	18	6	15	33	70	31	
brd	Alp La Schera	29	21	22	14	0	29	29	24	21	53	73	36	70

Tab. 6: Similarity (%) of the species composition between different spring brooklets (Jaccard 1908), 2011-2012 observations. Values ≥ 64 % (in bold) are considered as similar (Smukalla & Friedrich, 1994).

						sp	oring b	rookl	ets
		Val Cluozza	God dal Fuorn	Pluogls	Val dal Fuorn	Alp Purcher	Dschembrina	Spöl rhight bank	Spöl left bank
	God dal Fuorn	18							
	Ova dals Pluogls	18	33						
ets	Val dal Fuorn	45	14	33					
ķ	Alp Purcher	15	30	40	29				
0,0	Dschembrina	13	0	0	20	11			
so D	Spöl right bank	20	38	22	25	8	0		
spring brooklets	Spöl left bank	20	38	20	21	18	0	67	
	Plan dal Acqua Suot	10	29	13	8	10	0	14	14

philic species such as *Amphinemura standfussi*, *Leuctra muranyii*, *L. armata*, *N. sinuata* and *Dictyogenus fontium*, beside species with larger ecological preferences. Within the group of torrents, only Ova da Val Chavagl, Ova da l'Acqua and Ova Stabelchod had a species composition considered as similar (Tab. 5). Ova Cluozza had also a similar composition to two of these torrents. No similarity between the species composition has been demonstrated for rivers, nor for spring brooklets with the exception of two in the Val Spöl (Tab. 6).

Ecological traits of the stonefly fauna

Most species are considered as being stenoecious, particularly regarding temperature tolerance, and most are cold stenotherm (Fig. 2). Cold and very cold stenotherm species were found in all stream types, as for example larvae of *Perlodes intricatus* (Tab. 3). However, five species, *Nemoura cinerea*, *Protonemura nitida*, *Nemurella pictetii*, *Leuctra inermis* and *Leuctra major* are eurytherm. The larvae of these species develop in habitats with wider temperature ranges (Tab. 3), such as lower Spöl, Clemgia and Ova dal Fuorn up to Il Fuorn as well as some tributaries (*Protonemura nitida*, *Leuctra inermis*), brooks with high amount of particulate organic matter (Champlönch, *Nemoura cinerea*) or deep in the sediment pores of rivers (*Leuctra major*). *Nemurella pictetii* has been found in some springs as well as in the lakes of Macun. These species occur all over Switzerland down to low altitudes.

The high amount of rheophilic species (Fig. 2) has to be related with the prevalence of steep streams and fluctuating discharge during summer. Species with no current preference (e.g. *Amphinemura standfussi*, *Protonemura nitida*) or limnophilic species (e.g. *Nemoura cinerea*, *Nemurella pictetii*) occur mainly in lakes, spring brooklets, but also in streams with slight slopes and calm, shallow riverbanks (e.g. transition to fens/marshland).

Temperature and current preferences of occurring species were not very different between stream types (Fig. 3). Rivers and torrents have a very low amount of limnophilic or current indifferent species. Cold stenotherm species were frequent




Fig. 2: Number of stonefly species in the SNP with distinct current (upper; n = 32) and temperature (lower; n = 33) preferences.

in springs and brooks, and their abundance was higher than in larger streams. The situation is opposite for warm tolerant species or temperature indifferent species.

Temporal development of the stonefly fauna

Aubert (1965) recorded 38 stonefly species. Four of them had been located outside the SNP in the neighbouring regions: Amphinemura sulcicollis, Perla grandis (syn. Perla maxima), Leuctra mortoni, Leuctra moselyi. A control of the «catalogue des espèces» in Aubert (1965) and the location of sampling sites gave evidence for two further species sampled outside of the SNP limits: Leuctra sesvenna and Leuctra handlirschi. L. sesvenna is mentioned in Aubert (1965) from the Aua da Sesvenna, Marangun, between 2300 and 2500 m asl. This location is situated outside the SNP, just east of Val Nüglia and Val Foraz. L. handlirschi is mentioned in Aubert (1965) only from the Aua da Zeznina, at Val Zeznina at 1930 m asl, located outside the SNP, just north of the Macun lakes. Aubert (1965) thus finally recorded 32 stonefly species for the SNP. Descriptions of new species have been published since then. Material of the concerned species, collected before 1965 by Nadig or Aubert (today in museum collections in Lausanne or Chur), has been revised during the Red List Project (Lubini et al. 2012b) or the present work and recorded in the national database. Unverified data was put aside (e.g. Leuctra spp. larvae, or specimens cited in publications, but not found in the corresponding museum collections). Corrections and nomenclature changes have been mentioned in Tab. 2.

The current stonefly inventory of 34 species includes two more than nearly 60 years ago. Examining changes in composition and distribution provided evidence that one species, *Leuctra teriolensis*, has not been collected in the SNP again. In fact, the only verified record before 1965 in the SNP is from the river Ova Spin at




Fig. 3: Average composition of species (%) relative to their current (upper; n = 32) and temperature (lower; n = 33) preferences in different stream types.

Ova Spin. It means that the species was rare and could not be confirmed until now in the SNP. It was recently collected near the SNP, at Tschierv (Val Müstair) and at Susch (Inn Valley) (CSCF, Neuchâtel). The species is, however, common in the Alps. Three species collected in 2011 and 2012 were new for the SNP: *Leuctra moselyi*, *Perla grandis* and *Rhabdiopteryx harperi*. *R. harperi* was recently taken out from the *R. alpina*-species complex and described as new by Vinçon & Murányi (2009). Due to the lack of adult male specimens in Aubert's material, its presence in the SNP before 1965 cannot be confirmed or invalidated. Concerning *Perla grandis* and *Leuctra moselyi*, both species are widespread in Switzerland, mainly in streams below 1400 m as shown from *P. grandis* larvae records (Fig. 4). Both species were collected recently in the SNP with altitudinal limits of 1960 m asl (Ova dal Fuorn) and 1510 m asl (lower Spöl), respectively.

DISCUSSION

The results of the present study emphasized the stability of the stonefly fauna in the SNP over more than half a century. With one exception, all species, including endemic and rare species, were still sheltered in the protected area. Except for the River Spöl, this could be attributed to low impact on streambed morphology and the pristine runoff regimes, which are, in addition to high permeability of the riverbank substrate, important biodiversity drivers (Oswald 1990; Burgherr *et al.* 2003). Refuge habitats were provided by high streambed structure diversity combined with different stream types, which enhance ecosystem stability.

With 34 Plecoptera species, the present diversity reached the expected level, taking into account the harshness of the environmental conditions induced by nival flow regimes, dynamic stream morphology, low water temperatures, biogeographic

region of the central Alps; all are important drivers of the stonefly species composition. Biodiversity of the three aquatic insect orders Ephemeroptera, Plecoptera and Trichoptera in Switzerland is the highest between 700 and 1300 m asl (Altermatt *et al.* 2013) and then decreases with increasing altitude. These results were supported by different studies, like the pre-alpine Sense catchment study between 577 and 1400 m asl that registered 52 stonefly species (Zurwerra *et al.* 2000). Similar situations were observed in the Italian western Alps (Malicky *et al.* 2007) and the Isar catchment in Germany (Hering 1995).

Similarity of total Plecoptera species composition between streams was very low, except for a few streams of various types. Presence of some species seemed highly influenced by typology, like e.g. several crenophilic species present in a small group of similar brooks. Other factors may be important, like stream morphology or distance between sites. The three torrents with similar species composition had large gravel beds and catchment areas limited to a few km², but differed in their exposition (Haller 2013). The only similar spring brooklets in the Val Spöl had similar morphology and were situated very close. Springs in the SNP are of various types due to very different chemical and physical characteristics (Schlüchter *et al.* 2013). An extensive study of 20 springs and their direct surroundings in the Park, taking into account the entire macroinvertebrate fauna, was conducted recently (Felder 2013). Neighbouring springs were shown to have similar faunistical assemblage in contrast to springs situated farther away.

The recently confirmed presence of *Perla grandis* in the SNP needs a focus on its altitudinal distribution in Switzerland. First evidence of its presence in the larval stage above 1800 m asl was provided in the 1990s (national database, CSCF, Neuchâtel). Proportionally, *P. grandis* has been found more often at altitudes \geq 1400 m asl after 1990 than before. The development of *P. grandis* larvae takes up to 3 years until emergence in spring, preferring streams with water temperatures from 8

Perla grandis (N = 589)




Fig. 4: Distribution (%) of *Perla grandis* records at the larval stage clustered by altitudinal classes (m asl) (national database, CSCF, Neuchâtel, October 2013).

to 20 °C, while maximum growth occurs during the warm season (Frutiger & Imhof 1997). As the species was not found before 2008 in the SNP despite the important collecting effort (Aubert 1965) and as grown larvae were found in 2012, we conclude that *P. grandis* colonized the Ova dal Fuorn. As water temperature has major influence on egg development duration and larval growth (Lillehammer et al. 1989), influence of temperature increase has to be considered. There is clear evidence for warming in the SNP since 1917, with a raise of annual air temperature of 1.6 °C per 100 years and a more pronounced seasonal air temperature increase of 1.7 to 5 °C (Bundesamt für Meteorologie und Klimatologie Meteoschweiz, 2013). Warming is higher in spring, during increase of larval growth that follows winter growth slowdown. As water temperature is related to air temperature (e.g. Mohseni & Stefan 1999), stream temperatures increased from 0.8 to 1 °C in lowland rivers (Hari et al. 2006). In small mountain streams, in absence of glacier melt-water as in the SNP, warming is also visible and mainly due to solar and long wave radiation, heat exchange with the sediment and dissipation of kinetic energy (Meier et al. 2003). Increase of water temperature in the SNP streams due to climate change is therefore highly probable and the possible impact on the freshwater fauna has to be documented further. Issues are for example the ability of species to maintain viable populations in the SNP and the consequences on cold stenotherm species living at high altitudes, which are expected to come under increasing physiological stress because they will not be able to migrate to higher elevations. The immigration of species from lower locations is expected to be a major driver of the freshwater biodiversity in the Alps (Burgherr et al. 2003; Robinson & Oertli 2009). Migration of eurytherm species to higher locations has already been observed as illustrated by the following examples. The eurytherm mayfly *Habrophlebia eldae*, has immigrated from Italy to the canton of Ticino (CH) (Wagner et al. 2007). North of the Alps, in the catchment area of the River Rhine, the stonefly Leuctra geniculata recently spread to higher locations (Vittoz et al. 2013).

The importance of long-term monitoring in freshwater systems, especially in protected areas, has been recently emphasized (Robinson et al. 2011). Biodiversity monitoring in Switzerland already includes Plecoptera species diversity on a systematic sampling grid covering the whole country (Altermatt et al. 2013). As some changes are only perceptible at the species level, the present SNP stonefly species data is of unique value since it allows long term comparisons with a historical data set over more than half a century time span. A SNP Plecoptera monitoring program would therefore benefit from both Swiss large scale program and the detailed SNP database (box 13.1 in Scheurer et al. 2014). Such a data base is also valuable for the assessment of climate change effects, the control monitoring after rehabilitation measures (Mürle & Ortlepp 2012) and other possible impacts. Alpine streams in Switzerland are particularly threatened by the country's much higher hydroelectric power production density (kWh/year and km²) than in other neighbouring countries (Kissling et al. 2007) and in fact they are coming under increasing pressure through the planned energy transition from nuclear to non-fossil. The actual data base on the SNP is an important basis for a future reference system for the Swiss freshwater typology (Schaffner et al. 2013).

ACKNOWLEDGMENTS

The project has been encouraged and financially supported by the SNP scientific commission; thanks to Thomas Scheurer as well as to Ruedi Haller from the SNP management who authorised the fieldwork. The SNP kindly prepared the map in Fig. 1. We are grateful to Uta Mürle and Johannes Ortlepp (Hydra, Öschelbronn, D) who collected stonefly adults with enthusiasm and kindly made their collected specimens and data available to us. The CSCF in Neuchâtel provided historical species data and altitude datasets. We thank Tom Gonser who helped with the English text. Thanks to Gilles Vinçon and Michel Sartori for their reviews and constructive inputs.

BIBLIOGRAPHY

- Altermatt, F., Seymor, M. & Martinez, N. 2013. River network properties shape α-diversity and community similarity patterns of aquatic insect communities across major drainage basins. Journal of Biogeography 40(12): 2249–2260.
- Aubert, J. 1965. Les Plécoptères du Parc National Suisse. Résultats des recherches scientifiques entreprises au Parc National Suisse 55: 223-271.
- Bundesamt für Meteorologie und Klimatologie Meteoschweiz 2013. Die Klimaentwicklung im SNP von den frühen Messungen bis zum Jahre 2099. *In*: Haller, H., Eisenhut, A., Haller, R. (eds), Atlas des Schweizerischen Nationalparks. Die ersten 100 Jahre, pp. 214-217. Nationalparkforschung Schweiz 99/1. Haupt, Bern.
- Burgherr, P., Klein, B., Robinson, C. & Tockner, K. 2003. Surface Zoobenthos. *In*: Ward, J.V. & Uehlinger, U. (eds), Ecology of a Glacial Flood Plain. pp. 153-174. Kluwer Academic Publishers, Dordrecht, Boston, London.
- Eisenhut, A. 2013. Die Gewässer im SNP. *In*: Haller, H., Eisenhut, A. & Haller, R. (eds), Atlas des Schweizerischen Nationalparks. Die ersten 100 Jahre, pp. 28-29. Nationalparkforschung Schweiz 99/1. Haupt, Bern.
- Felder, S. 2013. Faunistische Charakterisierung alpiner Quellen im Nationalpark. Masterarbeit, Universität Basel. 53 pp.
- Frutiger, A. & Imhof, A. 1997. Life cycle of *Dinocras cephalotes* and *Perla grandis* (Plecoptera: Perlidae) in different temperature regimes. *In*: Landolt, P. & Sartori, M. (eds), Ephemeroptera & Plecoptera. Biology Ecology Systematics. pp. 34-43. Mauron + Tinguely & Lachat SA, Fribourg, Switzerland
- Gonseth, Y., Wohlgemuth, T., Sansonnens, B. & Buttler, A. 2001. Die biogeographischen Regionen der Schweiz, Erläuterungen und Einteilungsstandard. Bundesamt für Umwelt, Umwelt Materialien 137, 48 pp.
- Graf, W., Lorenz, A.W., Tierno de Figueroa, J.M., Lücke, S., López-Rodríguez, M.J. & Davies, C. 2009. Distribution and ecological preferences of European freshwater organisms. Volume 2, Plecoptera Schmidt-Kloiber, A. & Hering, D. (eds). Pensoft Publishers, Sofia-Moscow, 262 pp.
- Haller, R. 2013. Norden oder Süden, flach oder steil. Grundsätzliches zur Exposition und Hangneigung. *In*: Haller, H., Eisenhut, A. & Haller, R. (eds), Atlas des Schweizerischen Nationalparks. Die ersten 100 Jahre, pp. 20-21. Nationalparkforschung Schweiz 99/1. Haupt, Bern.
- Hari, R.E., Livingstone, D.M., Siber, R., Burckhardt-Holm, P. & Güttinger, H. 206. Consequences of climatic change for water temperature and Brown trout populations in Alpine rivers and streams. Global Change Biolology 12:10-26.
- Hering, D. 1995. Die Wasserinsektenfauna einiger nordalpiner Flussauen (Insecta: Ephemeroptera, Plecoptera, Coleoptera, Trichoptera). Lauterbornia 22: 31-49.
- Indermühle, N. & Oertli, B. 2007. Mise en place d'un monitoring de la biodiversité des étangs du cirque de Macun (Parc National Suisse): les macroinvertébrés aquatiques. Nationalpark-Forschung in der Schweiz 94: 73-182.
- Jaccard, P. 1908. Nouvelles recherches sur la distribution florale. Bulletin de la Société Vaudoise des Sciences Naturelles XLIV, 163: 223-269.
- Kissling, M., Brosi, D. & Cadonau, G. 2007. Wasserkraftnutzung und Restwasser. Rechtsfragen, Vollzug und Entschädigungslösungen im Interesse der Fliessgewässer. Schweizerische Greina-Stiftung, 101 pp.
- Lillehammer, A., Brittain, J.E., Saltveit, S.J. & Nielsen, P.S. 1989. Egg development, nymphal growth and life cycle strategies in Plecoptera. Holarctic Ecology 12: 173-186.
- Lubini, V., Knispel, S. & Vinçon, G. 2012a. Les Plécoptères de Suisse. Identification et distribution.

 Fauna Helvetica 27, CSCF & SEG, Neuchâtel, 270 pp.

- Lubini, V., Knispel, S., Sartori, M., Vicentini, H. & Wagner, A. 2012b. Listes rouges Ephémères, Plécoptères, Trichoptères. Espèces menacées en Suisse, état 2010. OFEV, Bern & CSCF, Neuchâtel. L'environnement pratique n°1212, 111 pp.
- Malicky, H., Vinçon, G. & Delmastro, G.B. 2007. Nuovi dati sulla distribuzione di Plecotteri e Tricotteri sulle Alpi occidentali italiane (province di Cuneo e Torino) (Insecta: Plecoptera, Trichoptera). *In*: Delmastro, G.B., Gaggino, A., Giachino, P.M., Morisi, A. & Rastelli, M. (eds), Ricerche sugli ambienti acquatici del Po Cuneese Interreg IIIA Aqua Memorie del'Associazione Naturalistica Piemontese 8: 75-95.
- Meier, W., Bonjour, C., Wüest, A. & Reichert, P. 2003. Modelling the effect of water diversion on the temperature of mountain streams. Journal of environmental engineering 129: 755-764.
- Mohseni, O. & Stefan, H.G. 1999. Stream temperature/air temperature relationship: a physical interpretation. Journal of Hydrology 218: 128–141.
- Moog, O. (ed.) 1995. Fauna Aquatica Austriaca, Lieferung 1995. Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Wien.
- Moog, O. (ed.) 2002. Fauna Aquatica Austriaca, Lieferung 2002, Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien.
- Mürle, U. & Ortlepp, J. 2012. Fliessgewässermonitoring im Schweizerischen Nationalpark. Nationalpark-Forschung in der Schweiz 98: 29-54.
- Nadig, A. 1942. Hydrobiologische Untersuchungen in Quellen des Schweizerischen Nationalparkes im Engadin (unter besonderer Berücksichtigung der Insektenfauna). Ergebnisse der wissenschaftlichen Untersuchungen des schweizerischen Nationalparks 1: 267-432.
- OFEV 2011. Liste des espèces prioritaires au niveau national. Espèces prioritaires pour la conservation au niveau national, état 2010. Office fédéral de l'environnement, Berne. L'environnement pratique n°1103, 132 pp.
- Oswald, R. 1990. Anthropogene Einflüsse auf die Benthos-Invertebraten alpiner Fliessgewässer und deren Selektionseffizienz auf das Nahrungsspektrum von Bachforellen im Rhein (Kanton Graubünden, Schweiz). Dissertation Universität Zürich, 64 pp.
- Robinson, C.T. & Oertli, B. 2009. Long-term Biomonitoring of Alpine Waters in the Swiss National Park. Eco.mont. Journal of protected mountain areas research and management 1(1): 23-34.
- Robinson, C.T., Doering, M. & Seelen, L. 2011. Use of protected areas for freshwater biomonitoring case studies in Switzerland. Eco.mont. Journal of protected mountain areas research and management 3(2): 13-22.
- Schaffner, M., Pfaundler, M. & Göggel, W. 2013. Typologie des cours d'eau suisses. Une base pour l'évaluation et le développement des cours d'eau. Office fédéral de l'environnement, Berne. Connaissance de l'environnement n° 1329, 63 pp.
- Scheurer, T., Rehsteiner, U., Weibel, R. & Baur, B. 2014. Fazit und Ausblick nach 100 Jahren Nationalparkforschung. Nationalparkforschung Schweiz 100/I: 355-377. Haupt, Bern.
- Schlüchter, C., Clausen, M., Hämmig, C., Margreth, A., Pointner, E., Steiner, B., Strasky, S. & Vetter, H. 2013. Quellen. Natürliche Wasseraustritte im Gelände. *In*: Haller, H., Eisenhut, A. & Haller, R. (eds), Atlas des Schweizerischen Nationalparks. Die ersten 100 Jahre, pp. 30-31. Nationalparkforschung Schweiz 99/1. Haupt, Bern.
- Smukalla, R. & Friedrich, G. 1994. Ökologische Effizienz von Renaturierungsmassnahmen an Fliessgewässern. Landesamt Nordrhein Westfalen. Materialien Nr. 7, 462 pp.
- Stucki, P. 2012. Directives pour la conservation de matériel macrozoobenthos (MZB). Musée de zoologie de Lausanne-Aquabug-CSCF. Rapport, 4 pp.
- Vinçon, G. & Graf, W. 2011. Two new alpine *Leuctra* in the *L. braueri* species group (Plecoptera, Leuctridae). Illesia 7(9): 92-103.
- Vinçon, G. & Murányi, D. 2009. Revision of the *Rhabdiopteryx neglecta* species group (Plecoptera: Taeniopterygidae). Aquatic Insects, Supplement 1, 31: 203-218.
- Vittoz, P., Cherix, D., Gonseth, Y., Lubini, V., Maggini, R., Zbinden, N. & Zumbach, S. 2013. Climate change impacts on biodiversity in Switzerland: A review. Journal for Nature Conservation 21(3): 154-162.
- Wagner, A., Lubini, V. & Vicentini, H. 2007. *Habrophlebia eldae* Jacob & Sartori, 1984 (Ephemeroptera: Leptophlebiidae) neu für die Schweizer Fauna. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 80: 91-98.
- Zurwerra, A., Bur, M., Turcsanyi, B., Maier, K.J. & Tomka, I. 2000. Benthische Wirbellosenfauna des Sensesystems (Kt. Freiburg). Mitteilungen der Schweizerischen Entomologischen Gesellschaft 73: 115-142.

(received April 30, 2015; accepted May 9, 2015; published December 31, 2015)