

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 88 (2015)

Heft: 3-4

Artikel: First records of *Loborhynchapion amethystinum* (Miller, 1857) from
Switzerland, new biological insights and remarks on *Loborhynchapion*
obtusum (Desbrochers des Loges, 1866) (Coleoptera, Apionidae)

Autor: Giusto, Carlo / Germann, Christoph

DOI: <https://doi.org/10.5169/seals-583854>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

First records of *Loborhynchapion amethystinum* (Miller, 1857) from Switzerland, new biological insights and remarks on *Loborhynchapion obtusum* (Desbrochers des Loges, 1866) (Coleoptera, Apionidae)

CARLO GIUSTO¹ & CHRISTOPH GERMANN^{2,3}

¹ Via Milite Ignoto 46/9, I-16036 Recco (Genova), Italy; carlo.giusto.apion@gmail.com

² Naturmuseum Solothurn, Klosterplatz 2, CH-4500 Solothurn, Switzerland.

³ Naturhistorisches Museum der Burgergemeinde Bern, Bernastrasse 15, CH-3005 Bern, Switzerland; germann.christoph@gmail.com

The first records of *L. amethystinum* from the Cantons of Valais and Grisons in Switzerland are reported. However, the record of *L. obtusum* from Switzerland remains doubtful. Both species are illustrated and their morphology, ecology and ethology are compared. *Astragalus monspessulanus* is quoted for the first time as host plant of *L. amethystinum* and new data on the larval and pupal microhabitat are provided.

Keywords: Curculionoidea, distribution, ecology, ethology, adventive species.

INTRODUCTION

In his «Nachträge zur Fauna coleopterorum Helvetiae besonders aus dem Gebiete des berner Seelandes, des Jura und der Walliser Alpen», published in 1888, Rätzer quoted *Apion obtusum* Desbrochers des Loges, 1866 from the Val d'Entremont (Swiss Valais) among 14 listed apionid species. Later, the same record, also reported by Favre (1890) and Stierlin (1898), has been generically quoted from Switzerland by Alonso-Zarazaga (2011) and Pelletier (2014) who cited the same species as *Loborhynchapion obtusum*.


Because the examination of most collections from Switzerland, including those of August Rätzer (1845–1907) stored in the Naturhistorisches Museum der Burgergemeinde Bern, did not reveal any specimens of *L. obtusum*, Germann (2010) correctly included the species as doubtful in his checklist of the Swiss weevils.

The first recordings of *Loborhynchapion amethystinum* (Miller, 1857) in the Swiss Cantons of Valais and Grisons give us the opportunity to compare the morphology of the two species and to analyze their distribution, ecology and ethology.

MATERIAL AND METHODS

Measurements and acronyms

Abbreviations of measurements (in alphabetical order) are as follows: Bl = body length measured in dorsal view, from the base of rostrum to the apex of elytra, in a position in which they are at the same level; Le = length of elytra from anterior margin to apex, in dorsal view; Lp = length of prothorax from front margin to base along



Figs 1–4: 1–2: habitus of *Loborhynchapion amethystinum* from Binntal: Giesse Camping. 1) male, 2) female.— 3–4: habitus of *Loborhynchapion obtusum* from Cima Ciantiplagna. 3) male, 4) female. Scale bar: 1 mm.

midline, in dorsal view; Lr = length of rostrum from apex (excluding mandibles) to fore margin of eye, in lateral view; We = width of elytra at their widest point, in dorsal view; Wmsr = width of mesorostrum at the widest point, in dorsal view; Wp = width of prothorax at the widest point, in dorsal view. Mean values are given in square brackets.

In the text the following acronyms have been used (in alphabetical order): CG = Carlo Giusto's collection, Recco, Italy; NMBE = Naturhistorisches Museum der Burgergemeinde Bern, Switzerland.

RESULTS

***Loborhynchapion amethystinum* (Miller, 1857)**

(Figs 1, 2, 5–7)

Specimens examined: Switzerland: Canton of Valais – 1 ♂: 053_08.3, SCHWEIZ, VS, Binntal, Binn, Campingplatz, rechtes Ufer, (S-exp.), N 658.700-W 135.740, 1,500 m, 28.VI.2008, Ch. Germann leg., on *Astragalus monspessulanus* L., (NMBE); 6 ♂♂, 8 ♀♀, 1 larva, 1 ♀ pupa: Giesse (Camping) (VS) 46°22'15" N-08°12'09" E, 1,500–1,550 m, 4–5.VII.2015, Ch. Germann & C. Giusto leg., on *Astragalus monspessulanus* L. (CG); 2 ♂♂: 256_15.1 SCHWEIZ, VS, Binn, Giesse Camping, oberhalb, S-exp. Hang, [above, south exposed slope] 1,500 m, 4.7.2015, Ch. Germann leg. (NMBE); 1 ♀: 254_15.2 SCHWEIZ, VS, Binn, Giesse Camping, oberhalb, S-exp. Hang, 1,500 m, 9.6.2015, Ch. Germann leg. (NMBE); 2 ♂♂, 1 ♀: Binntal: env. of Binn (Gallery) (VS) 46°41'10" N-08°10'20" E, 1,360 m, 5.VII.2015, Ch. Germann & C. Giusto leg., on *Astragalus monspessulanus* L. (CG); 2 ♂♂, 3 ♀♀, 2 larvae, 2 ♀♀ pupae: 256_15.2 SCHWEIZ, VS, Binn, E Santigläis (bei Tunneneingang), 656.465 / 134.691, 1,350 m, 5.7.2015, Ch. Germann leg. (NMBE).

Canton of Grisons – 1 ♂: 012_06_10. SCHWEIZ, GR, Ftan, Umgb., N813.774 /E186.189, 1,700 m, beating *Astragalus glycyphyllos*, 19.6.2006, Ch. Germann leg. (NMBE).


Distribution: Sibero-European species distributed from central Europe to Russian Far East.

In particular, it is recorded from the following states: Switzerland (Canton of Valais, Canton of Grisons), Germany, Austria (Carinthia, Styria, Burgenland, Lower Austria, Vienna), Czech Republic (South Moravian Region), Slovakia (Bratislava Region, Trenčín Region, Košice Region), Hungary (Komárom-Esztergom, Pest, Budapest, Bács-Kiskur), Ukraine (Lviv Oblast, Chernivtsi Oblast, Vinnytsia Oblast), Romania (Sibiu), Bulgaria (Sofia City Province), Turkey (Central Anatolian Region), Syria, Iran, Azerbaijan (Nakhchivan Autonomous Republic), Uzbekistan (Bukhara), Kazakhstan, Russia (Southern Federal District, Volga Federal District, Urals Federal District, Siberian Federal District, Far Eastern Federal District), Mongolia, China (Xinjiang).

Remarks: In Europe—excluding the vague and unconfirmed indication of Hamburg for Germany (Seidlitz, 1891a, 1891b) that seems very unlikely—its presence is mostly localized, even if not common, in south-eastern Czech Republic, in western Slovakia and in eastern and south-eastern Austria. In view of this, the new records from Switzerland, especially those from the Canton of Valais, considerably extend the distribution area of this species westwards.

Host plants: Fabaceae: *Astragalus australis* (Linnaeus) Lamarck, *Astragalus austriacus* Jacquin, *Astragalus danicus* Retzius, *Astragalus glycyphyllos* Linnaeus, *Astragalus monspessulanus* Linnaeus, *Astragalus onobrychis* Linnaeus.

Remarks: Baudyš (1912), Lengerken (1941) and Scherf (1963, 1964) also cited *Trifolium pratense* Linnaeus as host plant of *L. amethystinum*, but, as already



Figs 5–10: 5–7: head and prothorax of *Loborhynchapion amethystinum* from Binntal: Giesse Camp- ing. 5, 7) male, 6) female.— 8–10: head and prothorax of *Loborhynchapion obtusum* from Cima Cian- tiplagna. 8, 10) male, 9) female.

pointed out by Wagner (1914), this record is wrong and must be attributed to *Ischnopteron* (*Chlorapion*) *virens* (Herbst, 1797).

Astragalus monspessulanus is here quoted for the first time as host plant of *L. amethystinum*.

Ecology: All the specimens from Valais (Binntal: Giesse Camping) have been collected on a south facing slope in a montane *Pinus sylvestris* forest (Fig. 11) whereas the twelve specimens found at the entrance of the tunnel, just out of Binn, have been collected on a south-western facing rocky slope surrounded by a montane *Pinus sylvestris* forest (Fig. 12).

These observations agree well with bibliographical data indicating xerothermic grassy steppes, forest meadows and alpine heaths on sandy or gravelly soils as principal ecological niches inhabited by this species.

L. amethystinum mostly prefers hilly areas up to 1,000 meters a.s.l., but, as shown by the Swiss specimens, it can also reach 1,700 meters. The unusual finding of two specimens from Erciyes Mountain (Turkey) (Lodos *et al.* 2003), one of which found under stone at 2,700 m a.s.l., is extremely interesting and indicates the ability of this species to reach high altitudes, even if, at the same time, the eccentricity of the collecting locality in respect to the main distribution raises doubts about its correct identification.

Ethology: The limited bibliographical information about its ethology indicates that adults of *L. amethystinum* live on the exposed part of the host plant (Wagner 1906, 1914, 1916, Dieckmann 1977) and that, according to Stejskal (2004), in Ječmeniště (South Moravian Region, Czech Republic), they are common and active on the host plant, *A. austriacus*, especially during the night. Larvae develop inside the seeds of the pods (Wagner 1906, 1914, 1916, Urban 1923, Hoffmann 1958, Osella 1973, Köstlin 1973, Dieckmann 1977).

Our ethological observations agree with those of the adults that we found on the exposed parts of *A. monspessulanus* where traces of imago feedings were evident (Fig. 13), but totally contrast with those relative to larvae and pupae. Indeed we observed the pre-imaginal stages in the stems of the same plant inside of larval and pupal cells (Figs 14–17). Despite the observation in vitro of more than 60 pods for about two months, no specimen has been reared from the seeds of *A. monspessulanus*.

Phenology: Adults stay on host plants from May to October.

***Loborhynchapion obtusum* (Desbrochers des Loges, 1866)**

(Figs 3, 4, 8–10)

Specimens examined: Italy: Piedmont – 1 ♂, 1 ♀: Cima Ciantiplagna (Torino), 2,750 m, 30.IX.1989, C. Giusto leg., sub *Oxytropis helvetica* Scheele (CG); 1 ♀: I–Piemonte: Valle di Susa: Cima Ciantiplagna, 2,600 m, 3.X.1982, M. Meregalli leg. (CG).

Distribution: W-Alpine species.

Known only from France (Savoie, Hautes-Alpes) and Italy (Piedmont).

Host plants: Fabaceae: *Oxytropis helvetica* Scheele (synonym of *Oxytropis gaudinii* Bunge).



Figs 11–12: habitats of *Loborhynchapion amethystinum*: 11) surroundings Giesse Camping, — 12) surroundings Binn: entrance of the tunnel.

Ecology: A species apparently restricted to cacuminal areas in western Alps from 2,000 m up to 2,750 m a.s.l. The pair of photographed specimens were collected on the south facing slope near the summit.

Ethology: The specimens from Cima Ciantiplagna were found under little stones partially covering the procumbent stems of *O. helvetica*. Until now, larva and pupa remain unknown.

Phenology: The few known specimens have been collected from August to October.

CONCLUSIONS

It is absolutely evident that our knowledge about the two aforementioned species is very unbalanced in favour of *L. amethystinum*. This is apparently due to the presence of *L. obtusum* in a restricted areal, to its monophagy and to the fact that this species seems confined to the cacuminal zone in the western Alps. *L. amethystinum*, on the contrary, is oligophagous and more euryoecious, and has been able to colonise an extended area in the Palaearctic Region. Apart from the ecological and the biological differences listed above, it is possible to recognize the two species as follows:

- Species, on average, larger (Bl ♂: 2.41–2.65 [2.55] mm, ♀: 2.66–2.93 [2.80] mm); body surface shiny; dorsal piliform scales thin and inconspicuous; male rostrum stout with mesorostral dilatation obtusely rounded, female prorostrum cylindrical (Lr ♂: 0.72–0.84 [0.79] mm, ♀: 0.90–1.03 [0.97] mm; Wmsr ♂: 0.24–0.27 [0.26] mm, ♀: 0.18–0.21 [0.20] mm; Lr/Wmsr ♂: 2.74–3.32 [3.08], ♀: 4.56–5.26 [4.88]; Bl/Lr ♂: 3.11–3.46 [3.24], ♀: 2.70–3.05 [2.86]); prothorax bell-shaped with superficial, thin and sparse punctation (Lp ♂: 0.55–0.63 [0.59] mm, ♀: 0.53–0.66 [0.62] mm; Wp ♂: 0.59–0.67 [0.63] mm, ♀: 0.57–0.71 [0.60] mm; Lp/Wp ♂: 0.90–0.98 [0.93], ♀: 0.91–0.99 [0.93]); elytra (Le ♂: 1.63–1.81 [1.74] mm, ♀: 1.81–2.03 [1.95] mm; We ♂: 1.01–1.14 [1.10] mm, ♀: 1.12–1.30 [1.24] mm; Le/We ♂: 1.44–1.68 [1.58], ♀: 1.52–1.67 [1.58]) *Loborhynchapion amethystinum* (Miller, 1857)
- Species, on average, smaller (Bl ♂: 2.36 mm, ♀: 2.48–2.53 mm); body surface moderately shiny, almost opaque; dorsal piliform scales clearly visible; male rostrum stout with mesorostrum obtusely dilatated, female prorostrum feebly tapering to apex (Lr ♂: 0.69 mm, ♀: 0.82–0.84 mm; Wmsr ♂: 0.24 mm, ♀: 0.18 mm; Lr/Wmsr ♂: 2.88, ♀: 4.56–4.67; Bl/Lr ♂: 3.42, ♀: 3.01–3.03); prothorax trapezoidal or feebly bell-shaped with superficial rather dense punctation (Lp ♂: 0.52 mm, ♀: 0.58–0.60 mm; Wp ♂: 0.55 mm, ♀: 0.60–0.65 mm; Lp/Wp ♂: 0.95, ♀: 0.92–0.97); elytra (Le ♂: 1.64 mm, ♀: 1.80 mm; We ♂: 1.02 mm, ♀: 1.16–1.18 mm; Le/We ♂: 1.61, ♀: 1.53–1.55).... *Loborhynchapion obtusum* (Desbrochers des Loges, 1866)

Considering that *O. helvetica* is recorded for the southern Swiss Valais, as shown in Info Flora (2015), the presence of *L. obtusum* in this area must be considered possible and probable. Therefore, in light of this, it is desirable that further field researches be carried out at high altitudes to verify this hypothesis and so, finally, confirm or refute Rätzer's quotation.

ACKNOWLEDGMENTS

We are grateful to all our colleagues who, in different ways, have contributed to the writing of this note, in particular, many thanks to Enzo Colonnelli (Rome), Giulio Gardini (Genoa) and Roberto Poggi (Museo civico di Storia naturale «Giacomo Doria», Genoa) for their critical reading of the manuscript, to Eva Sprecher (Naturhistorisches Museum Basel) for bibliographical researches, to Peter Sprick (Hannover) for his advice concerning records from Germany, to Stefano Zoia (Milan) for his endless advices about the realisation of the photos and to Mark I. Russell (Anglesey) for the linguistic correction of the text.



Figs 13–17: 13–14: *Astragalus monspessulanus* from Binntal: Giesse Camping: 13) leaf with traces of imago feedings of *Loborhynchapion amethystinum*, 14): infested stem. — 15–16: larva of *Loborhynchapion amethystinum* in its cell. — 17: pupa of *Loborhynchapion amethystinum* in its cell.

REFERENCES

- Alonso-Zarazaga, M.A. 2011. Apionidae. In: Löbl, I & Smetana, A. (eds), Catalogue of Palearctic Coleoptera. Vol. 7. Curculionoidea 1., pp. 77–83 + 148–176. — Apollo Books, Stenstrup.
- Baudyš, E. 1912. Tři nové kálky Apiony vyvolané. — Časopis české společnosti entomologické 9: 143–147.
- Dieckmann, L. 1977. Beiträge zur Insektenfauna der DDR: Coleoptera – Curculionidae (Apioninae). — Beiträge zur Entomologie 27(1): 7–143.
- Favre, E. 1890. Faune des Coléoptères du Valais et des régions limitrophes. Avec la collaboration du Dr. Edouard Bugnion. — Zurcher & Furrer, Zürich, xlvii + 448 pp.

- Germann, C. 2010. Die Rüsselkäfer (Coleoptera, Curculionoidea) der Schweiz – Checkliste mit Verbreitungsangaben nach biogeografischen Regionen. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 83: 41–118.
- Hoffmann, A. 1958. Faune de France, 62, Coléoptères Curculionides (Troisième Partie). — Lechevalier, Paris: 1209–1839.
- Info Flora 2015. Das nationale Daten- und Informationszentrum der Schweizer Flora. <https://www.infoflora.ch/it/flora/1162-oxytropis-helvetica.html#synonyms>. Accessed in October 2015.
- Köstlin, R. 1973. Beiträge zur Insekten-Faunistik Südwestdeutschlands. Die Gattung *Apion*, mit einem Anhang über *Apion*-Funde ausserhalb des eigentlichen Beobachtungsgebietes, soweit sie kontrolliert werden konnten. — Mitteilungen des Entomologischen Vereins Stuttgart 8: iii + 198 pp.
- Lengerken, H. von. 1941. Von Käfern erzeugte Pflanzengallen. — Entomologische Blätter 37(3): 121–159.
- Lodos, N., Önder, F., Pehlivan, E., Atalay, R., Erkin, E., Karsavuran, Y., Tezcan, S. & Aksoy, S. 2003. Faunistic studies on Curculionidae (Coleoptera) of Western Black Sea, Central Anatolia and Mediterranean regions of Turkey. — İzmir, 83 pp.
- Osella, G. 1973. Una nuova specie di *Apion* Herbst della fauna italiana e ridefinizione dell'*Apion obtusum* Desbrochers (Coleoptera Curculionidae). — Atti del Museo civico di Storia naturale di Trieste 28: 197–209.
- Pelletier, J. 2014. Apionidae Schoenherr, 1823. In: Tronquet, M. (coord.), Catalogue des Coléoptères de France, pp. 636–645. — Association roussillonnaise d'Entomologie, Perpignan.
- Rätzer, A. 1888. Nachträge zur Fauna coleopt. Helvetiae besonders aus dem Gebiete des berner Seelandes, des Jura und der Walliser Alpen. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 8[1888–1893](1): 20–42.
- Scherf, H. 1963. Die Brutpflanzen der Gattung *Apion* Hbst. — Entomologische Blätter 59(1): 39–45.
- Scherf, H. 1964. Die Entwicklungsstadien der mitteleuropäischen Curculioniden (Morphologie, Biologie, Ökologie). — Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 506: 1–335.
- Seidlitz, G. 1891a. Fauna Transsylvanica. Die Käfer (Coleoptera) Siebenbürgens. — Hartungsche Verlagsdruckerei, Königsberg, [10 unnumbered] + lvi + 914 pp., 1 pl.
- Seidlitz, G. 1891b. Fauna Baltica. Die Käfer (Coleoptera) der Deutschen Ostseeprovinzen Russlands. Zweite neu bearbeitete Auflage, Sechster Theil. — Hartungsche Verlagsdruckerei, Königsberg, [10 unnumbered] + xlvi + 818 pp., 1 pl.
- Stierlin, G. 1898. Fauna coleopterorum helvetica. Die Käfer-Fauna der Schweiz nach der analytischen Methode bearbeitet. II. Theil. — Bolli & Böcherer, Schaffhausen, xii + 662 pp.
- Stejskal, R. 2004. Contribution to the knowledge of beetles (Coleoptera) of dry grasslands – weevils (Curculionoidea) of Ječmeniště near Znojmo (southern Moravia). — Entomofauna Carpathica 16: 74–82.
- Urban, C. 1923. Die Nahrungspflanzen der Apionen. — Entomologische Blätter 19(4): 171–176.
- Wagner, H. 1906. Beiträge zur Kenntnis der Gattung *Apion* Herbst. II. — Münchener koleopterologische Zeitschrift 3: 13–34.
- Wagner, H. 1914. Berichtigende Notizen. — Entomologische Blätter 10: 144–145.
- Wagner, H. 1916. Familie Curculionidae. In: Schaufuss, C., Calwer's Käferbuch. Einführung in die Kenntnis der Käfer Europas. Sechste Auflage. Vol. 2, pp. 1030–1198. — E. Schweizerbart'sche Verlagsbuchhandlung Nägele & Dr. Sprösser, Stuttgart.

(received March 11, 2015; accepted after revision October 15, 2015; published December 31, 2015)

