**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

**Entomological Society** 

Herausgeber: Schweizerische Entomologische Gesellschaft

**Band:** 88 (2015)

**Heft:** 3-4

**Artikel:** Scaphisoma poussereaui sp. nov. from La Réunion (Coleoptera:

Staphylinidae: Scaphidiinae), a range extension of the S. tricolor

species group

Autor: Löbl, Ivan

**DOI:** https://doi.org/10.5169/seals-583859

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

## Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 23.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# MITTEILUNGEN DER SCHWEIZERISCHEN ENTOMOLOGISCHEN GESELLSCHAFT BULLETIN DE LA SOCIETE ENTOMOLOGIQUE SUISSE

88: 367–370, 2015 doi:10.5281/zenodo.34001

Scaphisoma poussereaui sp. nov. from La Réunion (Coleoptera: Staphylinidae: Scaphidiinae), a range extension of the S. tricolor species group

# Ivan Löbl

Muséum d'histoire naturelle, route de Malagnou 1, 1207 Genève, Switzerland; ivan.lobl@bluewin.ch

Scaphisoma poussereaui sp. nov. from La Réunion is described. This species is a member of the S. tricolor species group known so far only from Southeast Asia. It may be readily distinguished from other Mascarene congeners by the elytra with shortened sutural striae.

Keywords: Scaphisoma pousseraui sp. n., La Réunion, new species

#### INTRODUCTION

The mycophagous genus *Scaphisoma* Leach, 1815 is almost world-wide in distribution and with 636 species currently recognized as valid the most species-rich genus of the subfamily (Löbl, 1997 and subsequently published descriptions). Though the Malagasy and Afrotropical Scaphidiinae remain inadequately studied, the Mascarene taxa have been studied in comparatively more detail by Vilson (1944) and Löbl (1977). Eight species of *Scaphisoma* are known from La Réunion, two of them (*S. obliquemaculatum* Motschulsky, 1863 and *S. nigrofasciatum* Pic, 1915) are widely distributed Oriental species, *S. montanellum* Vinson is common to Mauritius and La Réunion, and five species are endemic to La Réunion (Löbl, 1977). Here I describe a new species of *Scaphisoma* from La Réunion collected by J. Poussereau (Dax, France). The new species belongs to the *S. tricolor* species group known from Southeast Asia (Löbl 1975) and is very distinctive from the remaining Mascarene congeners.

## MATERIAL AND METHODS

The body length is measured from the anterior pronotal margin to the inner apical angle of the elytra. The maximal length and width ratios of the antennomeres are given, measured on antennae mounted on slides. Characters of metanepisterna refer to their exposed parts. The abdominal ventrites are counted from the first visible one (i.e., the third morphological sternite). The statement about abdominal microsculpture does not refer to the intersegmental membranes. The sides of the aedeagi refer to their morphological sides, with the ostium situated dorsally, while it is in the resting position rotated 90°.

The labels of the types are reproduced verbatim, different labels under a specimen are separated by a slash. The holotype bears a printed red «HOLOTYPE» label, each paratype a printed yellow «paratype» label, and each a respective printed identification label with «det. I. Löbl, 2015».

The material is housed in the following collections:

CPJP Private collection of Jacques Poussereau, Dax, France MHNG Muséum d'histoire naturelle, Geneva, Switzerland MNHN Muséum national d'Histoire naturelle, Paris, France


**TAXONOMY** 

# Scaphisoma poussereaui sp. nov.

(Figs 1-3)

Holotype: ♂, La Réunion 974 Coll. J. Poussereau / Chemin de Centure Maison Boyer J.P. Souche Jamblon 19 02 2015 (MNHN).

Description. Male. Length 1.85–2.05 mm, width 1.27–1.46 mm. Head, mouthparts, antennae, pronotum, hypomera, and tarsi ochreous. Anterior two thirds of elytra dark brown to blackish, apical third of elytra yellowish. Mesoventrite and metaventrite with epimera and anepisterna reddish-brown. Coxae, femora, tibiae and abdomen light brown, slightly lighter than metaventrite. Antennae with relative length/width of antennomeres as: III 13/8: IV 50/9: V 73/9: VI 60/10: VII 65/12: VIII 56/9: IX 68/12: X 65/15: XI 68/14. Pronotum and elytra not microsculptured. Pronotum with lateral margins evenly rounded, lateral margin carinae not visible in dorsal view, lateral striae very densely and distinctly punctate, disc with very fine and fairly dense punctation. Hypomera not microsculptured, with punctation very fine, hardly visible at 25 times magnification. Exposed part of scutellum fairly large, about 0.05 mm long. Elytra with lateral margins evenly rounded, lateral carinae in dorsal view concealed near bases, exposed posterior basal fourth to third; lateral striae very densely and fairly coarsely punctate; apical margins rounded, indistinctly crenulated near inner angles, inner apical angles situated posterior level of outer apical angles; sutural margins not raised, sutural striae fairly shallow, starting posterior to level of scutellar tip, subparallel; adsutural areas flat, with single puncture row; discal punctation of elytra dense and coarse except on yellowish apical area, fairly irregular, with punctures distinctly delimited, puncture intervals mostly distinctly larger than puncture diameters; punctation on light apical area very fine. Hind wings appearing fully developed. Mesanepisterna not microsculptured. Mesepimera as interval to mesocoxae and about 3 times as long as wide. Metaventrite with strigulate microsculpture; punctation very fine, distinct punctures in impressed antecoxal rows and punctures bordering submesocoxal lines excepted; punctures also fairly coarse on small areas near inner margin of metacoxae. Middle part of metaventrite convex, shallowly impressed apically. Submesocoxal lines parallel, submesocoxal areas about 0.03 mm long, about as sixth of interval to metacoxae. Metanepisterna convex, impressed toward suture, below level of metaventrite along suture, narrowed anteriad, with margin mostly oblique, rounded near anterior angles. Protibiae and metatibiae straight, mesotibiae distinctly curved posterior between basal third and apex, about as thick as protibiae, thicker than metatibiae. Protarsomeres 1 strongly widened, about as wide as apices of protibiae. Protarsomeres 2 and 3 much narrower than protarsomere 1. Mesotarsomeres 1 strongly



Figs 1–3. Aedeagus of *Scaphisoma pousserreaui* sp. nov., in ventral view (1), in lateral view (2), median lobe, without parameres in dorsal view (3). Scale bar = 0.2 mm.

widened, wider than apices of tibiae. Mesotarsomere 2 slightly narrower than 1, mesotarsomeres 3 weakly widened. Abdomen with strigulate microsculpture, very finely punctate except on a narrow area posterior to the intercoxal process and at margins of submetaxocal line. Submetacoxal lines convex, submetacoxal areas 0.06 mm, almost as third of interval to apical margin of sternite 1. Sternite 6 flattened in middle, extended apically to form large, broadly rounded lobe. Aedeagus (Figs 1–3) 1.28 mm long. Basal bulb weakly sclerotized. Ventral branch of apical process of median lobe strongly inflexed lateroventrally, somewhat flattened, weakly narrowed toward tip in dorsal view. Parameres each with large and strongly sclerotized subbasal apophyses bearing membranous lobe. Internal sac with proximal, strongly

bent, large tooth joined to moderately sclerotized area bearing three rods; apical part of internal sac very densely and very finely denticulate.

Female. With most characters as in male, but: Pronotum slightly lighter than or as dark as anterior two thirds of elytra. Apical elytral margins almost oblique, with inner angles slightly prominent, tooth-like and very finely crenulated. Mesotibiae weakly bent, about as thick as metatibiae. All tarsomeres narrow. Sternite 6 not lobed.

*Etymology*. The new species is named in honour of its collector, Jacques Poussereau, Dax, France.

Comments. The species is a member of the S. tricolor group defined in Löbl (1975: 272) containing the following species: S. affectum Löbl, 2015, S. aspectum Löbl, 2015, S. borneense Pic, 1916, S. chujoi Löbl, 1982, S. hajeki Löbl, 2012, S. luctans Löbl, 1986, S. maculiger Löbl, 1875, S. renominatum Löbl, 1975, S. tarsale Löbl, 2015, S. testaceomaculatum (Pic, 1915) and S. tricolor Heller, 1917. Two additional species, S. dentipenne Löbl, 1971 and S. philippinense Oberthür, 1883, known only from females, may also be members of the group. The group was defined mainly by having conspicuously large aedeagi with moderately sclerotized basal bulb, usually bearing a ventral ridge supporting robust external muscles, the apical process of median lobe is bifid, strongly sclerotized and asymmetrical, the parameres are complex and the expanded membranous apex is supported by strongly sclerotized apophyses, and the internal sac is complex, often containing a curved proximal tooth. The species of the S. tricolor group have comparatively large bodies, long antennae, the metaventrite with antecoxal puncture rows, the abdominal microsculpture strigulate, the male sternite 6 with a large, broadly rounded lobe. Having shortened sutural striae of the elytra is another diagnostic feature for some species and in a number of examined undescribed Philippine species (the striae are elongate in S. affectum, S. aspectum, S. chujoi, S. hajeki, S. maculiger and S. testaceomaculatum). The apical margins of the elytra are sexually dimorphic for some species, and males may also differ from females by their pronotal coloration. Scaphisoma poussereaui is readily distinguished from all members of the group by its elytral coloration and the aedeagal characters, in particular by the shape of the ventral branch of the apical process inflexed strongly ventrolaterally. In addition to these characters, S. poussereaui may be easily distinguished from its Mascarene congeners by the elytra with short sutural striae.

# ACKNOWLEDGMENTS

My cordial thanks are due to Jacques Poussereau, Dax, France, who made his collections available for study and to Richard A.B. Leschen, Auckland, New Zealand, for comments on the manuscript.

## **REFERENCES**

Löbl, I. 1975. Beitrag zur Kenntnis der orientalischen Scaphisomini (Coleoptera, Scaphidiidae). — Mitteilungen der Schweizerischen entomologischen Gesellschaft 48:269–290.

Löbl, I. 1977. Les Scaphidiidae (Coleoptera) de l'île de la Réunion. — Nouvelle Revue d'Entomologie 7: 39–52.

Löbl, I. 1997. Catalogue of the Scaphidiinae (Coleoptera: Staphylinidae). — Instrumenta biodiversitatis 1: i–xii, 1–190.

Vinson, J. 1943. The Scaphidiidae of Mauritius. — Mauritius Institut Bulletin 2: 177–209.

(received September 15, 2015; accepted October 22, 2015; published December 31, 2015)