

Taxonomic notes on Indian species of Echthroplexis Förster (Hymenoptera : Encyrtidae), with descriptions of two new species

Autor(en): **Shafee, S. Adam / Fatma, Anis**

Objekttyp: **Article**

Zeitschrift: **Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society**

Band (Jahr): **57 (1984)**

Heft 4: **Festschrift Prof. P. Bovey**

PDF erstellt am: **20.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-402134>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

Taxonomic notes on Indian species of *Echthroplexis* Förster (Hymenoptera: Encyrtidae), with descriptions of two new species

S. ADAM SHAFEE & ANIS FATMA

Department of Zoology, Aligarh Muslim University, Aligarh, India

Echthroplexis FÖRSTER is accepted as valid genus. *Mashhoodiella* HAYAT is synonymised with *Echthroplexis* FÖRSTER. Other generic synonymy is quoted. Authors who synonymised the genera are cited in brackets after authors of genera. Six new combinations (*E. albiclavatus* [AGARWAL] comb. n., *E. albiscutellaris* [KHAN] comb. n., *E. echthromorpha* [HAYAT] comb. n., *E. ferrierei* [HAYAT *et al.*] comb. n., *E. indicus* [AGARWAL] comb. n., and *E. nigritus* [AGARWAL] comb. n.) are established. Two new species: *E. tumkurensis* sp. n. and *E. longipedicellus* sp. n. are described and illustrated. Types deposited in Zoological Museum, Aligarh Muslim University, Aligarh, India.

Genus *Echthroplexis* FÖRSTER

Echthroplexis FÖRSTER, 1856: 33.

Type-species: *Coenocercus puncticollis* THOMSON.

Homalotylus MAYR, 1876: 752 (DE SANTIS, 1963: 162).

Type-species: *Encyrtus flaminius* DALMAN.

Nobrimus THOMSON, 1876: 137 (DE SANTIS, 1963: 162).

Type-species: *Encyrtus flaminius* DALMAN.

Mendozaniella BRETHES, 1913: 97 (TIMBERLAKE, 1919: 134).

Type-species: *Mendozaniella mirabilis* BRETHES.

Hemaenasioidea GIRAULT, 1916: 307 (TIMBERLAKE, 1919: 134).

Type-species: *Hemaenasioidea oculata* GIRAULT.

Lepidaphycus BLANCHARD, 1936: 13 (BLANCHARD, 1941: 16).

Type-species: *Lepidaphycus bosqi* BLANCHARD.

Neoaenasioidea AGARWAL, 1966: 71 (HAYAT *et al.*, 1975: 64).

Type-species: *Neoaenasioidea indica* AGARWAL.

Mashhoodiella HAYAT, 1972: 209. Syn. n.

Type-species: *Mashhoodiella echthromorpha* HAYAT.

Comments: HAYAT (1972) separated his genus *Mashhoodiella* from *Homalotylus* MAYR and *Isodromus* HOWARD on the basis of length and separated condition of funicle segments and length of fore wing venation etc. which are of specific value. Therefore, *Mashhoodiella* is treated here as synonym of *Echthroplexis* FÖRSTER.

Echthroplexis albiclavatus (AGARWAL) Comb. n.

Neoaenasioidea albiclavatus AGARWAL, 1970: 27.

Homalotylus albiclavatus (AGARWAL); HAYAT *et al.*, 1975: 69.




Fig. 1: A-K. *Echthroplexis nigritus* (AGARWAL), ♀.

(A) Head, facial view; (B) Head, dorsal view; (C) Mandible; (D) Maxillary and labial palpi; (E) Antenna; (F) Pronotum; (G) Fore wing; (H) Part of fore wing venation; (I) Apex of abdomen, dorsal view; (J) Subgenital plate; (K) Part of external genitalia.

L-N. *Echthroplexis flaminius* (DALMAN) ♀.

(L) Antenna; (M) Subgenital plate; (N) Part of external genitalia.

Host: Coccinellid larvae predaceous on coccids, *Pseudococcus* sp., *Nipaecoccus vastator* (MASKELL), *Centrococcus insolitus* (GREEN).

Distribution: INDIA: Shencottah, Shoranur, Joginder Nagar, Lakhimpur kheri, Aligarh.

Echthroplexis albiscutellaris (KHAN) Comb. n.

Neoaenasioidea albiscutellaris KHAN, 1976: 181.

Host: Aphid

Distribution: INDIA: Aligarh.

Echthroplexis echthromorpha (HAYAT) Comb. n.

Mashhoodiella echthromorpha HAYAT, 1972: 210.

Host: *Ceroplastes* sp.

Distribution: INDIA: Aligarh.

Echthroplexis ferrierei (HAYAT *et al.*) Com. n.

Homalotylus ferrierei HAYAT *et al.*, 1975: 67.

Host: Coccinellid larvae on *Cerococcus* sp.

Distribution: INDIA: Nasik.

Echthroplexis flaminius (DALMAN) (Fig. 1 L-N)

Encyrtus flaminius DALMAN, 1820: 340.

Echthroplexis flaminius (DALMAN); DE SANTIS, 1963: 164.

Material examined. - 10 ♀, 2 ♂, INDIA: Andhra Pradesh, Guntur, 10.VIII.1968, ex coccinellid larvae predaceous on *Nipaecoccus vastator* (MASKELL) on *Solanum* sp. (S. ADAM SHAFEE).

Echthroplexis indicus (AGARWAL) Com. n. (Fig. 2 D-F)

Neoaenasioidea indica AGARWAL, 1966: 73.

Homalotylus indicus (AGARWAL); HAYAT *et al.* 1975: 67.

Host: Coccinellid larvae on *Rastrococcus iceryoides* (GREEN).

Distribution: INDIA: Shoranur, Aligarh.

Echthroplexis longipedicellus sp. n. (Fig. 2 B, C)

Female.

Head dark with bluish green reflections, slightly longer than wide in facial view; frontovertex slightly more than twice as long as wide; ocelli red, arranged in acute triangle, lateral ocellus close to eye margin and separated by four times its diameter from occipital margin; eyes silvery white; malar space as long as eye width; maxillary and labial palpi 4 and 3 - segmented respectively; mandibles tridentate. Antennae (Fig. 2 B) dark except sixth funicle segment and club white; scape cylindrical, six and a half times as long as wide; pedicel slightly more than three times as long as wide, as long as following two funicle segments together; funicle segments 1-6 gradually decreasing in length distad; first funicle segment slightly less than twice as long as wide, sixth as long as wide; club unsegmented, obliquely truncated at apex, as long as preceding three funicle segments together.

Thorax dark; mesoscutum with white and scutellum with dark small hairs. Forewings hyaline with a broad infuscated patch medially; marginal vein as long as wide, stigmal vein longer than marginal and postmarginal veins together; marginal fringe short. Legs dark except tibial spur and basal four tarsal segments white; middle tibial spur as long as basitarsus.

Abdomen dark, about as long as thorax; subgenital plate reaching apex of abdomen; ovipositor concealed, arising from basal one-third of abdominal venter; third valvulae short (Fig. 2 C).

Body length: 2.5 mm.




Fig. 2: A. *Echthroplexis tumkurensis* sp. n., ♀: Antenna.
 B, C. *Echthroplexis longipedicellus* sp. n., ♀.
 (B) Antenna; (C) Part of external genitalia.
 D-F. *Echthroplexis indicus* (AGARWAL), ♀.
 (D) Antenna; (E) First valvifer; (F) Second valvifer with third valvula.

Comments: In the key to species of *Homalotylus* MAYR given by HAYAT *et al.* (1975) *E. longipedicellus* sp. n. runs close to *E. flaminius* (DALMAN), from which it can be separated by its having antennae with sixth funicle segment white, pedicel more than three times as long as wide, funicle segments 1-5 distinctly longer than wide.

Holotype ♀, 4 ♀ Paratypes, INDIA: Uttar Pradesh, Aligarh, University campus, ex *Coccinellid* larvae feeding on *Aonidiella orientalis* (Newstead), on *Dalbergia sissoo*, 12.IX.1981 (ANIS FATMA).

Echthroplexis nigritus (AGARWAL) Comb. n. (Fig. 1 A-K)

Neoaenasioidae nigritus AGARWAL, 1970: 27.

Homalotylus nigritus (AGARWAL); HAYAT et al., 1975: 65.

Material examined. - 4♀, INDIA: Uttar Pradesh, Aligarh, University campus, ex Coccinellid larvae predaceous on Coccids, on *Dalbergia sissoo* 4.X.1983 (ANIS FATMA).

Echthroplexis terebratus HAYAT et al.

Echthroplexis terebratus HAYAT et al., 1975: 69.

Host: *Nipaecoccus vastator* (MASKELL).

Distribution: INDIA: Ranipet.

Echthroplexis tumkurensis sp. n. (Fig. 2 A)

Female.

Head reddish brown, slightly longer than wide in facial view; frontovertex about twice as long as wide; ocelli white, arranged in equilateral triangle, lateral ocellus separated by twice its diameter from occipital margin and close to inner orbital margin; antennae inserted below lower level of eyes, inter-antennal space as wide as width of frons between eyes at median ocellus; malar space as long as eye width. Antennae (Fig. 2 A) yellowish brown except club white; scape cylindrical, slightly more than five times as long as wide; pedicel slightly less than twice as long as wide, distinctly longer than first funicle segment; funicle segments 1-6 gradually increasing in width distad, first as long as wide, sixth slightly wider than long; club entire, obliquely truncated apically, as long as preceding three funicle segments together.

Thorax reddish brown; mesoscutum with complete parapsidal furrows meeting medially on scuto-scutellar suture; mesoscutum and scutellum with numerous small setae. Forewings hyaline except broad infuscated patch medially; marginal vein very short, postmarginal vein shorter than stigmal vein. Hind wings hyaline. Legs reddish brown except tibial spur and tarsal segments of middle legs white; mid tibial spur as long as basitarsus.

Abdomen reddish brown, slightly shorter than thorax; cerci present on apical one-third of abdominal dorsum; ovipositor hidden, arising from basal half of abdominal venter.

Body length: 1.6 mm.

Comments: The new species is more closely related to *Echthroplexis indicus* (AGARWAL), from which it can be separated by having unexerted ovipositor, legs uniformly reddish brown except tibial spur and tarsal segments of middle legs white.

Holotype ♀, INDIA: Tumkur, ex Coccinellid larvae feeding on *Coccus* sp., on *Mangifera indica*, 5.VII.1968 (S. ADAM SHAFEE).

ACKNOWLEDGEMENTS

The authors are deeply indebted to Prof. NAWAB H. KHAN, Chairman, Department of Zoology, for providing research facilities. One of us (A. FATMA) is thankful to U.G.C., New Delhi, for providing financial assistance.

LITERATURE

- AGARWAL, M. M. 1966. *Three undescribed genera and species of Encyrtidae (Hymenoptera: Chalcidoidea) parasitic on Coccids.* Proc. Indian Acad. Sci. (B) 63: 67-79.
- AGARWAL, M. M. 1970. *Some new Chalcidoid parasites recorded from Aligarh (India) (Hymenoptera: Encyrtidae).* Mushi 44: 25-29.
- BLANCHARD, E. E. 1936. *Apuntes sobre Chalcidoideos argentinos nuevos y conocidos.* Rev. Soc. Ent. Argent., Buenos Aires 8: 7-32.
- BLANCHARD, E. E. 1941. *Nuevos parásitos del bicho de cesto Oeceticus kirbyi, GUILD.* Rev. Soc. Ent. Argent. Buenos Aires 11: 3-21.
- BRETHES, J. 1913. *Himenopteros de la America meridional.* An. Mus. Nac. Hist. Nat. Buenos Aires 24: 35-165.
- DALMAN, J. W. 1820. *Forsok till uppstallning af insect-familjen Pteromalini, i Synnerhet med afseende pa dei Sverige funne arter.* K. Svenska Ventensk. Akad. Handl. 1820.
- DE SANTIS, L. 1963. *Encirtidos de la Republica Argentina.* An. Comm. Invest. Cient. Prov. BS. Aires. 4: 9-422.
- FÖRSTER, A. 1856. *Hymenopterologische Studien Heft 2, Chalcidiae und Proctotrupii.* Aachen. 1-152.
- GIRault, A. A. 1916. *New miscellaneous Chalcidoid Hymenoptera with notes on described species.* Ann. ent. Soc. Amer. 9: 291-308.
- HAYAT, M. 1972. *Descriptions of two new genera and species of Encyrtidae (Hymenoptera: Chalcidoidea), with notes on some described species.* Acta ent. bohemoslov. 69: 207-214.
- HAYAT, M., ALAM, S. M. & AGARWAL, M. M. 1975. *Taxonomic survey of Encyrtid parasites (Hymenoptera: Encyrtidae) in India.* Alig. Musl. Univ. Publ. (Zool. Ser.) Indian Ins. Typ. 9: 1-112.
- KHAN, M. Y. 1976. *New species of the genus Neoaenasioidea AGARWAL (Hymenoptera: Encyrtidae).* J. Bombay Nat. Hist. Soc. 73: 181-183.
- MAYR. 1876. *Die europäischen Encyrtiden.* Verh. Zool.-bot. Ges. Wien. 25: 675-778.
- THOMSON, C. G. 1876. *Hymenoptera Scandinaviae.* 4. London, 1-259.
- TIMBERLAKE, P. H. 1919. *Revision of the parasitic Chalcidoid flies of the genera Homalotylus MAYR and Isodromus HOWARD, with description of two closely related genera.* Proc. U. S. nat. Mus. 56: 133-194.

(received July 2, 1984)