

**Zeitschrift:** Alpine entomology : the journal of the Swiss Entomological Society  
**Herausgeber:** Swiss Entomological Society  
**Band:** 7 (2023)

**Artikel:** Planning insect survey in alpine ecosystems  
**Autor:** Pellet, Jérôme  
**DOI:** <https://doi.org/10.5169/seals-1053259>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Planning insect surveys in alpine ecosystems

Jérôme Pellet<sup>1,2</sup>

<sup>1</sup> *n+p wildlife ecology, Place St-François 6, CH-1003 Lausanne, Switzerland*

<sup>2</sup> *Department of Ecology and Evolution, University of Lausanne, Lausanne CH-1015, Switzerland*

<https://zoobank.org/032B0AE3-92C0-43C3-AE98-9EBB41AD434E>

Corresponding author: Jérôme Pellet (jerome.pellet@nplusp.ch)

Academic editor: Claudia Buser ♦ Received 12 August 2023 ♦ Accepted 25 October 2023 ♦ Published 30 October 2023

## Abstract

Most biological survey programs rely on multi-species inventories (e.g. birds, amphibians, butterflies, dragonflies). These programs usually rely on multiple visits during pre-defined time windows. The implicit goal of this popular approach is to maximize the observed species richness. Here, we present a novel method to optimize the timing of survey windows using a framework maximizing the detectable species pool. We present a proof of concept using 20 years of entomological records in Switzerland using butterflies, dragonflies, and grasshoppers. The general framework presented can potentially be applied to a wide range of biological survey schemes. It offers a new practical tool for adaptive entomological monitoring under climate change.

## Key Words

Lepidoptera, Odonata, Orthoptera, altitudinal levels, phenology, adaptive monitoring

## Introduction

Standing at the core of complex ecological food webs, insects provide insights into the health and stability of ecosystems. They are thus widely used as bioindicators at local, regional, and international scales (McGeogh 1998; Thomas 2005; Buckland and Johnston 2017; Chowdhury et al. 2023). By surveying and monitoring insect diversity, we gain a better understanding of the intricate relationships between species and their habitats across time, enabling us to develop environmentally sound conservation strategies and evaluate the efficiency of public policies (Yoccoz et al. 2001).

Many of the ongoing entomological survey programs aim at estimating species richness among taxa. Even though recent technologies (e.g. computer vision, acoustic monitoring, radar, and molecular methods) offer new perspectives (van Klink et al. 2022), visual encounters remain the most widespread approach. This is especially true for several popular taxa that are widely surveyed in alpine ecosystems, such as butterflies/day-flying moths, dragonflies/damselflies, and crickets/grasshoppers, all of

which can be readily identified or photographed in the field. Even though these taxa do not contain an overwhelming number of species compared to other taxa, surveying them remains a costly endeavor.

Entomological visual surveys are usually based on repeated visits across the activity period of the focal taxon. This is necessary because individual species fluctuate in abundance asynchronously during a year (the adult activity or flight periods of various species of insect typically only partly overlap within a focal taxon, see Pellet 2008). These multiple visits aim to maximize the chance of encountering all potentially present species. Monitoring schemes therefore very often rely on pre-defined time windows surveys that are assumed to maximize the observed species richness of the community under scrutiny.

Here, we present a novel approach to identify the best time windows for surveying alpine entomological communities by optimizing the encounter probabilities of every species with as few visits as possible. Using 20 years of observations for three popular taxa, we provide evidence-based, data-driven, guidance for alpine insect survey planning.

## Material and methods

We first extracted all observations of Lepidoptera (limited to butterflies and day-flying moths), Odonata (dragonflies and damselflies), and Orthoptera (crickets and grasshoppers) from info fauna, the Swiss biological records center ([www.infofauna.ch](http://www.infofauna.ch)) for the period spanning 2003–2022. The data was then organized into three matrices (one for each taxon) containing (i) the species name, (ii) the year the observation was made, (iii) the altitudinal levels of the observation, (iv) 52 columns corresponding to the weeks of the calendar year. These weekly columns were then filled with the total number of adult individuals of a given species that had been observed each year at a given altitudinal level.

Species detectability in a given week at a given altitudinal level was first assumed to follow  $P(X_{s,t}) \cong 1 - e^{-N_{s,t}}$ , where  $P(X_{s,t})$  is the probability of detecting species  $s$  during week  $t$  and  $N_{s,t}$  is the number of observations of species  $s$  during week  $t$ . That is, the more abundant a species is, the more likely it is that a single individual of that species will be observed. In short, we ended up with an expected number of species being potentially observed at every altitudinal level, week, and year.


Our optimization algorithm then worked through the following steps, iterating years and altitudinal levels,

finding - by exhaustion of all possibilities - the combination of survey weeks maximizing the sum of  $P(X_{s,t})$  (i.e. the number of species likely to be detected). For convenience, we tested 5 scenarios representing an increasing number of annual surveys (from 1 to 5). We then used this data to plot the best time windows - from a single week to a combination of 5 different weeks - that maximize the species richness likely observed by an observer.

The first draft of the introduction, discussion, and abstract of this paper has been adapted with PerplexityAI (2023). Prompts included the first version of the texts along with requests to (i) shorten paragraphs, (ii) improve clarity and (iii) correct any grammatical errors. All outputs from PerplexityAI (2023) were then reviewed and edited before being taken into consideration.

## Results

The optimized survey windows for 3 taxa and 3 altitudinal levels are described in Fig. 1. Each of the 9 sub-figures illustrates the best periods to maximize detectable species richness under 5 survey intensity scenarios (from a single annual survey to 5 annual surveys).



**Figure 1.** Optimal time windows to maximize potential species richness in entomological surveys for 3 taxa at 3 altitudinal levels assuming between 1 and 5 surveys each. The mean of the 2003–2022 period is represented with a white dot, the colored bars represent the standard deviation. A single survey aiming at maximizing the potential species richness of butterflies in the lowland (lower left sub-figure) would have to take place between weeks 26 and 30 of the year (first half of July). If two surveys are planned, then they should ideally take place on week 23 (early June  $\pm 1$  week) and on week 28 (mid-July  $\pm 2$  weeks).

For Odonata at the subalpine level (top middle sub-figure), a single visit should be made on the last week of July (the white dot representing the median best week). Depending on yearly variability, this best week can span anywhere between mid-July and the end of August. If two surveys are envisioned, then the first one should occur in mid-July and the second one in early August.

As expected, higher elevations translate into later survey windows, the amplitude of the shift being about 2 weeks between the lowland and the subalpine levels. Fig. 1 also shows, with little surprise, that Orthoptera tend to be more detectable later in the year than Odonata and Rhopalocera, the latter two groups having a larger spring/early summer species pool.

Running the algorithm for the 1983–2022 period (data not represented in Fig. 1) yielded valuable insights into changes in the timing of the optimal survey windows between the two 20-year periods. On average, all groups showed an advance of the best time windows of 0.9 weeks. That is, the best time windows moved about one week early between the two time periods. More specifically, Rhopalocera and Orthoptera showed a bigger advancement (1.1 week) than Odonata (0.6 week). The advancement of the timing was also larger at the subalpine level (1.6 weeks) than at lower elevations (0.9 and 1.2 week for the lowland and mountain levels respectively). There was, however, no significant change in the standard deviations of the best time windows for any taxon or altitudinal level.

## Discussion and conclusion

Insect surveys represent technically and logistically challenging operations that can prove costly (Field et al. 2007). In a world of limited financial resources, optimizing survey periods allows for a better balance of resources between monitoring investments and management actions, which constitute the final aim of most natural resources public policies (Field et al. 2007). This approach requires that the goals and scope of the surveys be explicitly formulated (Anderson 2001). In our case, we postulated, as in many ongoing programs, that maximizing observed species richness was the objective. Maximizing species cumulative detection probabilities across multiple surveys increases the chance of obtaining relevant species community data, as well as identifying species/habitat relationships or detecting trends in occupancy (Pollock et al. 2002; MacKenzie and Royle 2005; Mourguiart et al. 2020). This optimization approach focusing on both detection probability and estimates of occupancy has proven useful in other groups in the past (e.g. amphibians, Barata et al. 2017 or mammals, Baumgardt et al. 2019). In short, our approach potentially increases the return on investment for multiple species survey schemes.

By using a large 20-year-long dataset across multiple altitudinal levels, we closed the loop of active adaptive monitoring, where data collected in the past is used to

improve future efforts (Lindenmayer et al. 2011). This is especially important under climate changes that shift both habitat suitability and phenological periods of insects and other cold-blooded species (Vitasse et al. 2021; Buckley 2022). As we have shown here, the overall phenological shift in 20 years is about 1 week. It is consequently a necessity to regularly adapt existing survey programs (Halsch et al. 2021; Hill et al. 2021). Further optimization could consider not only changes in emergence timing but also changes in voltinism or shifts in altitudinal ranges.

## Acknowledgments

Gerard Maze and Marine Bugnon provided help with the construction of the model and its implementation in both Python (parsing and optimization algorithms) and R (summary statistics and figure production). DO! L'agence prepared the pictograms of Fig. 1. Yves Gonseth, Benedikt Schmidt, and Christian Monnerat provided feedback during the initial stages of this project.

## References

- Anderson DR (2001) The need to get the basics right in wildlife field studies. *Wildlife Society Bulletin* 29: 1294–1297.
- Barata IM, Griffiths RA, Ridout MS (2017) The power of monitoring: optimizing survey designs to detect occupancy changes in a rare amphibian population. *Nature Scientific Reports* 7: 16491. <https://doi.org/10.1038/s41598-017-16534-8>
- Baumgardt JA, Morrison ML, Brennan LA, Pierce BL, Campbell TA (2019) Development of multispecies, long-term monitoring programs for resource management. *Rangeland Ecology & Management* 172: 168–181. <https://doi.org/10.1016/j.rama.2018.07.010>
- Buckland ST, Johnston A (2017) Monitoring the biodiversity of regions: Key principles and possible pitfalls. *Biological Conservation* 214: 23–34. <https://doi.org/10.1016/j.biocon.2017.07.034>
- Buckley LB (2022) Temperature-sensitive development shapes insect phenological responses to climate change. *Current Opinion in Insect Science* 52: 100897. <https://doi.org/10.1016/j.cois.2022.100897>
- Chowdhury S, Dubey VK, Choudhury S, Das A, Jeengar D, Sujatha B, Kumar A, Kumar N, Semwal A, Kumar V (2023) Insects as bio-indicators: A hidden gem for environmental monitoring. *Frontiers in Environmental Science* 11: 1146052. <https://doi.org/10.3389/fenvs.2023.1146052>
- Field SA, O'Connor PJ, Tyre AJ, Possingham HP (2007) Making monitoring meaningful. *Austral Ecology* 32: 485–491. <https://doi.org/10.1111/j.1442-9993.2007.01715.x>
- Field SA, Tyre AJ, Possingham HP (2005) Optimizing allocation of monitoring effort under economic and observational constraints. *Journal of Wildlife Management* 69: 473–582. [https://doi.org/10.2193/0022-541X\(2005\)069\[0473:OAOMEU\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2005)069[0473:OAOMEU]2.0.CO;2)
- Hill GM, Kawahara AY, Daniels JC, Bateman CC, Scheffers BR (2021) Climate change effects on animal ecology: butterflies and moths as a case study. *Biological Reviews* 96: 2113–2126. <https://doi.org/10.1111/brv.12746>

- Halsch CA, Shapiro AM, Fordyce JA, Nice CC, Thorne JH, Waetjen DP, Forister ML (2021) Insects and recent climate change. *PNAS* 118(2): e2002543117. <https://doi.org/10.1073/pnas.2002543117>
- Lindenmayer DB, Likens GE (2010) The science and application of ecological monitoring. *Biological Conservation* 143: 1317–1328. <https://doi.org/10.1016/j.biocon.2010.02.013>
- Lindenmayer DB, Likens GE, Haywood A, Miezi L (2011) Adaptive monitoring in the real world: proof of concept. *Trends in Ecology and Evolution* 26: 641–646. <https://doi.org/10.1016/j.tree.2011.08.002>
- McGeogh MA (1998) The selection, testing, and application of terrestrial insects as bioindicators. *Biological Reviews* 73: 181–201. <https://doi.org/10.1111/j.1469-185X.1997.tb00029.x>
- MacKenzie DI, Royle JA (2005) Designing occupancy studies: general advice and allocating survey effort. *Journal of Applied Ecology* 42: 1105–1114. <https://doi.org/10.1111/j.1365-2664.2005.01098.x>
- Mourguiart B, Couturier T, Braud Y, Mansons J, Combrisson D, Besnard A (2020) Multi-species occupancy models: an effective and flexible framework for studies of insect communities. *Ecological Entomology* 46: 163–174. <https://doi.org/10.1111/een.12991>
- Pellet J (2008) Seasonal variation of butterflies' detectability in Pollard walks. *Journal of Insect Conservation* 12: 155–162. <https://doi.org/10.1007/s10841-007-9075-8>
- PerplexityAI (2023) Perplexity [Large language model]. <https://www.perplexity.ai>
- Pollock KH, Nichols JD, Simons TR, Farnsworth GL, Bailey LL, Sauer JR (2002) Large scale wildlife monitoring studies: statistical methods for design and analysis. *Environmetrics* 13: 105–119. <https://doi.org/10.1002/env.514>
- Possingham HP, Wintle BA, Fuller RA, Joseph LN (2012) The conservation return on investment from ecological monitoring. In: Lindenmayer DB, Gibbons P (Eds) *Biodiversity monitoring in Australia*. CSIRO, Collingwood, 49–62.
- Thomas JA (2005) Monitoring change in the abundance and distribution of insects using butterflies and other indicator groups. *Proceedings of the Royal Society B, Biological Sciences* 360: 339–357. <https://doi.org/10.1098/rstb.2004.1585>
- Vitasse Y, Ursenbacher S, Klein G, Bohnenstengel T, Chittaro Y, Delestrade A, Monnerat C, Rebetez M, Rixen C, Strebel N, Schmidt BR, Wipf S, Wohlgemuth T, Yoccoz NG, Lenoir J (2021) Phenological and elevational shifts of plants, animals and fungi under climate change in the European Alps. *Biological Reviews* 96: 1816–1835. <https://doi.org/10.1111/brv.12727>
- Yoccoz NG, Nichols JD, Boulinier T (2001) Monitoring of biological diversity in space and time. *Trends in Ecology & Evolution* 16: 446–453. [https://doi.org/10.1016/S0169-5347\(01\)02205-4](https://doi.org/10.1016/S0169-5347(01)02205-4)