

Zeitschrift: Alpine entomology : the journal of the Swiss Entomological Society
Herausgeber: Swiss Entomological Society
Band: 7 (2023)

Artikel: Description of the larva of Isoperla orobica Ravizza, 1975 (Plecoptera, Perlodidae), with notes on ecology and distribution in Switzerland
Autor: Lubini-Ferlin, Verena / Teufl, Pia / Novotny, Christoph
DOI: <https://doi.org/10.5169/seals-1053260>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Description of the larva of *Isoperla orobica* Ravizza, 1975 (Plecoptera, Perlodidae), with notes on ecology and distribution in Switzerland

Verena Lubini-Ferlin¹, Pia Teufl², Christoph Novotny², Laurent Vuataz^{3,4}

1 Eichhalde 14, CH-8053 Zürich, Switzerland

2 Institute of Hydrobiology and Aquatic Ecology Management, University of Natural Resources and Life Sciences, Gregor Mendel Strasse 33, A-1180 Vienna, Austria

3 Muséum cantonal des sciences naturelles, Département de zoologie, Palais de Rumine, Place de la Riponne 6, CH-1005 Lausanne, Switzerland

4 University of Lausanne (UNIL), Department of Ecology and Evolution, CH-1015 Lausanne, Switzerland

<https://zoobank.org/2922E5D9-5FC0-40E1-89BC-2AC742B70F42>

Corresponding author: Verena Lubini-Ferlin (lubini@sunrise.ch)

Academic editor: Dávid Murányi ♦ Received 5 July 2023 ♦ Accepted 21 October 2023 ♦ Published 2 November 2023

Abstract

The larva of *Isoperla orobica* Ravizza, 1975, an endemic Alpine species, is described with information about the distribution and ecology in Switzerland. New barcodes were generated to support the association between life stages. The species is associated with springs and spring brooklets. In Switzerland, the species occurs almost exclusively on the southern slopes of the Alps in the cantons Valais, Ticino and Grisons.

Key Words

Stoneflies, morphology, COI, Alps, endemism, taxonomy, larval description

Introduction

Isoperla Banks, 1906 is the most species-rich genus among the stonefly family Perlodidae, with at least 190 valid species recognized from the Holarctic and Oriental Realms (DeWalt et al. 2023). Sixty-three species are reported in Europe, with most of them in the Mediterranean region (DeWalt et al. 2023). Eight species are reported from Switzerland (Roesti 2021), belonging to five different species groups: *acicularis*: *Isoperla carbonaria* (Aubert, 1953), *I. lugens* (Klapálek, 1923); *grammatica*: *I. grammatica* (Poda, 1761), *I. felderorum* (Roesti, 2021); *obscura*: *I. obscura* (Zetterstedt, 1840); *oxylepis*: *I. orobica* (Ravizza, 1975), *I. oxylepis* (Despax, 1936); *rivulorum*: *I. rivulorum* (Pictet, 1841) (Muranyi 2011). *Isoperla lugens* and *I. orobica* are endemic species to the Alps (Graf et al. 2009). In Switzerland, all but two species (*I. orobica*, *I. felderorum*) are well known also in terms of their larval stage (Lubini et al. 2012). This study aims to newly describe the larva of *I. orobica*, which is

currently only known in the adult stage. This will enhance our knowledge of the distribution of *Isoperla* species in the context of the Swiss monitoring project of aquatic insects. Furthermore, the distribution and ecology of the species in the Alps are discussed.

Materials and methods

All specimens were caught by hand, the larvae by kicknet sampling or by picking specimens directly from the substrate, and the adults by sweep net or “Japanese umbrella” (beating sheet). The specimens were stored in 80% ethanol and deposited in the collection of the Muséum cantonal des sciences naturelles, Lausanne, Switzerland. Mouthparts, abdominal tergites and cerci were mounted on slides and photographed. To distinguish *I. orobica* from sympatric species such as *I. carbonaria* and *I. rivulorum*, we examined comparative material, referring to Lubini et al. (2012) regarding most morphological

characters. The terminology of the maxilla follows Sandberg (2011). Local and altitudinal distribution as well as the phenological data were provided by the National Center for Wildlife Data and Information in Switzerland (<https://infofauna.ch>), the Italian distribution by R. Fochetti (Tuscia University).

To provide additional evidence supporting the association between larvae and adults, we conducted genetic sequencing of a 658 bp fragment from the mitochondrial gene cytochrome oxidase subunit 1 (COI hereafter), which corresponds to the standard animal barcode region. We conducted this sequencing on three specimens of *I. orobica*, consisting of two larvae and one adult male (Table 1), using the non-destructive DNA extraction procedure described in Vuataz et al. (2011). The Polymerase Chain Reaction (PCR), purification and sequencing steps were conducted according to the methodology described in El Alami et al. (2022). Forward and reverse sequencing reads were assembled and edited in Codon Code Aligner 10.0.2 (Codon-Code Corporation, Dedham, MA), and aligned in Jalview 2.11.2.7 (Waterhouse et al. 2009). To tentatively augment our dataset, we searched for COI sequences associated with *I. orobica* in the GenBank and Barcode of Life Data System (BOLD) databases as of September 21, 2023, but no additional sequences were available.

Material examined

Isoperla orobica: Switzerland: 6 larvae, spring brooklet, tributary to river Taferna (VS), 46°15'53"N, 8°02'09"E, 1580 m, 6.VII.2017, leg. Verena Lubini, 1 larva DNA, GBIFCH00658280; 8 larvae, same locality, same date, leg. Remo Wüthrich; 4 larvae, same locality, 7.VII.2022, leg. Verena Lubini; 4 larvae, tributary to Fiume Bello, Alpe Rompiago (TI), 46°05'25"N, 8°59'44"E, 1131 m, 28.VI.2018, leg. Verena Lubini; 4 larvae, tributary to Riale Vadina, Alpe di Neggia (TI), 46°06'46"N, 8°50'55"E, 1289 m, 27.VI.2018, leg. Remo Wüthrich; 1 larva, 3 ♂, tributary to Riale Vadina, Rotiscia (TI), 46°07'05"N, 8°51'01"E, 996 m, 27.VI.2018, leg. Remo Wüthrich, 1 ♂ DNA, GBIFCH00658286; 6 larvae, same locality, 29.VI.2022, leg. Verena Lubini and Viviane Uhlmann; 1 larva DNA, GBIFCH00658284, tributary to Riale Vadina, Alpe di Neggia (TI), 46°06'47"N, 8°51'11"E, 1250 m, 29.VI.2022, leg. Verena Lubini; spring brooklet, tributary to river Laggina, Alte Staful (VS), 46°09'30"N, 8°03'24"E, 1550 m, 7.VII.2017, leg. Remo Wüthrich. Italy: 1 larva, 1 ♂, spring brooklet, Valchiusella, 45°32'4"N, 7°40'48"E, 1200 m, 6.VIII.2016, leg. Verena Lubini.

Table 1. Newly sequenced *Isoperla orobica* specimens from Switzerland, including collection details and GenBank accession numbers.

Specimen catalogue nb	Stage	GPS Coordinates	Altitude	Date	Collector	GenBank ID
GBIFCH00658280	Larva	46°15'53"N, 8°02'09"E	1580 m	6.VII.2017	V. Lubini	OR600533
GBIFCH00658284	Larva	46°06'47"N, 8°51'11"E	1250 m	29.VI.2022	V. Lubini	OR600534
GBIFCH00658286	Adult male	46°07'05"N, 8°51'01"E	996 m	27.VI.2018	R. Wüthrich	OR600535

Isoperla carbonaria: Switzerland: 3 larvae, Val Giona, Büee (TI), 46°06'18"N, 8°50'18"E, 1135m, 5.VI.2000, leg. Verena Lubini; 1 larva, Val Muggio (TI), 45°53'53"N, 9°02'32"E, 544 m, 5.VI.2000, leg. Verena Lubini; 1 larva, Valle Verzasca, Gerra (TI), 46°19'37"N, 8°48'08"E, 819 m, 14.IV.2023, leg. Paola Gandolfi.

Isoperla rivulorum: Switzerland: 1 larva, Alp Prüma, Val Roseg (GR), 46°27'29"N, 9°52'52"E, 1913 m, 27.III.2000, leg. Verena Lubini; 1 larva, spring brooklet, tributary to Rigia, Gruonholz (SZ), 47°02'25"N, 8°30'22"E, 1176 m, 19.VI.2022, leg. Emil Birnstiel; 1 larva, Munt San Franzesch (GR), 46°22'45"N, 10°06'03"E, 2150 m, 27.IV.2001, leg. Verena Lubini; 1 larva, river Landquart, Grüschi (GR), 46°58'27"N, 9°38'02"E, 593 m, 30.III.1993, leg. Verena Lubini.

Results

The alignment of the three COI sequences was 100% complete, with no missing data, and all sequences were identical, except for a single site that was ambiguous in the GBIFCH00658280 sequence (Table 1). These findings strongly support the conspecificity of the sequenced larvae and adult. As of September 21, 2023, a BLAST search on our sequences showed the closest match to be a sequence with a 94.1% similarity, originating from a specimen collected in Croatia and identified as *I. rivulorum* (Hlebec et al. 2021), while all other results were less than 94% similar, confirming that our sequences are the first *I. orobica* barcodes to be publicly deposited.

Description of the mature larva

Body length 8.9–11.3 mm (n = 14). General colour light brown, abdomen notably darker than head and thorax (Fig. 1A, B, 6C), ventral coloration pale (Fig. 1D, E). Pilosity typical of the genus, thoracic segments covered by dark brown clothing hairs and blunt marginal spines.

Head light brown with indistinct pale markings; defined pale interocellar area lacking; frontoclypeus anterior to M-line pale; sometimes two darker sublateral markings at anterior margin of frontoclypeus; pale sublateral occipital spots posterior to epicranial suture; head trapezoid shaped with frontal area slightly elongated (Fig. 2A). Scape, pedicel and following antennomeres pale; palpi and mouthparts pale (Fig. 2B).

Lacinia bidentate, sub-quadrata shaped with only slightly rounded inner margin; subapical tooth slightly longer than half of the length of apical tooth; 1 group A submar-




Figure 1. *Isoperla orobica*, larval morphology. **A.** Habitus dorsal; **B.** Abdomen dorsal; **C.** Tergite 1–4; **D.** Abdomen ventral; **E.** Furcasternites; **F.** Pronotum; **G.** Left cercus lateral; **H.** Right cercus detail lateral. (Photos: Pia Teufl, Christoph Novotny).



Figure 2. *Isoperla orobica*, larval morphology. A. Head and pronotum dorsal; B. Head ventral; C. Left lacinia and galea dorsal view; D. Right lacinia dorsal view; E. Right mandible dorsal view; F. Right mandible ventral view (Photos: Pia Teufl, Christoph Novotny).

ginal setae present, below apical tooth and adjacent to subapical tooth; 1 thin marginal seta below apical tooth and 1 thin marginal seta proximal to subapical tooth; 6–10 stout marginal setae, space between setae continuously decreasing towards base, almost all stout marginal setae of equal length except for the most proximal setae, which are shorter; sparse row of thin marginal setae (9–12) along proximal half of inner lacinial margin (Fig. 2C, D); scattered, sparse ventral surface setae on inner basal palm. Galea with four apical setae; inner and outer margin with sparse setae; galea reaching base of apical tooth. Mandible with one incisor and three molars, molar brush dense and long; dense setae dorsally and ventrally at inner margin; small area of setae laterally at the base of mandibles (Fig. 2E, F).

Pronotum light brown, lateral margins of disc lighter in colour; anterior and posterior margin of pronotum slightly darker; half as long as wide (ratio 1:2); spines on protonal margin distinct; longest spines as long as one fifth of the protonal width with shorter spines in between (Fig. 1F).

Mesonotum and metanotum with indistinct darker markings; wingpads pale; outer margin of wingpads slightly concave (Fig. 1A). Thoracic sternites pale; furcasternites and furcal pits hardly contrasting (Fig. 1E).

Abdominal terga mostly brown with variable dark median marking; first tergite lighter; posterior margin

of tergites 1–10 with long blunt spines, longest spines longer than half of tergal lengths (Fig. 1C). Sternites pale, distal segments slightly darker.

Legs light brown; dense row of acute spines covering dorsal margin of femur; spines short on proximal third and longer (length: half of femur width) on two distal thirds of femur; long silky setal fringe (almost as long as femur width) covering dorsal margin of femur (Fig. 3B); spines and setae on surface of femur (Fig. 3A).

Double row of dense spines (length: one third of tibia width) on dorsal margin of tibia (Fig. 3C) with longer ones towards distal third; dorsal side of tibia with sparse and long setal fringe (length: as long as tibia width). Tarsi with sparse setal fringe on dorsal margin (Fig. 3D). All claws with one basal tooth each.

Cercus pale yellow with a thin brown ring on proximal 1/4 of cercomeres; cercomeres cylindrical; medial cercomeres twice as long as wide. Cercal apical fringes dense, circumferential setae on medial cercomeres approximately one fourth of segment length, dorso-medial and ventromedial setae longer (length: more than half of segment length); cercus without dorsal setal fringe (Fig. 1G, H). Paraproct blunt with widely rounded apex.




Figure 3. *Isoperla orobica*, larval morphology. **A.** Front left femur; **B.** Front left femur, dorsal margin; **C.** Front left tibia; **D.** Front left tarsus (Photos: Pia Teufl, Christoph Novotny).

Distribution and ecology

Isoperla orobica is an endemic rheophilic species from the western and central Alps (Graf et al. 2009; Fochetti 2020), distributed only in northern Italy (Consiglio 1979; Ravizza 2023) and Switzerland (Fig. 4). In Switzerland, the species is predominantly observed along the southern slopes of the Alps, with its current distribution primarily centered on the Simplon Pass, as well as in certain mountainous regions within the canton Ticino (Val Colla, Valle di Vira, Val Serdina) and the canton Grisons (Val Mesocco) (<https://infofauna.ch>). Apart from the southern slopes of the Alps there is only one location in the central Alps, which is situated in the catchment of the river Rhone in the canton Valais, Switzerland.

The larvae were found in springs, spring brooklets and in brooks of the epirhithral zone mainly between 900 m and 1500 m, rarely above 2000 m. The substrate varies from organic to coarse stony bottoms in moderate to steep slopes (Fig. 5). The flight periods of the adult insects extend from late June to October at altitudes ranging from 900 m to 2030 m. The accompanying fauna consists of *Dictyogenus fontium* (Ris, 1896), *Chloroperla susemicheli* (Zwick, 1967), *Siphonoperla italicica* (Au-



Figure 4. Present distribution of *I. orobica* in Switzerland and Italy. Only one location is in the drainage area of the river Rhone (red star), all other locations (red squares) are in the drainage basin of the river Po. The Swiss locations represent areas of 5×5 km with a total of 30 sampling stations (<https://infofauna.ch>).

bert, 1953), *Leuctra insubrica* (Aubert, 1949), *L. armata* (Kempny, 1899), *L. braueri* (Kempny, 1898), *L. caprai* (Festa, 1939), *L. dolasilla* (Consiglio, 1955), *L. schmidti*




Figure 5. Habitat of *I. orobica* in Switzerland, upstream views. **A, B.** Spring brook, tributary to river Taferna (Rhone basin, canton Valais); **C.** River Vadina, tributary to Lake Maggiore (canton Ticino); **D.** Brook in Simplon pass, near Stalde (canton Valais) (Photos: Verena Lubini).

(Aubert, 1946), *L. vinconi* (Ravizza & Ravizza Dematteis, 1993), *Nemoura sinuata* (Ris, 1902) and *Protonemura lateralis* (Pictet, 1836).

Diagnosis

For comparison between *I. orobica* and the two sympatric species *I. rivulorum* and *I. carbonaria*, body coloration patterns and characteristics of the maxillae were considered (Table 2). In terms of coloration, *I. orobica* differs from both other species (Fig. 6, Lubini et al. 2012)

in having a uniformly brown colour and lacking a pale interocellar area on the head.

The lacinia of all three species is bidentate and generally sub-quadrata, but differences in the number of thin marginal setae on the lacinia and pilosity of the galea can aid in distinguishing these taxa. *Isoperla orobica* has 9–12 thin marginal setae compared to 25–35 in *I. rivulorum* (Table 2, Fig. 7). Additionally, *I. orobica* has setae on both the inner and outer margins of the galea; whereas *I. carbonaria* only has setae on the outer margin. Furthermore, *I. orobica* lacks the dense setal fringes on the tibiae, as is the case in the other two species.

Table 2. Verification table of morphological characters for *Isoperla orobica*, *I. rivulorum* and *I. carbonaria* (SMS = stout marginal setae; TMS = thin marginal setae).

Species	Body color	Interocellar area	Lacinia SMS	Lacinia TMS	Galea setae
<i>Isoperla orobica</i>	uniform	indistinct	6–10	9–12	inner & outer
<i>Isoperla rivulorum</i>	contrasting	distinct	7–8	25–35	inner & outer
<i>Isoperla carbonaria</i>	contrasting	distinct	7–8	10–16	outer



Figure 6. Larval habitus. **A.** *I. carbonaria*; **B.** *I. rivulorum*; **C.** *I. orobica* (Photos: Verena Lubini).




Figure 7. Lacinial morphology. **A.** *I. carbonaria*; **B.** *I. rivulorum* (Photos: Laurent Vuataz).

Acknowledgments

We are grateful for the photos of Figures 1 to 3 provided by the Institute of Hydrobiology and Aquatic Ecology Management, University of Natural Resources and Life Sciences, Vienna. Remo Wüthrich and Viviane Uhlmann helped collect larvae and adults, and Céline Stofel conducted the laboratory work for COI sequencing. The National Center for Wildlife Data and Information in Switzerland (<https://infofauna.ch>) provided the Swiss data, Romolo Fochetti the Italian data. We further thank Wolfram Graf for expert advice. Finally, we express our sincere gratitude to the reviewer, Chris Verdome, for his valuable comments which helped to improve our manuscript.

References

- Consiglio C (1979) La distribuzione dei Plecotteri italiani. Biogeographia – The Journal of Integrative Biogeography 6(1): 383–393. <https://doi.org/10.21426/B66110071>
- DeWalt RE, Maehr MD, Hopkins H, Neu-Becker U, Stueber G (2023) Plecoptera Species File Online. Version 5.0/5.0. <http://PlecopteraSpeciesFile.org/>
- El Alami M, Benlasri M, Sartori M, Vuataz L, Ghamizi M (2022) A new species of the genus *Prosopistoma* Latreille, 1833 (Ephemeroptera, Prosopistomatidae) from Morocco. ZooKeys 1117: 203–218. <https://doi.org/10.3897/zookeys.1117.83539>
- Fochetti R (2020) Endemism in the Italian stonefly-fauna (Plecoptera). Zootaxa 4722(4): 381–388. <https://doi.org/10.11646/zootaxa.4722.4.7>
- Graf W, Lorenz AW, Tierro de Figueroa JM, Lücke S, López-Rodrígues MJ, Davies C (2009) Distribution and Ecological Preferences of European Freshwater Organisms. Vol. 2. Plecoptera. Pensoft Publishers. Sofia, Bulgaria, 262 pp.
- Hlebec D, Sivec I, Podnar M, Skejo J, Kučinić M (2021) Morphological and molecular characterisation of the Popijač's Yellow Sally, *Isoperla popijaci* sp. nov., a new stenoendemic stonefly species from Croatia (Plecoptera, Perlodidae). ZooKeys 1078: 85–106. <https://doi.org/10.3897/zookeys.1078.66382>
- Lubini V, Knispel K, Vinçon G (2012) Die Steinfliegen der Schweiz. Bestimmung und Verbreitung. Les plécoptères de Suisse. Identification et distribution. Fauna Helvetica 27. CSCF & SEG, Neuchâtel, 270 pp.
- Muranyi D (2011) Balkanian species of the genus *Isoperla* Banks, 1906 (Plecoptera: Perlodidae). Zootaxa 3049(1): 1–46.
- Ravizza C (2023) The Stonefly communities of three brooks of the Soana Valley (Graian Alps), (Insecta Plecoptera). Rivista piemontese di Storia naturale 44: 87–101.
- Roesti C (2021) Die Steinfliegen der Schweiz. Haupt Verlag, Bern, 631 pp.
- Sandberg JB (2011) The *Isoperla* of California (Plecoptera: Perlodidae); larval descriptions and a key to 17 western Nearctic species. Illesia 7(22): 202–258.
- Vuataz L, Sartori M, Wagner A, Monaghan MT (2011) Toward a DNA Taxonomy of Alpine *Rhithrogena* (Ephemeroptera: Heptageniidae) Using a Mixed Yule-Coalescent Analysis of Mitochondrial and Nuclear DNA. PLoS ONE 6(5): e19728. <https://doi.org/10.1371/journal.pone.0019728>
- Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ (2009) Jalview Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics 25(9): 1189–1191. <https://doi.org/10.1093/bioinformatics/btp033>