

Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	49 (1969)
Heft:	2
Artikel:	The stable association enstatite-forsterite-chlorite in amphibolite facies ultramafics of the Lepontine Alps
Autor:	Trommsdorff, Volkmar / Evans, Bernard W.
DOI:	https://doi.org/10.5169/seals-38593

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Stable Association Enstatite-Forsterite-Chlorite in Amphibolite Facies Ultramafics of the Lepontine Alps

By *Volkmar Trommsdorff* (Basel) *) and *Bernard W. Evans* (Berkeley, Cal.) **)

With 2 text figures

Abstract

Ultramafic rocks with the paragenesis forsterite-enstatite-chlorite-magnetite occur in the sillimanite and calcite-anorthite zones of the Lepontine Alps. Textural and chemical evidence indicates that this assemblage formed during the Alpine metamorphism. Some oxidation prior to or during the last recrystallization caused the silicates to be very magnesian (e. g. olivine $Fa_{4.5}$). The paragenesis formed under conditions of low water activity in the amphibolite facies.

Geological Outline

Ultramafic rocks are an important constituent of the metamorphic series of the Lepontine Alps. They have given rise to numerous studies and to various interpretations reflecting the change in geological thought during the past sixty years (GRUBENMANN 1908; HEZNER 1909; KÜNDIG 1926; P. NIGGLI 1936; DAL VESCO 1953; KNUP 1958; MÖCKEL 1968). The Lepontine ultramafics are frequently associated with metamorphic Mesozoic carbonate sediments and form lenses and bodies from a few meters to several kilometers in size. They generally show a marked schistosity and lineation completely in agreement with the surrounding marbles, schists and gneisses – a fact already well understood by GRUBENMANN (1908). Together with the accompanying rocks, the ultramafics were subject to strong Alpine deformation and to out-lasting crystallization or recrystallization of various magnesian silicates, such as talc, anthophyllite, forsterite, and, in the zones of highest grade, enstatite.

*) Mineralogisch-Petrographisches Institut, Bernoullianum, 4000 Basel.

**) Department of Geology and Geophysics, University of California, Berkeley, California 94720, USA.

Enstatite-Forsterite-Schists of Val Cama

A typical example of enstatite-forsterite schist occur in Val Cama, Grisons (Eastern Lepontine Alps), in an area mapped and described by BRUGGMANN (1965). The grade of metamorphism in Val Cama is given by the occurrence of the parageneses anorthite-calcite (SCHWANDER and WENK 1967) and diopside-calcite-quartz in the adjacent marbles, and by the occurrence of predominant kyanite and locally of fibrolitic sillimanite in pelitic schists. Currently available experimental and calculated data on mineral equilibria indicate that the maximum temperature of metamorphism may have just exceeded 600° C. An estimate of pressure is much more uncertain – a value around four kb seems reasonable and not conflicting with estimates for the Western Bergell Alps (TROMMSDORFF 1968).

The ultramafic body at Val Cama forms a subvertical sheet about 100 m in thickness, accompanied on both sides by concordant marbles and calc-silicate rocks five to twenty meters in thickness. It outcrops over a length of more than two kilometers. Locally the ultramafics are inhomogeneous and



Fig. 1. Magnetite-chlorite-enstatite-forsterite schist, Val Cama. Specimen Mg 7. Magnification $\times 48$.

rich in hydrous phases such as talc, anthophyllite and tremolite, and these often occur in zoned veins cross-cutting the body. However, most of the ultramafic sheet is composed of magnetite-chlorite-enstatite-forsterite schist.

Petrography and Structural Analysis

One specimen of magnetite-chlorite-enstatite-forsterite schist, from Pro-magn, Val Cama, Switzerland (Coord. [Swiss net] 736.1/125.5; specimen Mg 7e), was studied in detail. The rock consists of a fine-grained, dark green matrix and of enstatite porphyroblasts forming slender prisms several cm in length oriented parallel to the foliation. In the matrix the dominant constituent is forsterite, forming a perfect mosaic of slightly elongated grains 0.5 to 1 mm in cross-section (Fig. 1). The marked schistosity of the rock is accentuated by evenly distributed colorless flakes of chlorite, strongly oriented parallel to the s-plane. Magnetite forms (up to 5 mm) xenomorphic and small idiomorphic grains evenly distributed over the rock. As a component of the matrix, the




Fig. 2. Orientation diagram for {001} of 100 chlorites (dots) and for [001] of 50 enstatites. The great circle corresponds to the macroscopic schistosity of the rock and L marks the lineation.
Specimen Mg 7, Val Cama.

enstatite is generally free of inclusions. The porphyroblastic enstatite, however, contains numerous inclusions of magnetite and forsterite aligned parallel to the foliation. Some of the large enstatites show slight undulatory extinction and occasionally ruptures filled with minor recrystallized granules. The grains of the matrix mosaic are absolutely undeformed.

An orientation diagram (Fig. 2) for (001) of the chlorites is consistent with the macroscopic foliation of the rock. In the same diagram, enstatite [001]-axes show a preferred orientation parallel to the macroscopic lineation and spread out on a great circle parallel to the s-plane. There is no detectable difference in orientation for small or large enstatite grains. Since the lineation and foliation of the ultramafic schists are in complete agreement with the structural properties of the accompanying metamorphic Mesozoic rocks, all the constituents of the matrix mosaic must have crystallized postkinematically during the Alpine metamorphism. The large enstatites are believed to have attained their shape and orientation by Alpine recrystallization.

Chemical Composition of Rock and Minerals

Aside from a high ratio of Fe_2O_3 to FeO , the composition of Mg 7e (Table 1) is comparable to typical harzburgites in the literature. Notable features are the low contents of CaO and alkalis. The high NiO and Cr_2O_3 helps confirm the ultimate igneous origin of the rock. Reduction of some of the Fe_2O_3 suffices to bring normative olivine and orthopyroxene compositions into line with other harzburgites, and, at the same time, reduces total normative orthopyroxene considerably (Table 1).

Table 1. *Chemical Analysis and Norm of Magnetite-Chlorite-Enstatite-Forsterite-Schist, Promagn, Val Cama, Specimen Mg 7e*

	Weight %	C.I.P.W. norms		Mg 7e, all Fe as FeO
SiO_2	43.56			
TiO_2	0.00			
Al_2O_3	1.65	Co	1.57	1.57
Cr_2O_3	0.52	Or	0.06	0.06
Fe_2O_3	4.91	Ab	0.25	0.25
FeO	4.23	An	0.05	0.05
MnO	0.11			
NiO	0.40	Hy	{ En 31.10 Fs 1.18	20.42 3.05
MgO	44.01	Ol	{ Fo 55.01 Fa 2.31	62.50 10.27
CaO	0.01	Mt	7.12	0.00
Na_2O	0.03	Cm	0.77	0.77
K_2O	0.008	Total	99.43	98.94
H_2O^+	1.12			
H_2O^-	0.01			
Total	100.628			

Analysts: A. G. LOOMIS, R. N. JACK, and J. HAMPEL.

Table 2. Microprobe Analyses of Olivine, Orthopyroxene, Chlorite and Magnetite in Specimen Mg 7e

	Olivine	Orthopyroxene	Chlorite	Magnetite
SiO ₂	41.3	57.9	33.17	0.00
TiO ₂	0.00	0.00	0.00	0.09
Al ₂ O ₃	0.00	0.05	13.68	0.50
Cr ₂ O ₃	0.00	0.01	1.35	7.8
V ₂ O ₃	0.00	0.00	0.00	0.25
Fe ₂ O ₃			1.37	61.79 ³⁾
FeO	4.46 ¹⁾	3.4 ¹⁾	1.29	24.03 ³⁾
MgO	53.3	38.1	36.19	3.9
NiO	0.41	0.065	0.22	0.82
MnO	0.10	0.17	0.01	0.18
CaO	0.00	0.03	0.00	0.00
H ₂ O ⁺			12.33 ²⁾	
Total	99.57	99.725	99.61	99.36

Atomic proportions:

Basis:	4 oxygens	3 oxygens	36 (O+OH)	32 oxygens
Si	0.994	0.988	6.285	0.000
Ti	0.000	0.000	0.000	0.020
Al	0.000	0.001	3.056	0.176
Cr	0.000	0.000	0.202	1.843
V	0.000	0.000	0.000	0.060
Fe ³⁺			0.195	13.901
Fe ²⁺	0.090	2.012	0.048	6.008
Mg	1.912	0.970	1.022	10.220
Ni	0.008	0.001	0.033	0.198
Mn	0.002	0.002	0.002	0.046
Ca	0.000	0.001	0.000	0.000
			OH 15.582	

Optical properties:

n_{α}	1.643		1.656		ω 1.579 \pm 0.001
n_{β}	1.661		1.660		
n_{γ}	1.679	\pm 0.001	1.666	\pm 0.001	
A_{γ}	0.036		0.010		0.004 \pm 0.001
$2 V_{\gamma}$	88 \pm 1°		70.3 \pm 0.5°		10°-15°

¹⁾ Total iron. There may be finite amounts of Fe₂O₃ in all three silicates in view of the presence of magnetite.

²⁾ F and Cl were not detected at the 0.1 and 0.01% levels respectively. SiO₂, Al₂O₃, Fe₂O₃, FeO, MgO and H₂O⁺ by A. G. LOOMIS.

³⁾ FeO and Fe₂O₃ calculated from total iron on the basis of a stoichiometric spinel with minor ulvöspinel substitution.

Microprobe analyses of the constituent minerals of Mg 7e (Table 2) revealed their homogeneous nature, the only exception being the small magnetites present as inclusions in the orthopyroxene. Olivine (Fa 4.5) and orthopyroxene (Of 4.8) compositions lie outside the range shown by igneous ultramafics, whether layered, orogenic or xenolithic. Their magnesian compositions are a response to metamorphism under relatively oxidizing conditions. As a result, the relative proportions of NiO and FeO (or MgO) deviate from the rather

well defined trend characteristic of terrestrial olivines of igneous origin (e. g., WHITE 1966; FORBES and BANNO 1966; SMITH 1966; MOORE and EVANS 1967).

The orthopyroxene is perhaps notable for its low contents of CaO, Al₂O₃, and Cr₂O₃. The partitioning of Ni between olivine and orthopyroxene suggests an equilibration temperature of 1310° C (HÄKLI 1968) a value drastically inconsistent with geological evidence¹⁾. The chlorite possesses an even lower Fe/Mg-ratio than olivine or orthopyroxene, and is a Cr-Al-variety, both elements being derived presumably from the original accessory chromite. The remaining Cr is present in a very Ti-poor magnetite, along with some of the Ni formerly present in silicate or sulphide phases.

Conclusions

Textural relationships argue strongly that the constituent phases of the enstatite-olivine rocks of Val Cama, as typified by specimen Mg 7, attained equilibrium during the last main metamorphic episode, the Alpine metamorphism. The chemical data, in addition, are consistent with a thorough phase of reequilibration at a late stage in the metamorphic history of the rock. As indicated by the parageneses kyanite-sillimanite-muscovite and diopside-calcite-anorthite in associated pelites and marbles respectively, the grade of metamorphism corresponds to medium amphibolite facies. Enstatite and forsterite together with magnetite and chlorite occur frequently in similar ultramafic rocks in the same metamorphic zone of the Lepontine region²⁾. In this case, it is clear that orthopyroxene is not indicative of PT conditions of the severity of granulite facies or pyroxene hornfels facies. Interestingly, enstatite is a possible phase in O'HARA's (1967) chlorite-amphibole-peridotite "facies", which he equates with parts of the amphibolite facies. The same holds for the chlorite-peridotite "facies" of ROST (1968). However, the designation "peridotite facies" seems unnecessary and possibly even confusing.

The stability at about 600° to 650° C of the pair enstatite-forsterite as opposed to anthophyllite-forsterite or talc-forsterite, requires the activity of H₂O to be less than about 0.25 of the activity of pure H₂O (GREENWOOD 1967). These conditions are still capable of stabilizing magnesian chlorite, since the latter can go to approximately 780° C at four kb water pressure (FAWCETT and YODER, 1966). H₂O-activity was not uniformly low during the meta-

¹⁾ Possibly the calibration of this thermometer is incorrect, and the dependence of K on Mg/Fe underestimated. For the exchange equilibrium between olivine and orthopyroxene ΔH and ΔG are apparently both large and opposite in sign, hence ΔS would be unusually large.

²⁾ For instance Alpe Arami, Val Gorduno (not including garnet bearing ultramafics); Corno di Gesero, west of Bellinzona; Val Agro, Verzasca; Val di Moleno; Cima Caneto, Centovalli.

morphism of the ultramafic rocks. Zoned anthophyllite-talc veins at Val Cama and other localities in the Leontine Alps (TROMMSDORFF, ms) point to local gradients in H₂O-activity. It seems unlikely that the enstatite-forsterite rocks of the Leontine formed by simple progressive metamorphism of serpentinite, unless the latter was rich in carbonates. Metamorphism of a dry protolith, or alternatively crystallization under conditions of high CO₂-activity (from the surrounding marbles) was much more probable.

The support of National Science Foundation grant GA 500 is gratefully acknowledged. Thanks are also due to J. Hampel for photomicrography, and to A. G. Loomis, R. N. Jack, and J. Hampel for chemical analyses.

References

- SMPM = Schweiz. Mineralogische und Petrographische Mitteilungen
- BRUGGMANN, H. O. (1965): Geologie und Petrographie des südlichen Misox. Diss. Univ. Zürich.
- DAL VESCO, E. (1953): Genesi e metamorfosi delle rocce basiche e ultrabasiche nell'ambiente mesozonale dell'orogene Pennidico. SMPM 33, 173-480.
- FAWCETT, J. J. and H. S. YODER (1966): Phase Relations of Chlorites in the System MgO-Al₂O₃-SiO₂-H₂O. Amer. Mineral. 51, 353-380.
- FORBES, R. B. and S. BANNO (1966): Nickel-iron Content of Peridotite Inclusion and cognate Olivine from an Alkali-Olivine Basalt. Amer. Mineral. 51, 130-140.
- GREENWOOD, H. J. (1963): The Synthesis and Stability of Anthophyllite. J. Petrol. IV, 317-351.
- (1967): Mineral Equilibria in the System MgO-SiO₂-H₂O-CO₂. In Researches in Geochemistry vol. 2, p. 542-567. P. H. Abelson editor, John Wiley and Sons, Inc. New York.
- GRUBENMANN, U. (1908): Der Granatolivinfels des Gordunotales und seine Begleitgesteine. Vjschr. Natf. Ges. Zürich, 53, 129-156.
- HÄKLI, T. A. (1968): An Attempt to apply the Makaopuhi Nickel fractionation Data to the Temperature Determination of a Basic Intrusive. Geochim. et Cosmochim. Acta 32, 449-460.
- HEZNER, L. (1909): Der Peridotit von Loderio (Tessin). Vjschr. Natf. Ges. Zürich, 54, 244-260.
- KNUP, P. (1958): Geologie und Petrographie des Gebietes zwischen Centovalli-Valle Vigezzo und Onsernone. SMPM 38, 83-238.
- KÜNDIG, E. (1926): Beiträge zur Geologie und Petrographie der Gebirgskette zwischen Calanca und Misox. SMPM 6, 3-100.
- MÖCKEL, J. R. (1968): Structural Petrology of the Garnet-Peridotite of Alpe Arami (Ticino, Switzerland), Leidse Geol. Med. 42, 61-130.
- MOORE, J. G. and B. W. EVANS (1967): The Role of Olivine in the Crystallization of the prehistoric Makaopuhi tholeiitic Lava Lake, Hawaii. Contr. Mineral. Petrol. 15, 202-223.
- NIGGLI, P., H. PREISWERK, O. GRÜTTER, E. BOSSARD und E. KÜNDIG (1933): Geologische Beschreibung der Tessineralpen zwischen Maggia- und Bleniotal. Beitr. geol. Karte Schweiz, NF. 71.

- O'HARA, M. J. (1967): Mineral Facies in Ultrabasic Rocks, p. 7–18 in Ultramafic and Related Rocks, P. J. Wyllie ed. John Wiley and Sons Inc. New York.
- O'HARA, M. J. and MERCY, E. L. P. (1966): Garnet-Peridotite and Eclogite from Bellinzona, Switzerland. *Earth and Planet. Sci. Lett.* I, 295–300.
- ROST, F. (1968): Über die Fazieseinstufung orogenotyper Peridotite und ihre Beziehungen zur Peridotitschale des Erdmantels. *Int. Geol. Congr. Praha*, vol. I, 187–196.
- SMITH, J. V. (1966): X-ray Emission Microanalysis of Rock-forming minerals. II. Olivines. *J. Geol.* 74, 1–16.
- SCHWANDER, H. und E. WENK (1967): Studien mit der Röntgen-Mikrosonde an basischen Plagioklasen alpiner Metamorphite. *SMPM* 47, 225–234.
- TROMMSDORFF, V. (1968): Die Wollastonitreaktion im westlichen Bergell. *SMPM* 48, 828–829.
- WHITE, R. W. (1966): Ultramafic Inclusions in Basaltic Rocks from Hawaii. *Contr. Mineral. Petrol.* 12, 245–314.

Manuscript received April 30th, 1969.