

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 56 (1976)

Heft: 3

Artikel: Heat flow measurements in perialpine lakes along the Swiss
geotraverse

Autor: Finckh, Peter

DOI: <https://doi.org/10.5169/seals-43719>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Heat Flow Measurements in Perialpine Lakes along the Swiss Geotraverse

By *Peter Finckh**)

Heat flow measurements in perialpine lakes were carried out to supplement the scanty information concerning the thermal conditions in the crust underlying central Europe. A probe was built which allowed the measurement of 11 temperature values at equally spaced sub-bottom intervals up to a maximum penetration of 10 m. Measurements of this type were made in Lake Zurich, Lake Zug, Lake Lucerne, Lago Maggiore, Lago di Como, Lago d'Iseo and Lago di Garda. After the retrieval of the probe a sediment core could be collected on samples of which thermal conductivity was measured after treating the material in such a way that exsolved gas bubbles were eliminated. Continuous seismic reflection and refraction profiles in all these lakes provided a set of parameters for gradient correction due to heat flow refraction. Additional corrections were applied on the basis of palynologically determined sedimentation rates and for the influence of the last period of glaciation. Final values obtained in this manner indicate rather high values of about $2.8 \cdot 10^{-6}$ cal/cm²·sec (117 mW/m²) north and south of the Alps. This is partially confirmed by earlier measurements in lakes but shows a considerable discrepancy with measurements from the central Alps.

*) Geologisches Institut, ETH-Zentrum, CH-8092 Zürich.