

Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	57 (1977)
Heft:	2
Artikel:	The Utzenstorf stony meteorite
Autor:	Bevan, A.W.R. / Easton, A.J.
DOI:	https://doi.org/10.5169/seals-44429

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Utzenstorf Stony Meteorite

By *A. W. R. Bevan* and *A. J. Easton* (London) *)

Abstract

The Utzenstorf, canton Bern, Switzerland, meteorite fell at 7 p.m. on August 16, 1928 at $47^{\circ} 7'$ N., $7^{\circ} 33'$ E. This olivine-bronzite, or H-group chondrite contains olivine ($\text{Fa}_{18.3}$), orthopyroxene ($\text{Fs}_{16.4}$), kamacite, taenite, troilite, chromite, phosphate and native copper. A chemical analysis with total Fe = 28.75% (wt) and $(\text{Fe} + \text{Mn}) \times 100 : (\text{Fe} + \text{Mn} + \text{Mg}) = 17.5$ (mol. %), is presented. On the basis of bulk composition, mineralogy and texture, Utzenstorf is classified as an H5 chondrite.

INTRODUCTION

On 16 August, 1928 at 7 p.m. local time a bright fireball travelling east to west and "about the size of the Moon" was seen by a number of witnesses in the area of the town of Utzenstorf, canton Bern, Switzerland. The fireball was visible for about four seconds and was accompanied by five or six detonations in rapid succession. Eleven days later, on the farm Stigli (lat. $47^{\circ} 7'$ N., long. $7^{\circ} 33'$ E.), a farmworker discovered three interlocking fragments of black crusted stone totalling 3422 grams in a hole approximately 25 cm deep. Some soil was adhering to the fragments which were washed clean under running water.

On 17 September, after reading a report of the fall in a local paper, Ernst Scheidegger, a teacher in Bern, travelled to Utzenstorf and returned the same day with the largest fragment of 2764 grams. He deposited it at the Naturhistorisches Museum where it was recognised as a meteorite (from GERBER, 1929 and HEY, 1966).

When fitted together the three fragments formed an incompletely crusted, orientated individual approximately $22 \text{ cm} \times 14 \text{ cm} \times 8 \text{ cm}$ which had broken on impact. Some small fragments had obviously become detached during atmospheric flight but these were not found.

*) Department of Mineralogy, British Museum (Nat. Hist.), London SW 7 5 BD.

The meteorite was described, analysed and figured by HUTTENLOCHER and HÜGR (1952) who classified it as an olivine-bronzite chondrite, and gave details of petrography and fusion crust. The present study reports a new chemical

Table 1. *Chlorination analysis of the Utzenstorf meteorite (B.M. 1975, M. 14)*

Non-magnetic attacked (sulphides)		wt%	Metal composition (kamacite)	
Fe	3.55		Fe	93.0%
Mn	(12 ppm)		Ni	6.6%
S	2.04		Co	0.4%
Cr	(6 ppm)		Ge	(52 ppm)
Ti	(34 ppm)		Ga	(6.8 ppm)
Ga	(<1 ppm)			
Ge	(<1 ppm)			
Magnetic attacked (Ni-poor metal and schreibersite)		Metal composition (kamacite)		
Fe	17.79		Fe	93.0%
Ni	1.26		Ni	6.6%
Co	0.085		Co	0.4%
Ge	(10 ppm)		Ge	(52 ppm)
Ga	(1.3 ppm)		Ga	(6.8 ppm)
P	n. f.			
Magnetic unattacked (Ni-rich metal)		Metal composition (taenite)		
Fe	0.42		Fe	49.3%
Ni	0.43		Ni	50.5%
Co	(17 ppm)		Co	0.2%
Ge	(1.03 ppm)		Ge	(121 ppm)
Ga	(0.35 ppm)		Ga	(41 ppm)
Non-magnetic unattacked (silicates, phosphates, oxides)		CIPW norm		
SiO ₂	35.93	Olivine	Fo	22.91
TiO ₂	0.28		Fa	6.34
Al ₂ O ₃	1.46	Hypersthene	En	22.45
Cr ₂ O ₃	0.50		Fs	5.43
Fe ₂ O ₃	0.67	Diopside	Wo	2.93
FeO	8.38		En	2.13
MnO	0.28	Plagioclase	Fs	0.53
MgO	23.06		Ab	7.18
CaO	1.70	Apatite	An	0.00
Na ₂ O	0.85		Or	0.61
K ₂ O	0.10	Chromite		
P ₂ O ₅	0.24			
NiO	0.06	Ilmenite		
H ₂ O ⁺	0.40			
C	0.30	Magnetite		
Ge	(0.5 ppm)			
Ga	(1 ppm)	Kamacite		
Total	99.79			
Total Fe = 28.75%		Taenite		
		Troilite		
		Rest		

n. f. = not found

Analyst: A. J. EASTON

analysis (Table 1) using an improved version of the chlorination technique of MOSS et al., (1967), and electron microprobe analyses of selected minerals in the meteorite (Table 2).

Table 2. *Electron microprobe analyses of selected minerals from the Utzenstorf meteorite*

	Olivine	Orthopyroxene	Chromite	
			¹⁾	²⁾
SiO ₂	39.51	57.07	—	—
TiO ₂	—	0.18	2.06	1.83
Al ₂ O ₃	—	0.19	6.91	7.13
V ₂ O ₅	—	—	0.85	0.89
Cr ₂ O ₃	—	0.02	57.46	56.99
FeO	17.28*)	10.83*)	28.18*)	28.51*)
MnO	—	—	1.46	1.48
MgO	43.34	30.93	3.46	2.87
CaO	0.03	0.60	—	—
Total	100.16	99.82	100.38	99.70
	(Fa _{18.3})	(Fs _{16.2} En _{82.6} Wo _{1.2})		

¹⁾ Chromite associated with metallic nickel-iron.

²⁾ Chromite in matrix.

*) Total Fe as FeO.

— = not looked for.

Analytical conditions: Cambridge Instruments 'Geoscan' microprobe. Standards employed, independently analysed minerals and pure metals, and the results were corrected according to the method outlined by SWEATMAN and LONG (1969) and using the BM-IC-NPL computer program compiled by MASON, FROST and REED (1969). Analyst: A. W. R. BEVAN.

MACRO AND MICROSCOPIC FEATURES

The meteorite is a grey crystalline chondrite. Terrestrial oxidation has turned the crust a dark brown colour and has also penetrated the interior producing patches of rust around metal particles. The oxidation, and relatively high bulk Fe₂O₃ content (0.67% wt) for an observed fall can probably be attributed to weathering as a result of heavy rain during the period between fall and recovery and to its being washed under running water. On a cut surface a few chondrules up to 2 mm can be distinguished; also abundant metal particles that are evenly distributed.

In thin section, numerous poorly defined chondrules commonly 0.4 mm across, and chondrule fragments, composed essentially of olivine or orthopyroxene occur in a microcrystalline matrix. The chondrules display a variety of morphological types including radiating pyroxene chondrules, and barred, and porphyritic olivine chondrules. Clinobronzite and plagioclase grains are rare and occur as small crystallites in the matrix. Metallic nickel-iron, troilite, chromite, phosphate and native copper are also present.

Nine olivine grains and six orthopyroxene grains were analysed by electron microprobe and found to be homogeneous with mean compositions $\text{Fa}_{18.3}$ and $\text{Fs}_{16.2} \text{En}_{82.5} \text{Wo}_{1.3}$ respectively. The olivine composition is consistent with that (Fa_{18}) obtained for Utzenstorf by MASON (1963) using X-ray diffraction.

Metallic nickel-iron is the most abundant opaque phase occurring as irregular and sometimes rounded grains up to 1 mm in size. Kamacite (α -Fe, Ni) with 6.6–6.8% Ni is more abundant than taenite (γ -Fe, Ni) with 48.9–49.5% Ni (analyses by electron microprobe). Cores of some taenite grains have decomposed to plessite ($\alpha + \gamma$ -Fe, Ni). Troilite occurs in association with metal and also as irregular masses up to 0.12 mm across. Grains near the fusion crust and within the heat affected zone have decomposed to fine-grained eutectic-like intergrowths.

Chromite occurs in various textural associations: coarse-grained chromite (0.10–0.15 mm) associated with metal and troilite; discrete grains (0.03–0.06 mm) in the matrix; and aggregates of small ($\sim 6 \mu\text{m}$) rounded grains also in the matrix. Analyses of coarse-grained and matrix chromite are given in Table 2. The compositions are consistent with that for H-group chondrites (BUNCH, et al., 1967; BUNCH and OLSEN, 1975). Some chromite grains enclose phosphates but were too small to analyse by electron microprobe.

Native copper is rare and occurs as aggregates of irregular grains ($\sim 5 \mu\text{m}$) in nickel-iron surrounded by troilite.

BULK CHEMISTRY

The chemical analysis reported in Table 1 was obtained from a 3.4 gram sample of Utzenstorf (B.M. 1975, M. 14). The chemical composition and calculated total Fe/SiO_2 and $\text{Fe}^\circ/\text{total Fe}$ ratios of 0.80 and 0.63, respectively, are consistent with an H-group classification of the meteorite (VAN SCHMUS and WOOD, 1967). Calculated Ni contents of both kamacite and taenite are in excellent agreement with those determined by electron microprobe and the normative silicate $(\text{Fe} + \text{Mn}) \times 100 : (\text{Fe} + \text{Mn} + \text{Mg})$ molecular ratio, 17.5, is close to that derived from the electron microprobe analyses of olivine and pyroxene.

The method of analysis (Moss, et al., 1967), uses a magnetic separation followed by selective attack of dry Cl_2 on kamacite and sulphide, leaving silicates and taenite essentially unaffected. This enables the major and some trace elements to be determined in a number of fractions of chondritic meteorites. Gallium and germanium concentrations in different fractions of Utzenstorf are given in Table 1. The ratio of Ga in metal to Ga in silicate, 6.1, is slightly high but within experimental error for the range of values for H 4–6 chondrites (2.8–6.0) (CHOU, et al., 1973). The high Ni concentration,

470 ppm, of the non-magnetic, unattacked fraction is probably the result of terrestrial oxidation. Schreibersite was not detected.

CONCLUSIONS

The composition of the olivine ($\text{Fa}_{18.3}$), abundance of free metal, and total Fe content of 28.75 wt% confirm HUTTENLOCHER and HÜGI's (1952) classification of Utzenstorf as an olivine-bronzite or H-group chondrite. The absence of polysynthetically twinned clinopyroxene and well developed plagioclase grains indicates that the meteorite belongs to petrologic grade 5 of the VAN SCHMUS and WOOD (1967) classification.

Acknowledgements

The authors thank Professor H. A. Stalder for making a sample of the meteorite available for study; and Drs R. Hutchison and A. L. Graham for advice and critically reading the manuscript.

References

BUNCH, T. E., K. KEIL and K. G. SNETSINGER (1967): Chromite composition in relation to chemistry and texture of ordinary chondrites. *Geochim. Cosmochim. Acta*, **31**, 1569–1582.

BUNCH, T. E. and E. OLSEN (1975): Distribution and significance of chromium in meteorites. *Geochim. Cosmochim. Acta*, **39**, 911–927.

CHOU, C.-L., P. A. BAEDECKER and J. T. WASSON (1973): Distribution of Ni, Ga, Ge and Ir between metal and silicate portions of H-group chondrites. *Geochim. Cosmochim. Acta*, **37**, 2159–2171.

GERBER, E. (1929): Vorläufige Mitteilungen über den Meteorstein von Utzenstorf (Unteremmental), Kanton Bern. *Mitt. naturf. Ges. Bern*, 1928, 25–31.

HEY, M. H. (1966): Catalogue of Meteorites, 3rd edition, British Museum (Nat. Hist.), London.

HUTTENLOCHER, H. and T. HÜGI (1952): Der Meteorit von Utzenstorf. *Mitt. naturf. Ges. Bern*, ser. 2, **9**, 67–128.

MASON, B. (1963): Olivine composition in chondrites. *Geochim. Cosmochim. Acta*, **27**, 1011–1023.

MASON, P. K., M. J. FROST and S. J. B. REED (1969): BM-IC-NPL computer program for calculating corrections in quantitative X-ray microanalysis. National Physical Laboratory, I.M.S. Report 1.

Moss, A. A., M. H. HEY, C. J. ELLIOTT and A. J. EASTON (1967): Methods for the chemical analysis of meteorites: II. The major and some minor constituents of chondrites. *Mineralog. Mag.*, **36**, 101–119.

SWEATMAN, T. R. and J. V. P. LONG (1969): Quantitative electron-probe analysis of rock forming minerals. *J. Petrol.*, **10**, 332–379.

VAN SCHMUS, W. R. and J. A. WOOD (1967): A chemical-petrologic classification for the chondritic meteorites. *Geochim. Cosmochim. Acta*, **31**, 747–765.