Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =

Bulletin suisse de minéralogie et pétrographie

Band: 62 (1982)

Heft: 3

Artikel: Petrology and geochemistry of Fe-Ti gabbros and plagiogranites from

the Western Alps ophiolites

Autor: Pognante, U. / Lombardo, B. / Venturelli, G.

DOI: https://doi.org/10.5169/seals-47981

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 20.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Petrology and Geochemistry of Fe-Ti Gabbros and Plagiogranites from the Western Alps Ophiolites

by U. Pognante¹, B. Lombardo² and G. Venturelli³

Abstract

The geochemistry and mineral chemistry of two suites of Fe-Ti gabbros and plagiogranites from the Western Alps meta-ophiolites have been investigated and compared to data from oceanic analogues and to Fe-rich differentiates from some stratiform continental intrusions. Low pressure (<6 kb) fractionation under initial reduced conditions and late stage relative oxidation associated with filter pressing and cumulus processes were operative during the production of the Fe-Ti gabbros. In the later stages the fractionation could have been assisted by immiscibility processes in which Zr doesn't display the expected partition between the two immiscible liquids.

The stratigraphic position of the Alpine intrusive differentiates indicate that these rocks were either concentrated at various levels of gabbroic magma chambers or expelled as dykes in the host peridotites.

Introduction and Geologic Setting

Fe-Ti gabbros are well known from the Atlantic and Indian oceanic crusts (MIYASHIRO et al., 1970; BONATTI et al., 1971; THOMPSON, 1973; ENGEL and FISHER, 1975; PRINZ et al., 1976) and from some ophiolite complexes (OHNENSTETTER et al., 1975; BECCALUVA et al., 1976, 1977; CHURCH and RICCIO, 1977; LOMBARDO et al., 1978; OHNENSTETTER and OHNENSTETTER, 1980; SERRI, 1980), but are typically absent in other ophiolite suites (e. g. Troodos). Accordingly the strong Fe-Ti enrichment during differentiation of basaltic liquids was recently considered to be a typical feature of some accreting plate margins where these rocks are usually associated with subordinate amounts of plagiogranites (SERRI, 1981).

In this paper new data for two representative suites (ROCCIAVRÉ and LANZO, Fig. 1) of Fe-Ti gabbros and plagiogranites from the Western Alps meta-ophi-

¹ Istituto di Petrografia, Università di Torino, via S. Massimo 24, I-10123 Torino, Italy.

² Centro di Studio sui problemi dell'orogeno delle Alpi Occidentali, Torino, Italy.

³ Istituto di Mineralogia, Università di Parma, Parma, Italy.




Fig. 1 Tectonic sketch map of the Western Alps with location of the Rocciavré (R) and Lanzo (L) ophiolite sections. Piedmont Ophiolite Nappe: 1) main ophiolite bodies and 2) undifferentiated metasediments with minor ophiolites; 3) Dora-Maira and Gran Paradiso continental units; 4) San Bernardo continental unit; 5) Sesia-Lanzo continental unit.

olites are presented, and the petrogenesis of these suites is discussed and compared with that of the present-day oceanic analogues and with the Fe-rich differentiates occurring in some stratiform continental intrusions.

The Alpine tectonic-metamorphic evolution and the petrographical and geochemical features indicate that both the Rocciavré and the Lanzo suites belong to the Piedmont Ophiolite Nappe. This nappe suffered an eclogitic to blueschist equilibration which is believed to reflect Late Cretaceous subduction of a Late Jurassic-Cretaceous oceanic lithosphere (DAL PIAZ, 1974).

Field Relationships

Fe-Ti gabbros occur in the Western Alps meta-ophiolites (Piedmont Ophiolite Nappe) as: 1) layered sequences within isotropic gabbros, 2) dykes cutting peridotites, 3) pods within olivine gabbros and troctolites (Lombardo and Pognante, in press). Plagiogranites are uncommon and occur associated to Fe-Ti gabbros or as dykes cutting peridotites.

This work deals with the Rocciavré and Lanzo Fe-Ti gabbro suites (Fig. 1) which still display clear relationships with the surrounding rocks and largely preserve the primary mineralogy and textures. Both suites were subjected to the

Alpine eclogitic metamorphism (400-480°C and 12-13 kb for Rocciavré; Pognante, 1981), but part of them largely escaped deformation and metamorphism only produced coronitic or incomplete pseudomorphic transformations.

The Rocciavré suite (Pognante, 1979, 1981) is a layered sequence included in a large body of isotropic olivine-orthopyroxene gabbros and consists of centimetric-metric layers of gabbros rich in ilmenite and Ti-magnetite (ilm-mt gabbros) interlayered with gabbro-norites. Small pods of plagiogranite rarely occur in close association with the ilm-mt gabbros. At Lanzo, ilm-mt gabbros, ilm-mt diorites and plagiogranites occur along with more abundant olivine gabbros as cross cutting dykes within tectonite peridotites. The latter consist of spinel lherzolites largely re-equilibrated into plagioclase lherzolites with minor dunites and harzburgites (BOUDIER, 1978).

Prior to the eclogite metamorphism the Fe-Ti gabbros and the plagiogranites suffered partial recrystallization under low pressure amphibolite facies conditions which we refer to an «oceanic» metamorphism. At Lanzo this stage predates the emplacement of basalt dykes and is associated with up to kilometric ductile shear zones.

Petrographic Features

The magmatic mineralogy of the analyzed gabbro-norites, ilm-mt gabbros and plagiogranites from the Rocciavré and Lanzo suites is summarized in Table 1. The gabbro-norites and the ilm-mt gabbros are medium grained to pegmatoid rocks with panallotriomorphic to ophitic textures, and mainly differ in the amount of Fe-Ti oxides and apatite. Fine grained varieties occur in the margins of some ilm-mt gabbro dykes from Lanzo. The Lanzo ilm-mt gabbros are distinguished by the occurrence of olivine and of primary brown amphibole which appear to be absent in the Rocciavré analogues. The plagiogranites consist of altered Na-rich plagioclase in granophyric intergrowth with quartz and of clinopyroxene (Rocciavré), or of albite and brown amphibole (Lanzo). The accessories are zircon, apatite, allanite, Fe-Ti oxides and sphene.

Microstructural considerations suggest that both in the gabbro-norites and in the ilm-mt gabbros the first phases to crystallize are olivine and plagioclase followed by clinopyroxene, orthopyroxene and finally by ilmenite, Ti-magnetite, apatite, zircon and amphibole.

Mineral Chemistry

Compositional data for the magmatic minerals from Rocciavré and Lanzo were obtained with an ARL-SEMQ electron microprobe and are presented in Tables 2-3.

Table 1 List of the analyzed samples (Tables 2-3-4). Abbreviations: gb = gabbro; nor = norite; plagiogr = plagiogranite; pl = plagioclase; ol = olivine; cpx = clinopy-

roxene; opx = magmatic min	roxene; opx = orthopyroxene; amph = amphibole; ilm magmatic minerals pseudomorphed by Alpine and/or «		ilmenite; mt = T ceanic» minerals.	i-magnetite; ap = apatite; zr The minerals are given in orc	roxene; opx = orthopyroxene; amph = amphibole; ilm = ilmenite; mt = Ti-magnetite; ap = apatite; zr = zircon; all = allanite; sph = sphene; qtz = quartz. Underlined: magmatic minerals pseudomorphed by Alpine and/or «oceanic» minerals. The minerals are given in order of decreasing abundance.
	Rock type	Locality		Field occurrence	Magmatic mineralogy
Rocciavré					
R327	gabbro	Sangone	valley	isotropic gabbro	
8881	=	Punta Vailere	ilere	=	b xd5
R299	ol gb-nor	Fonte Ne	Neiretto	layered sequence	cpx, 0],
R305	Ξ.	=	=	= .	cpx, opx, ol,
R320	gb-norite	=	=	=	cpx, opx, ilm
R322	Ξ	=	=	=	cpx,
R323	=	=	=	=	cpx, opx, ilm
R356	=	=	=	=======================================	cpx, opx,
8937	=	=	=	=	cpx, opx,
8940	=	=	₹.	Ξ	cpx, opx,
R321	ilm gb-nor	=	=	=	cpx, opx,
R319	ilm-mt gb	=	=	=	cpx, ilm,
R841	=	=	=	=	cpx, opx,
R849	Æ	=	=	=	cpx, opx,
8944	=	Ξ	=	=	cpx, ilm,
8945	=	=	=	=	cpx, ilm, mt
R296	plagiogr	=	=	=	pl, qtz, cpx, ap, zr, all, ilm
Lanzo					
VL17	ilm-mt gb	Monte Ar	rpone	chilled dyke	
VL18	=	=	=	dyke	
VL23	=	=	=	=	
VL55	plagiogr	Valdella	atorre	Ξ	
۷۲58	Ξ	2		= :	amph, sph, zr,
7162	=	z		= 0	amph, 11m, zr, sph
۷۲68	Ξ	i i		=	pl, amph, ap, ilm, sph, zr, all

Table 2 Representative microprobe analyses of magmatic minerals from the Rocciavré gabbro-norites and ilm-mt gabbros. Structural formulae are on the basis of 4 oxygens for olivine, 6 for pyroxenes and ilmenite, 8 for plagioclase.

	01	<u>P1</u>	Срх				0px		Ilm
	R305	R322	R305	R322	R841	R849	R305	R322	R841
SiO ₂	39.94	54.78	52.34	51.37	52.87	52.74	55.14	53.84	- ',
TiO ₂	<u></u>	0.09	0.79	1.04	0.56	0.92	0.39	0.32	48.63
A1 ₂ 0 ₃	0.03	28.38	2.52	2.30	1.18	2.20	1.29	1.15	
Cr ₂ 0 ₃	0.08	_	0.16	_	0.08	-	_	-	_
FeO tot	16.53	0.17	6.23	9.52	10.81	9.66	16.11	16.58	49.78
MnO	0.17	-	0.24	0.28	0.19	0.26	0.40	0.32	0.34
Mg0	42.84	-	14.97	14.73	14.50	14.31	26.39	25.48	0.66
Ca0	0.02	10.47	21.74	20.56	19.01	19.48	1.18	2.30	-,
Na ₂ 0	_	6.02	0.69	0.41	1.03	0.67	0.07	0.11	-
K ₂ 0	-	0.02	-	-	-	-	-	-	-
Tot.	99.61	99.93	99.68	100.21	100.23	100.24	100.97	100.10	99.39
\$i	1.012	2.475	1.936	1.915	1.971	1.954	1.974	1.959	_
Al	0.001	1.511	0.110	0.101	0.052	0.096	0.054	0.049	-
Ti	_	0.003	0.022	0.029	0.016	0.026	0.010	0.009	1.891
Cr_	0.002	-	0.005	-	0.002	-	-	-	_
Fe tot	0.350	0.006	0.193	0.297	0.337	0.299	0.482	0.505	2.151
Mn	0.004	-	0.008	0.009	0.006	0.008	0.012	0.010	0.015
Mg	1.618	=	0.826	0.814	0.806	0.791	1.408	1.382	0.050
Ca	0.001	0.507	0.862	0.821	0.759	0.773	0.045	0.090	-
Na	-	0.527	0.049	0.030	0.074	0.048	0.005	0.008	-
K		0.001	-	-	-	_	-	-	-
Tot.	2.988	5.030	4.011	4.016	4.023	3.955	3.990	4.012	4.107

Table 3 Representative microprobe analyses of magmatic minerals from the Lanzo ilm-mt gabbros and plagiogranites. Structural formulae are on the basis of 6 oxygens for clinopyroxene and ilmenite, 8 for plagioclase, 12 for magnetite and 23 for amphibole.

	Срх	 	Amph		<u>P1</u>	Ilm	Ti-Mt
	V1 17	V1 23	Vl 23	V1 68	V1 62	V1 23	V1 23
Si0 ₂	51.66	51.92	42.35	51.13	66.71	0.19	-
TiO ₂	0.94	0.99	3.89	2.02	-	51.70	10.77
A1 ₂ 0 ₃	2.32	2.47	10.80	5.62	20.48	0.18	1.94
Cr ₂ 0 ₃	=	0.31	-	0.11	=	- F	20 46.37
e0 tot	12.87	10.42	15.10	10.64	0.28	46.78 F	0 40.19
4n0	0.36	0.28	0.19	0.31	-	0.31	-
1g0	12.68	13.89	10.96	17.57	-	0.33	0.17
Ca0	18.71	18.54	10.82	7.40	0.30	-	
Na ₂ 0	0.60	0.87	3.57	2.89	12.25	-	-
20	-	-	0.05	0.07	0.35	-	
Tot.	100.14	99.69	97.73	97.76	100.37	99.49	99.44
Si	1.944	1.941	6.332	7.282	2.931	0.010	_
A1	0.103	0.109	1.903	0.943	1.061	0.011	0.259
Γi	0.027	0.028	0.437	0.216	-	1.971	0.918
r		0.009	_	0.012	-	- F	3+ e 3.956
e tot	0.405	0.325	1.888	1.267	0.010	1.983 F	e ²⁺ 3.811
in .	0.011	0.009	0.024	0.038	-	0.013	_
1g	0.711	0.779	2.445	3.731	-	0.025	0.02
Ca	0.754	0.743	1.733	1.135	0.014	-	
Na	0.044	0.063	1.035	0.798	1.044	-	-
K	-		0.012	0.015	0.020	-	-
Tot.	3.999	4.006	15.807	15.437	5.080	4.013	8.973




Fig. 2 Compositional variation of magmatic clinopyroxenes in the Rocciavré and Lanzo gabbros. Rocciavré: 1) isotropic gabbros; 2) gabbro-norites; 3) ilm-mt gabbros. Lanzo: 4) ilm-mt gabbros. The compositional fields of the Rocciavré pyroxenes (5) and olivines (black area), and of the Lanzo clinopyroxenes (6) are shown in the inset. Data from Tables 2-3 and 7 additional samples from Rocciavré.

Relics of *olivine* were only found in some Rocciavré gabbro-norites where it is a Mg-rich variety ranging from Fo_{82} to Fo_{84} (Fig. 2, Table 2).

Plagioclase occurs as rare relics in the Rocciavré gabbro-norites and in a few plagiogranites and ilm-mt diorites from Lanzo. In the gabbro-norites it displays an intermediate composition and occasionally shows slight zoning (An 45-55; Table 2). In the plagiogranites and in the ilm-mt diorites the plagioclases are albite (An 1-2; Table 3) and andesine (An 35) respectively.

At Rocciavré the *clinopyroxene* is diopside or endiopside (Ca₄₆Mg₄₅Fe₉) in the isotropic gabbros (Lombardo and Pognante, in press) and augite both in the gabbro-norites (Ca₄₄Mg₄₄Fe₁₂) and in the ilm-mt gabbros (Ca₄₄Mg₄₁Fe₁₅; Fig. 2, Table 2). A slightly Fe-richer augite (Ca₄₀Mg₄₀Fe₂₀; Fig. 2, Table 3) occurs in the Lanzo ilm-mt gabbros. A subsolidus trend toward Ca-richer compositions is obvious from Fig. 2 and is also inferred by the abundance of exsolution lamellae of Ca-poor pyroxene (Ca₂Mg₇₃Fe₂₅). The compositions of the analyzed clinopyroxenes are comparable to those from the Alps-Apennines gabbros (Serri, 1980; Lombardo and Pognante, in press; Mevel, pers. com.) and from the oceanic gabbros (Hodges and Papike, 1976; Prinz et al., 1976), which also exhibit only limited Fe-enrichment during differentiation. On the contrary these pyroxenes differ from those of some stratiform intrusions emplaced in the continental crust (e. g. Skaergaard and Bushveld; Wager and Brown, 1967) where the Fe-rich differentiates contain significantly Fe-richer pyroxenes.

The orthopyroxenes from the Rocciavré gabbro-norites are bronzites $(Ca_4Mg_{71}Fe_{25}; Fig. 2, Table 2)$ and display a subsolidus re-equilibration toward Ca-poorer compositions with thin exsolutions of clinopyroxene. A Mg-richer variety $(Ca_1Mg_{82}Fe_{17})$ occurs in some reaction rims around olivine.

In the Rocciavré gabbro-norites and in the Lanzo plagiogranites the Fe-Ti oxides are mainly ilmenite (Ilm₉₉Hem₁; Table 2). A more complex pattern is displayed by the ilm-mt gabbros; here in fact the oxides consist of largely altered Ti-magnetite (Mt₆₉Usp₃₁; Table 3) rich in exsolutions of ilmenite, and of discrete grains of ilmenite (Ilm₉₇Hem₃, Table 2; Ilm₉₉Hem₁, Table 3).

The amphiboles from the Lanzo plagiogranites are distinguished by high values of SiO_2 and of Na_2O (Table 3), and applying the nomenclature proposed by Leake (1978) are classified as magnesio-katophorites. The amphiboles from the Lanzo ilm-mt gabbros are rich in TiO_2 and Na_2O (Table 3), and are classified as titanian magnesio-taramites. Excepted the low K_2O values, the latter are similar to some amphiboles dredged in the Atlantic (PRICHARD and CANN, 1982).

Bulk Rock Chemistry

14 samples from Rocciavré and 5 samples from Lanzo have been analyzed for major and some trace elements (Table 4). SiO₂, TiO₂, Al₂O₃, Fe₂O₃tot, MnO, CaO, K₂O, P₂O₅, Sc, V, Cr, Ni, Y, Zr, Sr, Rb have been determined by X-ray fluorescence (Philips 1400). MgO, Ni, Cr were determined by atomic absorption, and Na₂O by flame photometry.

The wide compositional range displayed by the investigated rocks is apparent in the diagrams of Fig. 3 where several major and trace elements are plotted against FeOtot/MgO. In spite of the variable element mobilization during the different metamorphic events, in the AFM diagram (Fig. 4) the analyzed rocks preserve a tholeiitic trend. However, in contrast with the present-day oceanic gabbroids, the Alpine rocks show a moderate Na-enrichment probably obtained during metamorphism.

Low values of FeOtot/MgO (0.30-0.37), Ti, P, Zr, Y, V and high Cr, Ni characterize the Rocciavré isotropic gabbros, whereas higher values of FeOtot/MgO (0.61-1.25), Ti, P, Zr, Y and lower Cr, Ni distinguish the gabbronorites. A dramatic increase of FeOtot/MgO (3.2-5.6), Ti and V is shown by the ilm-mt gabbros and is mainly influenced by the segregation of ilmenite and Ti-magnetite. An exception is sample 8944 which displays very high FeOtot/MgO, P, Y but low V reflecting the abundance of ilmenite and apatite. The relatively high Ni values distinguishing the ilm-mt gabbros from Lanzo are related to the occurrence of olivine. The plagiogranites show high amounts of Si, Na, Zr, Y and low FeOtot/MgO, Mg, Fe, Ca, Ti, V and Sc. The very low content of K may be the effect of metamorphism.

Table 4 Major (wt%) and trace (ppm) element abundances in the gabbroic rocks from Rocciavré and Lanzo. -, not determined; (), less accurate data.

ROCCIAVRE	ROCCIAVRE	VRE	(Ppm)	or common			6400	010100		OCCIONIS	allu La	, II	dalices in the gardride rocks from Nocklavie and Lanzo, not determined, (), less accurate data	miled, ()	LANZO	ourate de	ģ		
	R327	8881	F320	R356	R299	8940	8937	R322	R323	R321	8945	R319	8944	R296	VL23	VL18	VL17	VL55	VL58
Sio	51.6	52.1	51.5	51.1	54.2	50.4	6.64	52.8	50.8	50.6	45.3	44.7	45.9	64.3	43.3	44.2	8.44	61.8	4.69
7 110,	0.23	0.26	0.48	0.48	0.53	0.58	0.48	0.53	97.0	1.3	5.2	6.4	3.1	0.79	5.7	6.5	5.4	0.14	0.20
A1 03	14.8	16.2	12.5	13.8	14.4	12.9	15.5	17.3	15.2	14.1	10.9	9.8	10.0	13.3	10.2	8.6	8.6	19.4	16.0
Fe ₀ tot	4.1	3.9	7.0	6.3	5.9	8.3	8.1	7.1	8.5	10.8	18.6	21.7	23.0	7.5	20.7	21.0	21.0	1.8	1.0
0 U	90.0	0.07	0.12	0.12	0.11	0.13	0.13	0.12	0.16	0.16	0.25	0.26	0.31	0.12	1	1	1	1	1
₩ ₉ 0	12.3	9.5	10.4	6.3	4.8	11.4	6.5	7.7	8.8	7.8	6.4	5.5	3.7	2.9	5.7	5.2	6.4	2.0	1.4
CaO	13.0	13.5	13.4	14.8	12.8	12.3	11.7	10.4	11.2	10.5	11.3	10.5	7.6	4.2	10.1	8.7	9.5	4.5	6.0
0 % 11	2.1	3.1	3.3	3.0	2.6	2.9	3.1	3.3	2.8	3.5	3.1	3.4	2.4	5.4	3.5	4.3	4.5	9.1	10.3
, 0 ²	< 0.05	0.05	1.0	0.04	0.04	0.04	0.05	0.04	0.05	0.09	<0.05	<0.05 ⋅	<0.05	90.0	0.05	0.10	90.0	0.04	< 0.05
P.05	0.07	0.07	0.08	0.08	0.08	0.11	0.09	0.12	0.08	0.10	0.08	0.22	2.0	0.50	0.04	0.11	0.05	90.0	0.18
L.0.1.	1.7	1.1	1.3	6.0	1.1	6.0	1.2	0.68	1.5	1.0	0.3	0.1	0.0	6.0	9.0	0.0	0.0	።	9.0
Š	36	35	94	1,7	0,4	53	0,4	1	13	ι	64	55	34	19	59	51	51	°°	, ,
>	120	130	179	1	1	į	1	Ĺ	ı	·C	911	1030	51	81	975	740	770	23	ထ
ວ	1760	345	178	06	140	96	95	20	54	75	==	~	^	7	18	9	^	* 1	25
;; ;	255	133	132	81	88	16	102	61	76	j	37	04	œ	20	105	86	23	48	59
>	80	=	16	16	11	22	18	15	21	23	25	55	(125)	(150)	25	35	30	37	70
Zr	5.5	o	11	52	23	36	53	29	56	34	20	115	79	415	77	87	55	187	550
Sr	158	182	120	147	158	144	157	167	171	186	153	69	123	88	170	184	165	1300	147
Rb	т	2	က	က	ო	4	3.5	ო	4	m	w	ю	4	2	ဗ	Ŋ	4		r




Fig. 3 Variation diagrams of some major (wt%) and minor (ppm) elements against FeOtot/MgO for the analyzed rocks. Rocciavré: (\Box) isotropic gabbros; (\blacksquare) gabbro-norites; (\blacksquare) ilm-mt gabbros; (\blacksquare) ilm-mt gabbros; (\blacksquare) plagiogranite. Lanzo: (\blacksquare) ilm-mt gabbros; (\blacksquare) plagiogranites. Data from Table 4.




Fig. 4 AFM diagram for the gabbros and plagiogranites from Rocciavré and Lanzo. The compositional field of the oceanic gabbros is also shown (Bonatti et al., 1971; Thompson, 1973; Prinz et. al., 1976). Symbols as in Fig. 3.


From the geochemical data here reported the fractionation range of the Rocciavré and Lanzo rocks appears to overlap with that of other gabbroids from the Alps-Apennines ophiolites (Beccaluva et al., 1977; Lombardo et al., 1978; Serri, 1980) and from modern ocean ridge areas (Miyashiro et al., 1970; Bonatti et al., 1971; Thompson, 1973; Engel and Fisher, 1975; Hodges and Papike, 1976; Prinz et al., 1976). Conversely both the ophiolite and ocean Fe-Ti gabbros differ from the Fe-rich differentiates of the stratiform continental intrusions (e. g. ferrodiorites from Skaergaard; Wager and Brown, 1967) which show higher Fe and lower Ti contents, and a strong iron enrichment of the silicates.


Petrogenesis

Magmatic evolution

Since gabbros mostly result from cumulate liquidus phases plus variable amounts of trapped liquid, their compositions should not define a liquid line-of-descent, but should only give a rough idea of the degree of fractionation of the parental magmas. However the chilled margins of Fe-Ti gabbro composition (sample VL17) in the Lanzo peridotites suggest that such liquids may exist and that gabbros cannot be always regarded as simple cumulus products.

The early stages of crystallization are characterized by the segregation of plagioclase, Cr-clinopyroxene, Mg-olivine and orthopyroxene, which produced





△ Fig. 5 Cr-Zr and Ni-Zr diagrams showing the evolutionary trend of the Rocciavré and Lanzo rocks. Symbols as in Fig. 3.

Fig. 6 Ti/100-Zr, Zr/Y-Zr, Y/Sc-Y diagrams showing the evolutionary trend for the Rocciavré and Lanzo rocks. Symbols as in Fig. 3.

large amounts of *isotropic gabbros*. The geochemical data, in particular the very low contents of Zr and Y, indicate that these rocks consist of little fractionated liquidus phases plus very subordinate amounts of trapped liquid.

In the *gabbro-norites* the abundance of liquidus orthopyroxene, the olivine-liquid reaction relations and the lower values of Mg, Ni, Cr point to derivation from a magma saturated in silica and more fractionated than that producing the isotropic gabbros.

The extreme composition displayed by the *ilm-mt gabbros* can be mainly related to fractionation, though cumulate processes where probably also operative. The evolutionary trends shown in Figs. 5-6 and the diagrams of Figs. 3-4 indicate an enrichment in Fe, Ti, V, Zr/Y, Y/Sc and a rapid fall of some compatible elements (Cr in particular) during differentiation. These trends suggest a crystallization under initial low fO₂ conditions and a crystal fractionation approaching surface equilibrium. A relative oxidation in the later stages leading to the formation of the ilm-mt gabbros seems however to be implied by the se-

gregation of Ti-magnetite with "Fe-poor" silicates, as shown experimentally by SPEIDEL and OSBORN (1967) and HILL and ROEDER (1974). This late stage oxidation would in turn suggest relatively reduced initial oxidation states in the parent magma (Morse et al., 1980).

The precipitation of the Fe-Ti oxides shifts the scarce residual liquid toward SiO₂- and alkali-rich compositions, leading to the formation of the *plagiogranites*. The abundance of primary amphibole both in the plagiogranites and in the ilm-mt gabbros, indicate that relatively high P H₂O conditions have accompanied the later stages at Lanzo.

An alternative or complementary mechanism to obtain these peculiar differentiation trends is liquid immiscibility (Khitarov and Pugin, 1978, with references). Recent experiments (McBirney and Nakamura, 1974; Dixon and Rutherford, 1979) indicate that in tholeite compositions immiscibility occurs, after about 95% of the original liquid has crystallized, producing a liquid strongly enriched in Fe, Ti, P and a SiO₂ enriched liquid. Part of the data obtained here, particularly the chemical similarities between the ilm-mt gabbros and plagiogranites and the two immiscible liquids, are in rough agreement with these experiments. However, if it is assumed that the ilm-mt gabbros represent liquids produced by immiscibility processes, Zr appears to be depleted in the Fe-Ti rich liquid, contrary to some experimental and theoretical patterns (Watson, 1976; Khitarov and Pugin, 1978; Ryerson and Hess, 1978). The contradiction may be resolved if the analyzed ilm-mt gabbros are regarded as immiscible liquids in which Zr has not the expected behaviour, as observed by Eby (1979) in some Monteregian Hills rocks produced by liquid immiscibility.

After their production the Fe-Ti rich and the plagiogranitic melts may have been trapped at various levels of the gabbroid magma chamber (e. g. Rocciavré) or expelled intruding as dykes the surrounding rocks (e. g. Lanzo).

Conditions of crystallization

The predominance of irregular and pegmatoid textures over rhytmic layerings, and the lack of troctolitic cumulates strongly suggest that the Rocciavré gabbro suite formed by «in situ» crystallization and by filter pressing processes in a turbolent environment rather than by undisturbed crystal settling.

Relatively low pressures of crystallization for both the Alpine suites are implied by the early appearance of plagioclase relative to pyroxenes, and by the stability of the olivine-plagioclase pair, which suggest pressures lower than 6-7 kb (Kushiro, 1973; Emslie, 1971). A pressure range of 2-5 kb is suggested for the Rocciavré rocks by some thermodynamic calculations involving the variations of the silica activity with \underline{P} and \underline{T} given the compositions of coexisting phases (Arculus and Wills, 1980; Buchanan et al., 1980). In this pressure

range an upper limit of about 1050°C (Holloway, 1973; Allen and Boett-Cher, 1978) is set, for the crystallization of the Fe-Ti gabbros, by the stability of igneous amphibole. Temperatures calculated for clinopyroxene-orthopyroxene pairs using the geothermometer proposed by Wood and Banno (1973) and Wells (1977) recalibration, range between 930 and 1050°C in the Rocciavré gabbro-norites, pointing to subsolidus re-equilibrations and then to slow cooling at depth. Temperatures for coexisting ilmenite and Ti-magnetite pairs using the geothermometer of Buddington and Lindsley (1964) are lower than 650°C and reflect subsolidus equilibration and/or the effects of metamorphism.

Conclusions

The petrological and geochemical study of the Fe-Ti gabbros and the plagiogranites from the Rocciavré and Lanzo suites confirms the close similarities between the intrusive rocks from the Western Alps ophiolites and those from present-day oceanic crust. Unlike the oceanic analogues, the stratigraphic position of the analyzed differentiated gabbroids can still be recognized. In particular these rocks are not segregated between the gabbro and the basalt sections, but appear to have crystallized at depth, either concentrated at various levels of gabbroid magma chambers, or expelled intruding as dykes the host peridotites.

In the formation of the Fe-Ti gabbros and plagiogranites, fractionation of a tholeiitic melt at relatively low pressure (<6kb) associated with filter pressing and cumulus processes, have been operative. The early stages of fractionation appear to have occurred under low fO₂ conditions, while the production of Fe-Ti gabbros in the later stages was probably obtained under relatively higher oxidation conditions. Immiscibility processes producing a peculiar partition of Zr in the SiO₂-rich liquid may have accompanied these later fractionation stages.

The ophiolite and ocean Fe-Ti rich differentiates show some significant geochemical and mineral chemistry differences with respect to those from stratiform intrusions emplaced into continental crust (e. g. Skaergaard). In fact the high fractionation of the former is controlled, in the later stages, by the precipitation of Fe-Ti oxides rather than by a significant iron enrichment of the silicates. Though these differences can be mainly explained in terms of different compositions of parent magmas, other factors could have acted during differentiation. In particular relatively reduced initial oxidation states are suggested for the parent magmas of the ophiolite and ocean Fe-Ti gabbros, possibly reflecting the lack of interaction with a continental crust.

Acknowledgements

The authors are indebted to Professors E. Callegari, V. Dietrich and G. B. Piccardo for helpful comments and discussions. This work was financially supported by the Centro di Studio sui problemi dell'orogeno delle Alpi Occidentali, C.N.R., Torino and by Ministero della Pubblica Istruzione (University of Parma), 1980. The mineral compositional data were collected in the electron microprobe laboratory of the Institute of Mineralogy and Petrology, University of Modena. The financial support from C.N.R. in the installation and operation of this laboratory is gratefully acknowledged.

References

- ALLEN, J.C., BOETTCHER, A. (1978): Amphiboles in andesite and basalt: II. Stability as a function of P-T-fH₂O-fO₂. Am. Mineral. 63: 1074–1087.
- ARCULUS, R.J., WILLS, K.J.A. (1980): The petrology of plutonic blocks and inclusions from the Lesser Antilles island arc. J. Petrol. 21/4: 743-799.
- BECCALUVA, L., MACCIOTTA, G., PICCARDO, G.B., VENTURELLI, G. (1976): Le rocce gabbriche dell'Appennino centro-settentrionale: II. Le caratteristiche chimiche. Ofioliti 1/2: 107-126.
- BECCALUVA, L., OHNENSTETTER, D., OHNENSTETTER, M., VENTURELLI, G. (1977): The trace element geochemistry of Corsican ophiolites. Contrib. Mineral. Petrol. 64: 11-31.
- BONATTI, E., HONNOREZ, J., FERRARA, G. (1971): Peridotite-gabbro-basalt complex from the equatorial Mid-Atlantic Ridge. Phil. Trans. Roy. Soc. London A 268: 385-402.
- BOUDIER, F. (1978): Structure and petrology of the Lanzo peridotite massif. (Piedmont Alps). Geol. Soc. Am. Bull. 89: 1574-1591.
- BUCHANAN, D.L., NOLAN, J., VILJOEN, E.A. (1980): Determination of Silica Activity in Bushveld Rocks. Contrib. Mineral. Petrol. 73: 311-319.
- BUDDINGTON, A.F., LINDSLEY, D.H. (1964): Iron-Titanium oxide minerals and synthetic equivalents. J. Petrol. 5/2: 310-357.
- Church, W. R., Riccio, L. (1977): Fractionation trends in the Bay of Island ophiolite of Newfoundland: polycyclic cumulate sequences in ophiolites and their classification. Can. J. Earth Sci. 14: 1156-1165.
- Dal Piaz, G.V. (1974): Le métamorphisme alpin de haute pression et basse température dans l'évolution structurale du bassin ophiolitique alpino-apenninique. Boll. Soc. Geol. It. 93: 437–468.
- DIXON, S., RUTHERFORD, M.J. (1979): Plagiogranites as late immiscible liquids in ophiolite and midocean ridge suites: an experimental study. Earth Planet. Sci. Lett. 39: 127-144.
- EBY, G. N. (1979): Mount Johnson, Quebec An example of silicate-liquid immiscibility? Geology 7: 491-494.
- EMSLIE, R.F. (1971): Liquidus relations and subsolidus reactions in some plagioclase-bearing systems. Carnegie Inst. 69: 148–155.
- ENGEL, C.G., FISHER, R.L. (1975): Granitic to ultramafic rock complexes of the Indian ocean ridge system. Western Indian ocean. Geol. Soc. Am. Bull. 86: 1553–1578.
- HILL, R., ROEDER, P. (1974): The crystallization of spinel from basaltic liquid as a function of oxygen fugacity. J. Geol. 82: 709-729.
- HODGES, F. N., PAPIKE, J.J. (1976) D.S.D.P. site 334: Magmatic cumulates from oceanic layer 3. J. Geophys. Res. 81/23: 4135-4151.
- HOLLOWAY, J.R. (1973): The system pargasite-H₂O-CO₂: a model for melting of a hydrous mineral with a mixed-volatile fluid-I. Experimental results to 8 Kbar. Geochim. Cosmochim. Acta 37: 651-666.
- KHITAROV, N.I., PUGIN, V.A. (1978): Liquid immiscibility in natural silicate systems. Geokhimiya 6: 803-820.

- Kushiro, I. (1973): Origin of some magmas in oceanic and circum-oceanic regions. Tectonophysics 17: 211–222.
- LEAKE, B.E. (1978): Nomenclature of amphiboles. Can. Mineral. 16: 501-520.
- LOMBARDO, B., POGNANTE, U. (in press): Tectonic implications in the evolution of the Western Alps ophiolite metagabbros. Ofioliti special issue: Ophiolites and Actualism.
- LOMBARDO, B., NERVO, R., COMPAGNONI, R., MESSIGA, B., KIENAST, J.R., MEVEL, C., FIORA, L., PICCARDO, G. B., LANZA, R. (1978): Osservazioni preliminari sulle ofioliti metamorfiche del Monviso (Alpi Occidentali). Rend. Soc. It. Miner. Petrol. 34: 253-305.
- McBirney, A.R., Nakamura, Y. (1974): Immiscibility in late-stage magmas of the Skaergaard intrusion. Carnegie Inst. 73: 348-352.
- MIYASHIRO, A., SHIDO, F., EWING, M. (1970): Crystallization and differentiation in abyssal tholeites and gabbros from mid-ocean ridges. Earth Planet. Sci. Lett. 39: 127-144.
- MORSE, S.A., LINDSLEY, D.H., WILLIAMS, R.J. (1980): Concerning intensive parameters in the Skaergaard intrusion. Am. Jour. Sci. 280-A: 159-170.
- Ohnenstetter, M., Ohnenstetter, D. (1980): Comparison between Corsican albitites and oceanic plagiogranites. Arch. Sc. Genève 33/2-3: 201-220.
- OHNENSTETTER, D., OHNENSTETTER, M., ROCCI, G. (1975): Tholeiitic cumulates in a high pressure metamorphic belt. Pétrologie 1/4: 291-317.
- POGNANTE, U. (1979): The Orsiera-Rocciavré metaophiolitic complex (Italian Western Alps). Ofioliti 4/2: 183-198.
- Pognante, U. (1981): Magmatic and metamorphic evolution of two Fe-Ti gabbroic series from the Piemonte ophiolite nappe in the Susa Valley areas, Italian Western Alps. Mem. Sci. Geol. Padova 35: 21-34.
- PRICHARD, H. M., CANN, J. R. (1982): Petrology and Mineralogy of dredged Gabbro from Gettysburg Bank, Eastern Atlantic. Contrib. Mineral. Petrol. 79: 46-55.
- PRINZ, M., KEIL, K., GREEN, J.A., REID, A.M., BONATTI, E., HONNOREZ, J. (1976): Ultramafic and mafic dredge samples from the equatorial Mid-Atlantic ridge and fracture zones. J. Geophys. Res. 81/23: 4087-4103.
- RYERSON, F.J., Hess, P.C. (1978): Implications of liquid-liquid distribution coefficients to mineral-liquid partitioning. Geochim. Cosmochim. Acta 37: 2135–2147.
- SERRI, G. (1980): Chemistry and petrology of gabbroic complexes from the northern Apennine ophiolites. Proceedings Int. Ophiolite Symp. Cyprus 1979: 296-313.
- SERRI, G. (1981): The petrochemistry of ophiolite gabbroic complexes: a key for the classification of ophiolites into low-Ti and high-Ti types. Earth Planet. Sci. Lett. 52: 203-212.
- Speidel, D.H., Osborn, E.F. (1967): Element distribution among coexisting phases in the system MgO-FeO-Fe₂O₃-SiO₂ as a function of temperature and oxygen fugacity. Am. Mineral. 52: 1139-1152.
- THOMPSON, G. (1973): Trace element distribution in fractionated oceanic rocks. 2. Gabbros and related rocks. Chem. Geol. 12: 99-111.
- WAGER, L.R., Brown, G.M. (1967): Layered igneous rocks. Oliver & Boyd, Edinburgh and London: 558 pp.
- WATSON, E.B. (1976): Two-liquid partition coefficients: experimental data and geochemical implications. Contrib. Mineral. Petrol. 56: 119-134.
- Wells, P.R.A. (1977): Pyroxene thermometry in simple and complex systems. Contrib. Mineral. Petrol. 62: 129-139.
- Wood, B.J., Banno, S. (1973): Garnet-Orthopyroxene and Orthopyroxene-clinopyroxene Relationships in Simple and Complex Systems. Contrib. Mineral. Petrol. 37: 161-174.