

**Zeitschrift:** Schweizerische mineralogische und petrographische Mitteilungen =  
Bulletin suisse de minéralogie et pétrographie

**Band:** 63 (1983)

**Heft:** 2-3

**Artikel:** Inconsistent chemical data on phlogopite : analyses and genesis of  
phlogopites from Madagascar

**Autor:** Stern, W.B. / Klein, H.H.

**DOI:** <https://doi.org/10.5169/seals-48730>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 02.05.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## **Inconsistent chemical data on phlogopite: analyses and genesis of phlogopites from Madagascar**

by *W. B. Stern*<sup>1</sup> and *H. H. Klein*<sup>2</sup>

### **Abstract**

Precambrian pyroxenites in South Madagascar contain phlogopites being today worldwide a main source of phlogopite used in the electrical industry. Literature data on the chemistry of phlogopites from the two mines of Ampandrandrave and Sakamasy are scarce, and highly inconsistent. At both mining sites well-located samples were collected and analyzed by means of energy dispersive X-ray fluorescence spectrometry. Significant, but small chemical differences between phlogopites from the two sites exist. The local chemical variation turns out to be small, often smaller than the analytical variation (error). Thus, no interdependence between chemical composition and sample localization was detectable.

Based upon new phlogopite analyses and literature data of pyroxenites and diopside from the two sites, a mass balance has been calculated assuming a metasomatic transformation of tectonically affected pyroxenites (metasomatic supply of 4 to 4.5% K<sub>2</sub>O, 0 to 1.5% MgO, ± Al<sub>2</sub>O<sub>3</sub>) as educt, and phlogopites (60 to 70% phlogopite, 5 to 10% fassaitic diopside, 20 to 30% calcite, and 0 to 5% anhydrite) + eluate (Si, Ca) as product, and keeping the volume of educt and product constant. Fluorine present in phlogopites between 1.5 and 3%, but not in pyroxenites, is supposed to be an essential agent in this metasomatic process, which was probably connected with the intrusion of discordant pegmatites present at Ampandrandrave and Sakamasy as well.

*Keywords:* phlogopite, chemical analysis, metasomatism, mineral deposits, Malagasy Republic

### **Introduction**

Madagascar is the main world source of all forms of phlogopite mica, and its output of sheet mica (splittings) amounts to 5% (1972), 10% (1968) respectively of the world splitting production of muscovite/phlogopite (PETKOV, 1972). The major part of the actually available phlogopite mica originates from mines located in the south of Madagascar, today particularly from Ampandrandrave and Sakamasy (fig. 1). Below-ground mining operations in Precambrian pyroxenites began in 1921, and 1935 respectively.

---

<sup>1</sup> Geochemical Labs / Mineralogical Institute, University, Bernoullistr. 30, CH-4056 Basel.

<sup>2</sup> The Swiss Insulating Works, CH-4226 Breitenbach.




Fig.1 Geological sketch map of southern Madagascar.

The whole production is consumed by the electro industry abroad, representing therefore an important factor for the Malagasy economy.

Published chemical analyses of phlogopite from Ampandandrave (N = 6) and Sakamasy (N = 1) are scarce (JAKOB & PARGA, 1932; DE LA ROCHE, 1963; JOO, 1972) and are remarkably inconsistent: whereas the analyses of JAKOB et al. and DE LA ROCHE (op. cit.) represent true phlogopite micas, part of the more recent data of JOO cannot be interpreted as end members of the trioctahedral




Fig. 2 Phlogopites from Ampandandrave and Sakamasy, data taken from literature.

Mg-Fe mica series (fig. 2). Are there striking distinctions due to an eventual inhomogeneity of the phlogopite deposits (in connection with the controversial genesis of these deposits, DE LA ROCHE, 1963) or due to other reasons?

According to the genetical history - metamorphous/metasomatic and/or magmatic - chemical gradients within one and the same deposit are conceivable; detailed indications of the sampling procedure are missing, however, in the cited literature. In general the chemical variation of phlogopite in its solid solution series within one and the same deposit is unknown (the same holds for muscovite, too).

### Geological outline

The productive phlogopite deposits of southern Madagascar are located in steeply dipping pyroxenite lamellae (width some few meters, length up to some kilometers) together with alternating beds of gneisses and leptynites of the Precambrian Androyan System affected by an archaic metamorphism in granulite facies (radiometric age 2100–2500 m.a., DELBOS, 1965).

Based above all on geochemical arguments the pyroxenites are interpreted since the works of LACROIX, 1941, DE LA ROCHE and JOO (op. cit.) as metasediments of formerly lagoonal marls.

The rock series is affected by a system of fissures – subparallel and perpendicular to the strike. The enriched occurrence of phlogopite in form of pockets and lenses is related to the tectonic pattern of the host rocks, but the phlogopite itself is tectonically not affected. Pegmatites strike discordantly through the rock series. Radiometric age determinations of the phlogopite (K/Ar, Rb/Sr) revealed 530 m.a. which corresponds to the last pan-african plutonic cycle (LEUTWEIN, 1964).

### Sampling

To clarify the question of the chemical variance of phlogopite from one and the same deposit, samples were collected by the authors in Ampandrandave and Sakamasy, and later analyzed. The test material covers all accessible tunnels in horizontal/lateral and vertical direction (fig. 3). 17 fresh, unaltered phlogopites from each mine were sampled. Help and assistance is gratefully acknowledged to M Fauret/Ampandrandave and M Renot/Sakamasy.

### Analysis

The analyses were carried out by means of energy-dispersive X-ray fluorescence spectrometry (EDS-XFA). Table I indicates the analytical parameters, STERN et al. 1982. The calculated variation ( $3\sigma$ , 99% confidence level) corresponds at Sakamasy to the analytical error (based upon counting statistics) for all main constituents (Si, Al, Fe, Mg, K and F) with exception of water, see table II. Thus, no chemical gradation or zonation whatsoever can be ascertained at Sakamasy. At Ampandrandave Fe and Mg vary within the range of the respective analytical errors, whereas the variation of the other elements is slightly outside the range of analytical error. The chemical variation, however, cannot be correlated with the locality of sampling regarding lateral or vertical directions – the variation seems to be random.




Fig. 3 The mining sites of Ampandrandava and Sakamasy (schematic) with indications on sampling.

The large range of chemical variation as reported by JOO (1972) for phlogopites originating from Sakamasy and Ampandrandava is not confirmed by these analytical series. Most probably the published data represent incorrect analyses and therefore are the consecutive crystal-chemical deductions (JOO, 1972, p. 12 ff.) meaningless.

Whereas the analysis of main constituents leads to results displaying quantifiable errors, the determination of volatile components remains fairly tricky.

Table I Analytical Parameters.

Apparatus	Energy dispersive X-ray fluorescence spectrometer TN-1710 of Tracor Northern. Detector 150 eV resolution. 2000 channel analyzer. Spectrometer designed at Geochem. Labs / Basle University ( Min.-petr. Institute )
Excitation	Ag tube, 21 kV, 0.05 mA. 200 sec counting time (life). Powdered and pressed samples ( STERN, 1979 ).
Analysis	software "Super-ML, matrix correction RH" by Tracor Northern, and "energy calibration, data reduction, plotting" by Gloor Instruments. Reference file 5 / RH 5 on fl 80. Silicate rocks and minerals as standards. Fluorine: ion selective electrodes ( CAMMANN, 1977 ).

Table II Average composition of 17 phlogopites from Ampandandrave and Sakamasy (EDS - XFA) wt-%.

Ampandandrave	$\bar{x}$ N=17	s	CoV*	CoV**	JAKOB et al. 1932
SiO <sub>2</sub>	38.28	0.426	3.3	1.0	38.4
Al <sub>2</sub> O <sub>3</sub>	16.01	.479	9.0	2.8	16.7
Fe <sub>2</sub> O <sub>3</sub>	6.37	.175	8.2	5.8	4.5
MnO	0.18	.015	25.	13.4	0.15
MgO	23.41	.340	4.4	4.5	23.4
K <sub>2</sub> O	9.97	.123	3.7	1.2	10.9
TiO <sub>2</sub>	1.15	.090	24.	2.7	1.69
H <sub>2</sub> O	3.20	.996	92.	11.8	3.2
F	1.53	.201	39.	18.7	.7
Sakamasy					
SiO <sub>2</sub>	38.11	0.026	0.2	1.0	
Al <sub>2</sub> O <sub>3</sub>	16.30	.400	2.5	2.8	
Fe <sub>2</sub> O <sub>3</sub>	6.97	.263	11.3	5.8	
MnO	0.07	.011	43.	13.4	
MgO	23.29	.369	4.8	4.5	
K <sub>2</sub> O	10.21	.062	1.8	1.2	
TiO <sub>2</sub>	.90	.068	23.	2.7	
H <sub>2</sub> O	2.55	.556	66.	11.8	
F	2.41	.183	23.	18.7	
* Coefficient of variation = $3\sigma \cdot 100 / \bar{x}$ evaluation based upon 17 single determinations					
** Coefficient of variation = $3\sigma \cdot 100 / \bar{N}$ based upon counting statistics, connected with analytical error					

The determined loss on ignition is not only dependent on the calcination temperature but also on the calcination time as well as on the grain size distribution of a sample (fig. 4). Since only calcination time and temperature are usually kept constant (they differ, however, from one laboratory to another), and since little is known about the actual grain size distribution, published data on volatile contents are inconsistent and subject to remarkable errors, see table III.

Phlogopites of Madagascar contain fluorine as a main component which is nowadays accurately determined by means of ionselective electrodes, whereas




Fig. 4 Weight loss of mica during calcination at different temperatures, static thermogravimetry. For small grain size fractions a creeping weight loss is observed (after KLEIN et al., 1982).

the old distillation methods gave often erroneous results (VGENOPOULOS, 1971). Whether fluorine is partially or entirely removed from phlogopite during calcination depends on the ignition routine used, fig. 5.

Differences in  $H_2O$  and volatile contents found by different laboratories analyzing the same, homogeneous sample, are due to the different calcination methods, whereas the variation of actual fluorine determinations may represent real chemical variations. Phlogopites from Sakamasy are significantly richer in

Table III Analytical variation of volatiles in silicate standards.

standard	NIM-G	NIM-S	NIM-L	NIM-N	NIM-P	NIM-S	
mean	0.77	0.34	2.77	0.64	0.19	0.20	Loss on ignition
relative error, %	11.5	34.1	3.1	186	226	383	
mean	0.58	0.29	2.44	0.41	0.35	0.38	water, $H_2O^+$
relative error, %	42.8	46.4	8.4	54	75	48	
mean	0.14	0.15	0.24	0.11	0.05	0.10	moisture, $H_2O^-$
relative error, %	77	86	62	94	99	79	

relative error = coefficient of variation =  $1\sigma \cdot 100 / \text{mean}$

Data basis : Report No. 1351 of the National Institute for Metallurgy,  
Johannesburg 1972




Fig. 5 Phenomena during calcination of phlogopite (SKY 1/581). Mica chips ground under norm conditions, average grain size less than 60  $\mu\text{m}$ . Experimentally determined points are symbolized by circles, schematic. Details of the calcination routine, like ignition time and temperature influence severely the loss on ignition (LoI) found. Above 1100°C and 1 hour calcination time fluorine and alcalies begin to disappear.

fluorine than those of Ampandandrave, variations, however, being considerable at both mines. A dependancy of the fluorine content in function of the sample location (depth, host rock etc.) could not be established.

### Results

Seventeen phlogopite analyses per mine from both, Ampandandrave and Sakamasy, display significant differences between the two deposits, but only little variations within a site being—in case of the main constituents—in the range of the analytical errors, table IVa/b. This proven homogeneity (fig. 6) of the two phlogopite deposits asks for a new genetical interpretation.

Phlogopites from Madagascar

Table IVa Chemical analyses (EDS - XFA) of phlogopites from Ampandandrave.

A M P A N D R A N D A V E		75-1	75-2	75-3	91-4	91-5	91-6	91-7	91-8	91-9	121-10	121-11	121-12	121-13	121-14	150-15	161-16	161-17
IX		5582	5583	5584	5585	5586	5587	5588	5589	5590	5591	5592	5593	5594	5595	5596	5597	6017
SiO <sub>2</sub>		38.6	38.9	39.0	38.3	38.6	38.2	37.9	37.8	37.9	38.3	38.7	38.3	37.7	38.8	38.3	37.6	37.7
Al <sub>2</sub> O <sub>3</sub>		16.4	16.0	16.4	16.1	16.0	16.3	15.8	15.5	16.5	16.2	14.7	15.6	15.9	16.7	16.4	16.0	15.6
Fe <sub>2</sub> O <sub>3</sub>		2.71	2.52	2.51	2.47	2.44	2.37	2.21	2.47	2.06	2.52	2.50	2.15	2.16	2.39	2.32	2.15	2.36
FeO		3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6	3.6
MnO		.16	.20	.16	.17	.18	.18	.16	.18	.21	.17	.16	.17	.19	.19	.18	.17	.17
MgO		23.6	23.7	23.6	23.3	23.1	23.6	23.2	23.0	23.5	23.4	24.1	23.9	23.2	23.3	23.6	22.9	22.9
CaO		.02	.0	.13	.06	.0	.05	.04	.0	.13	.12	.0	.03	.0	.0	.05	.0	.01
Na <sub>2</sub> O		.4	.4	.3	.3	.2	.3	.4	.4	.2	.3	.3	.3	.3	.2	.2	.3	.3
K <sub>2</sub> O		10.06	9.98	10.09	10.01	10.15	9.98	9.90	9.83	9.89	9.86	9.76	9.89	9.92	10.07	10.12	9.96	9.77
TiO <sub>2</sub>		1.26	1.33	1.21	1.22	1.13	1.18	1.23	1.15	.97	1.21	1.06	1.06	1.12	1.07	1.17	1.15	1.05
Rb <sub>2</sub> O		.07	.06	.06	.06	.06	.07	.07	.07	.04	.07	.07	.07	.06	.08	.07	.06	.05
H <sub>2</sub> O		1.5	1.2	1.3	3.9	3.3	3.2	4.2	4.5	3.6	3.9	3.5	3.8	3.7	2.9	4.0	3.3	3.6
Cl		.08	.06	.06	.04	.06	.05	.04	.05	.07	.08	.08	.04	.05	.06	.06	.03	.05
F		1.6	1.7	1.55	1.2	.95	1.7	1.7	1.55	1.65	1.4	1.45	1.65	1.7	1.55	1.45	1.6	1.65
O = F		.66	.70	.64	.49	.39	.70	.70	.64	.68	.57	.59	.68	.70	.64	.59	.66	.68
Sum		99.32	98.95	99.33	100.74	99.38	100.08	99.75	99.46	99.64	100.19	99.32	99.88	98.90	100.27	100.93	98.16	98.13

Table IVb Chemical analyses (EDS - XFA) of phlogopites from Sakamasy.

	57-1E	57-1W	57-2E	57-2W	57-3E	57-3W	57-4W	72-1E	72-2E	72-2W	72-3E	72-3W	72-4E	72-4W	72-5W	72-6W	72-7W
IX	6000	5599	6004	6005	6008	6009	6012	5598	6002	6003	6007	6006	6011	6010	6014	6015	6016
SiO <sub>2</sub>	38.3	38.4	38.2	38.1	38.4	38.0	37.9	38.4	38.7	38.1	38.4	38.0	38.0	37.9	37.7	37.8	37.4
Al <sub>2</sub> O <sub>3</sub>	16.4	16.7	16.9	16.3	16.9	16.4	16.3	16.3	15.7	16.9	15.8	16.0	15.9	16.0	16.6	15.8	16.2
Fe <sub>2</sub> O <sub>3</sub>	2.14	1.78	2.47	2.27	2.24	2.16	2.63	2.56	2.18	2.19	2.74	2.21	2.74	2.21	2.20	2.02	2.56
FeO	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2	4.2
MnO	.06	.09	.08	.08	.08	.08	.06	.07	.08	.09	.07	.05	.07	.08	.07	.08	.07
MgO	23.4	23.8	23.9	23.2	23.7	23.5	22.8	23.5	23.7	23.2	23.4	23.0	22.9	23.2	22.8	23.2	22.7
CaO	.0	.02	.16	.0	.0	.0	.07	.11	.03	.24	.0	.0	.12	.0	.0	.01	.02
Nb <sub>2</sub> O	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1	.1
K <sub>2</sub> O	10.20	10.27	10.29	10.22	10.25	10.25	10.14	10.31	10.23	10.20	10.22	10.19	10.09	10.19	10.11	10.25	10.14
TiO <sub>2</sub>	.85	.87	.75	.91	.86	.90	.84	.96	1.00	.82	.97	.95	.96	.94	.85	.98	.94
Rb <sub>2</sub> O	.05	.07	.07	.07	.06	.07	.06	.07	.06	.05	.08	.07	.08	.08	.05	.08	.09
H <sub>2</sub> O	2.3	3.1	2.0	2.8	2.1	1.9	2.5	3.9	1.9	2.6	2.2	2.6	2.4	2.4	3.2	2.0	2.0
Cl	.07	.08	.08	.08	.08	.07	.09	.07	.08	.07	.07	.07	.08	.06	.08	.07	.07
F	1.8	2.5	2.5	2.5	2.5	2.6	2.3	2.3	2.6	2.4	2.5	2.3	2.4	2.4	2.4	2.5	2.5
O = F	.74	1.03	1.03	1.03	1.03	1.07	.94	.94	1.07	.98	1.03	.94	.98	.98	.98	1.03	1.03
Sum	99.03	100.85	101.60	99.70	100.34	99.06	98.95	101.91	100.13	100.26	99.08	98.71	99.06	98.78	99.38	98.06	97.96




Fig. 6 Phlogopites from Ampandandrave and Sakamasy, new data, and data taken from literature (cross after JAKOB et al. (1932), DE LA ROCHE (1963) and JOO (1972)).

DE LA ROCHE (1963) assumes a metasomatic transformation of the pyroxenite by supply of potassium and magnesium in a fluid phase. His different attempts to balance the reaction are only partially satisfying since these calculations were based upon Al-free diopside. The actual pyroxenites consist up to 90% of fassaite containing 7 to 11%  $\text{Al}_2\text{O}_3$  (MAJMUNDAR, 1971), which affects the balance heavily.

JOO (1972) postulates a single phase genesis: transformation of lagoonous marns into pyroxenite by metamorphism in granulite facies and at the same time partial/differential transformation (hydatogenesis) of the pyroxenite by supply of marin-inherited potassium, and a fluid phase.

New data suggest a somewhat modified genesis, which is still based on the idea of LACROIX (1941):

- Precambrian pyroxenites - chemically and mineralogically identical with the actual ones (granulite facies metamorphism between 2100 and 2500 m.a.) - are effected by tectonic events: fissures subparallel and perpendicular to the strike of the host rocks are formed, inducing an increased mobility in the tectonically affected zones.







By means of a metasomatic supply relatively small amounts of  $K_2O$ ,  $MgO$ ,  $\pm Al_2O_3$  and volatiles have to be introduced. This metasomatic event may be connected with the intrusion of discordant pegmatites present at Ampandan-drave and Sakamasy as well.

- The correlations of Al-rich pyroxenites delivering the majority of the essential elements for phlogopite formation, and the intrusion of pegmatitic bodies may explain why coarse grained phlogopites are worldwide scarce. A further limiting factor is certainly the necessary supply of fluorine acting as a solvent, and being typical for all known coarse grained phlogopites all over the world (Canada, Ceylon-Sri Lanka, USSR / DUNN, 1961, RIMŠAITE, 1970, SKOW, 1962).

#### Acknowledgments

The authors are indebted to Messrs. Akesson, Bordier and collaborators of SOMIDA for logistic support in Madagascar, to Prof. H. de la Roche and Prof. M. Frey for critical reading, and to Mr. U. Naef for drafting work.

#### Bibliography

- BESAIRIE, H. (1965): Carte géologique de Madagascar 1 : 1 000 000. Service Géol. de Madagascar, Tananarive.
- CAMMANN, K. (1977): *Das Arbeiten mit ionenselektiven Elektroden*. Springer, Heidelberg.
- CLARK, S. P. (1966): Handbook of physical constants. Geol. Soc. America Mem. 97.
- DELBOS, L. (1965): Sur l'âge de quelques minéralisations de Madagascar. EXN. Bull. B.R.G.M. 1/1965.
- DE LA ROCHE, H. (1963): Etude géologique de l'extrême Sud-Est de Madagascar (zone d'intensité métamorphique élevée). Ann. Géol. de Madagascar 28, 1. Thèse Nancy 1958.
- DE LA ROCHE, H. (1963): Sur la composition chimique des phlogopites de Madagascar et sur la présence de variétés riches en fluor. Ann. Géol. de Mad. 33, 175.
- DUNN, J. A. (1961): Mica. Bull. Geol. Surv. India 102.
- JOO, J. (1972): Le mica industriel de Madagascar. Thèse Clermont.
- JAKOB, J., PARGA-PONDAL, I. (1932): Beiträge zur chemischen Konstitution der Glimmer. X. Mitt. Z. Krist. 82, 271.
- KLEIN, H. H., STERN, W. B., WEBER, W. (1982): On physical and chemical properties of ruby muscovite used in the electrical industry. Schweiz. Mineral. petrogr. Mitt., 62, 145.
- LACROIX, A. (1941): Les gisements de phlogopite de Madagascar et les pyroxenites qui les enferment. Ann. Géol. Serv. Mines Madagascar, fasc. X.
- LEUTWEIN, F. (1964): Etude géochimique et géochronologique d'un crystal de phlogopite de Miary, près de Bekily (Madagascar). C.R. Sem. Géol. 1964. Tananarive.
- MAJUMDAR, H. H. (1971): Fassaite from Madagascar. Canad. Mineral., 10, 899.
- PETKOV, B. (1972): Mica. In: Lefond, S. J.: Industrial Minerals and rocks. Amer. Inst. Mining, New-York.
- Report 1351 (1972): of the National Institute for Metallurgy. Johannesburg.

- RIMSAITE, J. (1970): Aionic and cationic variations in zoned phlogopite. *Contrib. Mineral Petrol.* 29, 186.
- SKOW, M. L. (1962): Mica, a material survey. US Dept. of the Interior, Bureau of Mines, Inf. Circ. 8125.
- STERN, W. B. (1979): Probleme der quantitativen röntgenspektrometrischen Analyse von Hauptkomponenten und Spuren in geologischen Proben. *Schweiz. Mineral. petrogr. Mitt.*, 59, 83.
- STERN, W. B., HÄNNI, H. (1982): Energy dispersive X-Ray spectrometry: a non-destructive tool in gemmology. *J. Gemmol.* 18, 285.
- VACHETTE, M., HOTTIN, G. (1977): Rajeunissement au pan-african des chaînes anosyennes, dans la région d'Esira (sud-est de Madagascar). *C.R. Acad. Sc. Paris, Sér. D*, 284, 1369.
- VGENOPOULOS, A. G. (1971): Der Fluorgehalt einiger gesteinsbildender Mineralien aus dem Lepontin. *Schweiz. Mineral. petrogr. Mitt.* 51, 363.

Manuscript received 12 october, 1983.