Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie				
Band:	68 (1988)				
Heft:	1				
Artikel:	Sr-bearing stilbite in a quartz-monzonite from Vathi, Kilkis, Northern Greece				
Autor:	Filippidis, A. / Kougoulis, C. / Michailidis, K.				
DOI:	https://doi.org/10.5169/seals-52050				

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Sr-bearing stilbite in a quartz-monzonite from Vathi, Kilkis, Northern Greece

By A. Filippidis¹, C. Kougoulis² and K. Michailidis¹

Abstract

Sr-bearing stilbite occuring in veinlets, fracture and open space fillings of the Vathi quartz-monzonite (Kilkis, Northern Greece) has been examined in some detail by means of chemical and X-ray analysis.

The studied stillbite belongs to the "Intermediate Zeolites" with $R = (Si / Si + Ti + Al + Fe^{3+}) = 0.747$ and has the following chemical formula:

$$K_{0.41}Na_{0.35}Ca_{3.89}Mg_{0.25}Sr_{0.09}Al_{9.09}Si_{26.87}O_{72}25.8H_2O.$$

The crystallographic parameters of the Sr-bearing stilbite and other seven different stilbites have been correlated with their chemical composition. The main influence to the cell parameters a and c of the stilbite comes from the monovalent cations Na⁺ and K⁺. The Sr-bearing stilbite in the Vathi quartz-monzonite found associated with pyrite, calcite and quartz, has crystallized from hydrothermal solutions of low temperature, not exceeding 200 °C.

Keywords: Zeolites, Sr-bearing stilbite, quartz-monzonite, crystallography and chemistry relations, Serbomacedonian massif, Northern Greece.

Introduction

Stilbite occurs commonly in association with other zeolites, in different rock types. It is found in sedimentary rocks (e.g. TSCHERNICH, 1972; STEPHENS and BRAY, 1973; NAKAMUTA, 1976), gneisses (e.g. PRIVETT, 1977), pegmatites (e.g. TENNYSON, 1960; ZABINSKI 1975), serpentinites (e.g. CHAUDHARI and SHARMA, 1971; AR-GENTI et al., 1986), basalts (e.g. MASON and GREENBERG, 1954; WALKER, 1960; AUMENTO, 1966; COMIN-CHIARAMONTI et al., 1979), gabbros (e.g. KIMATA et al., 1979) and granitic rocks (e.g. RAADE, 1969; PRIVETT, 1977).

Stilbite is quite similar to stellerite and barrerite in its X-ray diffraction trace. The most common of the three minerals is stilbite. The structure of stilbite was solved by GALLI and GOTTARDI (1966) and refinements were carried by SLAUGHTER (1970) and GALLI (1971). Usually in natural stilbites the two sites (1 and 1P) are filled by Ca, and Na occupies in part one site (2P).

The SrO content of stilbites was always found very low, not exceeding 0.06 wt% (GOT-TARDI and GALLI, 1985). In only one case a SrO content of 0.28 wt% is mentioned for a Hungarian stilbite (REICHERT and ERDELYI, 1935). The present work describes the mode of occurrence, the mineralogic properties and the genesis of a Sr-bearing stilbite in a quartz-monzonite from the Vathi area (N. Greece).

Occurrence

The investigated area of Vathi belongs geotectonically to the Serbomacedonian massif (Fig. 1) and specifically to the Vertiskos series (KOCKEL et al., 1971). This series is composed

¹ Department of Mineralogy-Petrology-Economic Geology, Aristotle University of Thessaloniki, GR-54006 Thessaloniki, Greece.

² Institute of Geology and Mineral Exploration (I.G.M.E.), Frankon 1, GR-54626 Thessaloniki, Greece.

of metamorphic rocks comprising biotite-muscovite, biotite-hornblende gneisses and micaschists with concordant intercalations of amphibolites and ultramafic rocks. These metamorphics are of at least lower paleozoic age (KOCKEL and WALTHER, 1968). Several volcanic bodies of rhyolitic to andesitic composition are extruded on the metamorphics. Subvolcanic to plutonic stock-like bodies, plugs and dykes ranging in composition from monzonite-monzodiorite to quartz monzodiorite and dacite-rhyodacite porphyries are also intruding the metamorphics. The magmatism in this area is considered to be of Upper Pliocene to lower quaternary age (MELIDONIS, 1972; KOCKEL et al., 1977; PANAGOS et al., 1978).

Drilling through an intrusive body (Fig. 1), which is surrounded by serpentinized ultramafic rocks, partly altered to talc and/or magnesite ceased at 504 m depth. The study of thin sections of the recovered drill core revealed that the intrusive has the following mineralogic assemblage:

quartz, K-feldspar, plagioclase (An=35%), biotite, Ca-amphibole, apatite, sphene, opaques±chlorite±calcite.

Chemical analysis on representative samples of the intrusive rock (table 1) gave the composition of a quartz-monzonite, according to STRECKEISEN and LE MAITRE (1979) classification. The quartz-monzonite shows a strong potassic alteration and porphyry copper type mineralization, consisting of pyrite, chalcopyrite and traces of molybdenite and galena. These phenomena are connected with the latest stages of the same magmatic event which gave the quartz-monzonite.

Stilbite was found to be more widespread between 52 and 55 m depth, from which the studied samples have been collected.



Fig. 1 Geological map of the investigated area.

SiO ₂ wt%	63.32
Ti0 ₂	0.35
A1 ₂ 0 ₃	17.58
Fe ₂ 03	1.13
Fe0	1.91
Mn0	0.05
MgO	2.57
CaO	3.28
Na ₂ 0	4.00
к ₂ 0	5.60
P205	tr.
s0 ₃	0.87
Total	100.66

Tab. 1 Chemical composition of the quartz-monzonite from Vathi, Kilkis (N. Greece).

Average of 3 analyses. Mixed sample from representative rock-samples selected from the drill-core (depth 180-240 m).

It occurs in veinlets (up to 1 cm wide), fractures and open spaces (up to 10 cm wide) of the quartz-monzonite. In vein and fracture-fillings, stilbite develops granular masses or interlocking small crystals, while in open spaces it develops radiating aggregates of tabular crystals (up to 8 mm long), which are actually cruciform twins. The aggregates are grouped in nearly parallel position to form sheaf-like aggregates (Fig. 2). Thin section studies revealed that stilbite shows good cleavage and its characteristic twinning in polycrystalline aggregates. It is creamy yellow in colour with brown tint and occurs associated with calcite, quartz, pyrite (Fig. 3) and secondary alkalifeldspar and biotite.

X-ray data

X-ray powder diffraction traces were taken with a Philips diffractometer, Ni-filtered CuK α radiation, using silicon as internal standard. The scanning speed was 1/1° and 1/4° per minute over the interval 4–104° of 20. The Xray pattern of Sr-bearing stilbite was carefully compared with the X-ray patterns of stilbite, stellerite and barrerite. The X-ray pattern differences between stilbite and stellerite indexed



Fig. 2 Scanning-electron microphotograph of Vathi Sr-bearing stilbite. Bar scale = $100 \,\mu$ m.



Fig. 3 Scanning-electron microphotograph of euhedral pyrite crystal as inclusion in the Sr-bearing stilbite. Bar scale = $71 \,\mu m$.

	Wt.	%	<u>Ca</u> 1	culated t	:o 72(0)		Cell parame	ters
Sample No	• V1	٧2		٧1	٧2		V1	V2
S10 ₂	57.29	57.57	Si	26.904	26,839	a (Å)	13.607(6)	13.601(5)
Ti02	0.02	bdl	Ti	0.007	-	ь (Å)	18.259(8)	18.252(7)
A1203	16.34	16.62	A1	9.044	9.132	c (Å)	11.286(5)	11.282(4)
Fe ₂ 0 ₃	0.01	0.02	· Fe ³	+ 0.004	0.007	v (Å ³)	2210.7	2208.8
MgŐ	0.33	0.38	Mg	0.231	0.264	β	127 ⁰ 96′	127 ⁰ 94′
CaO	7.69	7.81	Ca	3.869	3.901			
Sr0	0.31	0.34	Sr	0.084	0.092			
BaO	0.02	0.01	Ba	0.004	0.002			
NiO	0.03	0.03	Ni	0.011	0.011			
Na ₂ 0	0.35	0.41	Na	0.319	0.371			
K20	0.83	0.53	К	0.497	0.315			
H20*	16.78	16.28	H ₂ 0	26.28	25.31			
۲.			z	35.959	35.978			
Total	100.00	100.00	Х	5.015	4.956			
			R	0.748	0.746			

Tab. 2 Microprobe analyses and cell parameters of the Sr-Bearing stilbite from Vathi, Kilkis (N. Greece).

V1) Average of 3 analyses. V2) Average of 7 analyses. *) Estimated by difference. bdl = below detection limit. $Z = Si + Ti + Al + Fe^{3+}$. X = Mg + Ca + Sr + Ba + Ni + Na + K. $R = Si/(Si + Ti + Al + Fe^{3+})$. Figures in parentheses represent the estimated standard deviation; thus 13.607 (6) indicates estimated standard deviation of 0.006.

by PASSAGLIA et al. (1978) and GOTTARDI and GALLI (1985) were taken into consideration.

Stilbite is monoclinic with crystal symmetry C2/m (SLAUGHTER, 1970; GALLI, 1971), while stellerite and barrerite are orthorhombic with crystal symmetries Fmmm and Amma respectively (GALLI and ALBERTI, 1975). Natural stilbite crystals showing orthorhombic and monoclinic growth sectors have been mentioned by AKIZUKI and KONNO (1985).

The X-ray powder lines which are single in the orthorhombic system, split up in the monoclinic one (PASSAGLIA et al., 1978). In the most appropriate interval between 23° and 24° of 20 (CuK α -radiation), the studied stilbite exhibits two distinct peaks of the same height. According to PASSAGLIA et al. (1978) this is characteristic for monoclinic stilbite.

Cell parameter refinements (Table 2) on the Sr-bearing stilbite were performed with X-ray

powder diffraction data, using the 18 strongest and welldefined reflections and Si as internal standard. The values of the cell dimensions of the studied stilbite are close to those reported by other workers (e.g. GALLI and GOTTARDI, 1966; SIMONOT-GRANGE et al., 1970).

Chemistry of the Minerals

Microprobe analyses of Sr-bearing stilbite and some of the associated minerals: alkalifeldspar, biotite and pyrite are given in Tables 2, 3 and 4.

Stilbite from Vathi contains considerable amounts of Sr (on average 0.33 wt% SrO), which to our update knowledge is the highest Sr-content reported in literature. The major cation in the Vathi stilbite is Ca but considerable amounts of K, Mg and Na are present. The

	Wt <u>%</u>		Calc.	on 8(0)	11(0)
Sample No	٧2	V1		٧2	٧1
Si0 ₂	64.97	37.22	Si	2.990	2.818
TiO2	0.02	4.68	A1	1.007	1.121
A1203	18.56	12.56	Ti	0.001	0.267
FeÖ*	0.15	14.87	Fe ²⁺	0.006	0.942
Mn0	n.a.	0.16	Mn	-	0.010
MgO	0.07	15.16	Mg	0.005	1.711
CaO	bd1	n.a.	Sr	0.016	-
Sr0	0.61	n.a.	Ba	0.003	-
BaO	0.16	n.a.	Na	0.111	0.029
NiO	bdl	n.a.	К	0.846	0.942
Na ₂ 0	1.24	0.20	se Or	88.4	-
к,0	14.40	9.75	Ab	11.6	-
£			¥≊ An	0.0	-
Total	100.18	94.60			

Tab. 3 Microprobe analyses of alkali-feldspar (V2) and biotite (V1) from Vathi, Kilkis (N. Greece).

V2) Average of 3 analyses. V1) Average of 3 analyses.
*) FeO as total iron. bdl = below detection limit. n.a.
= not analysed.

chemical parameter $R = Si/(Si+Ti+Al+Fe^{3+})$, which represents the percentage of the tetrahedra occupied by Si, varies between 0.746 and 0.748. Based on the R values, the investigated stilbite is classified as "Intermediate zeolite" (0.625 < R < 0.750) after GOTTARDI (1978). The calculation of cation proportions on 72 oxygen atoms (Table 2) gives the formulae (average of samples V1 and V2):

 $\begin{array}{c} K_{0.41}Na_{0.35}Ca_{3.89}Mg_{0.25}Sr_{0.09}Al_{9.09}\\Si_{26.87}O_{72}25.8H_2O\end{array}$

Tab. 4	Microprobe	analyses	of	pyrite	from	Vathi,
Kilkis (N	N. Greece).					

Wt.%			Ca	Calc. on 2(S)			
Sample No	5 V1		4	٧1			
Cu	0.04		Со	0.01			
Ni	0.03		Fe	0.99			
Co .	0.43		S	2.00			
Fe	46.01		Σ	3.00			
S	53.56		Formula	Fe _{0.99} Co _{0.01} S ₂			
Total	100.07		···· • •··· • • • • • • • • • • • • • •				

V1) Average of 3 analyses.

PASSAGLIA et al. (1978), using the chemistry of many stellerites and stilbites on a triangular mole plot (Fig. 4), distinguish two separate fields for the minerals. Using the chemical data of table 2 and the triangular mole plot of PASSAGLIA et al. (1978), we see that the Vathi Sr-bearing stilbite falls in the stilbite area of Figure 4.

The alkali-feldspar ($Or_{88.4}Ab_{11.6}An_{0.0}$) contains considerable amounts of SrO (0.6 wt%).

The associated mica (Table 3) is a common tri-octahedral mica, namely biotite. Partial microprobe analyses on calcite gave CaO = 47.5, MgO = 0.1, FeO=0.0 and SiO₂ = 0.1 wt%.

The euhedral crystals of pyrite (Fig. 3) in association with the stilbite, contain around 0.4 wt% Co (Table 4).

Relationship between crystallographic parameters and chemical composition.

In this study the crystallographic parameters a, b, c, V and β of different stilbites have been correlated with their chemical composition. The following chemical variables were chosen:

 $Z = Si + Ti + Al + Fe^{3+},$

 $R = Si/(Si+Ti+Al+Fe^{3+}),$

X = Mg+Ca+Sr+Ba+Ni+Na+K,

M = Na + K (monovalent cations),

D = Ca+Mg+Sr+Ba+Ni (divalent cations),

Na, K, Z/X and M/X.

The correlation diagrams have been made using data of nine different stilbites from various places (see footnotes to Fig. 5).

The correlations of the crystallographic parameters b, V and β with the chemical composition show broad scattering. The crystallographic parameters a, b, c, V and β seem not to be effected by the divalent cations (Fig. 5).

The correlations a - (X, M, M/X, Z/X)and c - (X, M, M/X, Z/X) seem to be the best ones. Both parameters a and c are positively correlated with X, M, M/X and negatively with Z/X (Figs. 6 and 7). Thus, the main influence to the cell parameters a and c comes from the monovalent cations (M), these two parameters are increasing with increasing M (Figs. 6 and 7).

The correlations of β -angle with the chemi-



Fig. 4 The triangular mole plot $D_{3.7}Al_{7.4}Si_{28.6.}O_{72} - D_{4.8}Al_{9.6}Si_{26.4}O_{72} - M_{2.2}D_{3.7}Al_{9.6}Si_{26.4}O_{72}$ (after PASSAGLIA et al. 1978). The dashed line separates the area of stellerites and stilbites. 1) stilbite from Vathi, sample V1, 2) stilbite from Vathi, sample V2, M = Na+K (monovalent cations), D = Ca+Mg+Sr+Ba (divalent cations).

cal variables were not so discernible as in figures 6 and 7. Possibly the β -angle is correlated positively with the monovalent cations (Fig. 8). According to PASSAGLIA et al. (1978) the β -angle depends on the monovalent cations only, and K influences the β -angle in the same direction but less markedly than Na. The single correlations between cell parameters, Na and K show broad scattering except those of *a* with Na and K, which show that *a* increases with increasing Na and K content of stilbite (Fig. 8). These single correlations are based on data from fewer stilbites because some of them do not contain any Na or K.

Origin of the Sr-bearing stilbite

The mode of occurrence (veinlets, fracture and open space fillings) of the studied stilbite within the quartz-monzonite indicate an epigenetic origin. The association of the mineral with pyrite, calcite and quartz as well as the potassic alteration (secondary K-feldspar and biotite) and the porphyry copper mineralization of the quartz-monzonite suggest a hydrothermal origin.

The Sr content of the stilbite (0.31–0.34 wt% SrO, Table 2) derived mainly by hydrothermal activity on the Sr-bearing alkali-feldspars of



Fig. 6 Plot of a versus composition in stilbite. X=Mg+Ca+Sr+Ba+Ni+Na+K (divalent and monovalent cations), M=Na+K (monovalent cations), $Z=Si+Ti+Al+Fe^{3+}$. Stilbites used: see footnotes to Fig. 5.



Fig. 7 Plot of c versus composition in stilbite. For abbreviations and symbols see footnotes to Figs 5 and 6.



LIOU (1971). HARADA and TOMITA (1967) showed that stilbite above $200 \,^{\circ}$ C and water pressure >40 atms produced wairakite. A stability diagram for the above mentioned reactions is presented in Figure 9.

In the Vathi quartz-monzonite no Ca-analcime, laumonite, wairakite or other zeolites were observed. From the above it is concluded that the studied Sr-bearing stilbite has originated from a late stage hydrothermal activity, at temperature not exceeding 200 °C.

^a Fig. 8 Plot of β and a versus composition in stilbite. For abbreviations and symbols see footnotes to Figs 5 and 6.



Fig. 9 P-T fields for stilbite. (a) stilbite = laumontite + quartz + H_2O (LIOU, 1971); (b) stilbite = Ca-analcime + quartz + H_2O (JUAN and LO 1973); (c) stilbite = wairakite + SiO_2 + H_2O (HARADA and Томіта, 1967).

Acknowledgements

We would like to express our thanks to Drs: G. Christofides, A. Kassoli-Fournaraki, A. Tsirambidis and K. Christodoulou for critial comments on the manuscript. We express our gratitude to Mrs. V. Lymberopoulou for typing the manuscript, to Mrs I. Chalvatzi for drawing the figures, to H. Harryson (Inst. of Geology, Univ. of Uppsala, Sweden), who performed the microprobe measurements, to L. Liung (biomedical Center, Univ. of Uppsala, Sweden), who helped us with the scanning electron microscope.

References

- ABBONA, F. and FRANCHINI, A.M. (1970): Sulla disidratazione della stilbite. Atti Accad. Sci. Torino, 104, 309-321.
- AKIZUKI, M. and KONNO, H. (1985): Order-disorder structure and the internal texture of stilbite. Amer. Mineral., 70, 814-821.
- ARGENTI, P., LUCHETTI, G. and PENCO, A.M. (1986): Zeolite-bearing assemblages at the contact Voltri Group and Sestri-Voltagio zone (Liguria, Italy). N. Jb. Miner. Mh., 5; 229-239.

- AUMENTO, F. (1966): Thermal transformations of stilbite. Can. J. Earth Sci., 3, 351-366.
- CHAUDHARI, M. W. and SHARMA, B. L. (1971): On stilbite from Ranakpur. Journ. Geol. Soc. India, 12, 191-192.
- COMIN-CHIARAMONTI, P., PONGILUPPI, D. and VEZ-ZALINI, G. (1979): Zeolites in shoshonitic volcanics of the north-eastern Azerbaijan (Iran). Bull. Mineral., 102, 386-390.
- GALLI, E. (1971): Refinement of the crystal structure of stilbite. Acta Cryst., B27, 833-841.
- GALLI, E. and GOTTARDI, G. (1966): The crystal structure of stilbite. Miner. Petrogr. Acta, 12, 1-10.
- GALLI, E. and ALBERTI, A. (1975): The crystal structure of barrerite. Bull. Soc. Fr. Minéral. Cristallogr. 98, 331-340.
- GOTTARDI, G. (1978): In Natural Zeolites: occurrence, properties, use (Sand, L.B. & Mumpton, F.A., ed.). Oxford and New York, Pergamon Press, 31-44.
- GOTTARDI, G. and GALLI, E. (1985): Natural Zeolites.
- Berlin, Springer-Verlag, 284-300. HARADA, K. and TOMITA, K. (1967): A sodian stilbite from Onigajo, Mie prefecture, Japan, with some experimental studies concerning the conversion of stilbite to wairakite at low water vapor pressures. Amer. Mineral., 52, 1438-1450.
- JUAN, V.C. and Lo, H.-J. (1973): Stability field of stilbite. Geol. Soc. China (Formosa), Proc., 16, 37-49
- KIMATA, M., MATSUKURA, Y. and VATSU, E. (1979): Stilbite in the weathered gabbro. Ann. Rep. Inst. Geosci. Univ. Tsukuba, 5, 62-64.
- KOCKEL, F. and WALTHER, H. (1968): Zur geologischen Entwicklung des südlichen Serbo-Ma-zedonischen Massivs (Nordgriechenland). Bul-gar. Acad. Science. Commit. Geol., KH (vol.) XVII, Sofia.
- KOCKEL, F., MOLLAT, H. and WALTHER, H.W. (1971): Geologie des Serbo-Mazedonischen Massivs und seines mesozoischen Rahmens (Nord-Griechenland). Geol. Jb., 89, 529-551.
- KOCKEL, F., MOLLAT, H. and WALTHER, H.W. (1977): Erläuterungen zur Geologischen Karte der Chalkidiki und angrenzender Gebiete, «Bund. Geow. Rohst.», Hannover, 119 p. LIOU, J.G. (1971): Silbite-Laumontite equilibrium.
- Contr. Mineral. Petrol., 31, 171-177.
- MASON, B. and GREENBERG, S.S. (1954): Zeolites and associated minerals from southern Brazil. Arkiv Min. Geol., 1, 519–526.
- MELIDONIS, N.G. (1972): Der Bogen junger vulkanischer Gesteine zwischen Strymonikon und Metamorphosis (Zentr. Mazedonien). Mineral Exploration, I.G.M.E. No 5, 1-51 (in Greek).
- NAKAMUTA, Y. (1976): Zeolites and clay minerals in the Tertiary sedimentary rocks in Tsuyazaki area, Fukuoka Prefecture. J. Japan. Assoc. Min. Petr. and Econ. Geol., 71, 147-156. (Japanese with English summary)
- PANAGOS, A.G., PE, G.G. and VARNAVAS, S.P. (1978): The volcanic rocks of Strymonikon-Metamorphosis, Central Macedonia, Greece. Chem. Erde, 37, 50-61.
- PASSAGLIA, E., GALLI, E., LEONI, L. and ROSSI, G. (1978): The crystal chemistry of stilbites and stellerites. Bull. Mineral., 101, 368-375.

- PRIVETT, D.R. (1977): An unusual octahedral fluorite, stilbite, laumontite, calcite, and quartz assemblage in Danville, Virginia. Virginia Minerals, 23, 7-9.
- RAADE, G. (1969): Cavity minerals from the Permian biotite granite at Nedre Eiker church. Norsk. Geol. Tidsskr., 49, 227-239.
- REICHERT, R. and ERDELYI, J. (1935): Über die Minerale des Csodi-Berges bei Dunabogdany (Ungarn). Tscherm. Miner. Petr. Mitt., 46: 237-255.
- SIMONOT-GRANGE, M.H., COINTOT, A. and THRIERR-SOREL, A. (1970): Etude des systèmes divariants zéolite-eau. Cas de la stilbite. Comparaison avec la heulandite. Bull. Soc. Chim. Fr. 1970, 4286-4297.
- SLAUGHTER, M. (1970): Crystal structure of stilbite. Amer. Mineral., 55, 387-397.
- STEPHENS, J.D. and BRAY, E. (1973): Occurrence and infrared analysis of unusual zeolitic minerals from Bingham, Utah. Mineral. Record, 4, 67-72.
- STRECKEISEN, A. and LE MAITRE, R.W. (1979): A chemical approximation to the modal QAPF

classification of the igneous rocks. N. Jb. Miner. Abh., 136, 169-206.

TENNYSON, C. (1960): Berylliummineralien und ihre pegmatitische Paragenese in den Graniten von Tittling/Bayerischer Wald. N. Jb. Miner. Abh., 94: 1253-1265.

TSCHERNICH, R.W. (1972): Zeolites from Skookumchuck dam, Washington. Mineral. Record, 3, 30 - 34.

- WALKER, G.P.L. (1960): The amygdale minerals in the Tertiary lavas of Ireland. III. Regional distribution. Mineral. Mag., 32, 503-527. WISE, W.S. (1969): Heulandite with excess water: A
- correction. Mineral Mag., 37, 522.
- ZABINSKI, W. (1975): Stilbite from Strzegom (Lower Silesia). Mineral. Polonica, 6, 93-98.

Manuscript received September 10, 1987; revised manuscript accepted February 2, 1988.