

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 69 (1989)

Heft: 1

Artikel: Mantle periodites from the Austroalpine Mt. Mary nappe (Western Alps)

Autor: Cesare, B. / Martin, S. / Zaggia, L.

DOI: <https://doi.org/10.5169/seals-52778>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Mantle peridotites from the Austroalpine Mt. Mary nappe (Western Alps)

by B. Cesare¹, S. Martin² and L. Zaggia³

Abstract

A possible fragment of the Insubric continental upper mantle occurs as a slice of spinel - amphibole - harzburgite within the intermediate unit of the Austroalpine Mt. Mary nappe.

This contribution reports petrographic data as well as electron microprobe analyses of the mineral constituents, and major, trace and Rare Earth element abundances of the bulk rock.

The data are consistent with a metamorphic reequilibration of a former spinel lherzolite under pre-Alpine high temperature amphibolite/granulite facies conditions, followed by partial serpentinization during the Alpine orogeny.

Keywords: Austroalpine, Italian Western Alps, Mt. Mary nappe, peridotite, metamorphism, mineral chemistry, bulk rock composition.

1. Introduction

This paper records the first occurrence of spinel - amphibole - harzburgites in the Austroalpine continental crust of the Mt. Mary nappe, Italian Western Alps.

The peridotitic rocks form a small lens located at 2610m on the southern slope of the Mt. Mary - P.te de Genève ridge, between the Valpelline and Aosta Valleys (Fig. 1). They are tectonically associated with the kinzigitic paragneisses of the Intermediate unit of Mt. Mary nappe, which exhibit a polymetamorphic history and widespread mylonitic deformation (CANEPA et al., 1987 and in press). Additional fragments of serpentinized peridotites occur in the rock debris on the northern side of Mt. Mary (upper Comba Verzignoletta and Comba d'Arpisson).

Similar associations of peridotite tectonites with high-grade paragneisses have already been reported from other units of the Austroalpine nappe system; they occur for instance in the Upper unit of the Sesia-Lanzo (2nd Diorite-Kinzigitic zone; FRANCHI, 1903; BECCALUVA et al., 1978) and Dent Blanche (Valpelline unit; ARGAND, 1934; NICOT, 1977) composite nappes.

In the Southern Alps, spinel lherzolites occur as isolated slices associated to the lower continental

crust of the Ivrea-Verbano unit (Baldissero, Finero and Balmuccia peridotites, VOSHAGE et al., 1987 and refs. therein). All the above ultramafic rocks have been interpreted as "continental peridotites". They are believed to represent relics of lithospheric, continental, uppermost mantle.

In contrast, the largely preserved spinel and plagioclase peridotites of the Lanzo massif (POGNANTE et al., 1985) and of Balangero (SANDRONE and COMPAGNONI, 1986), which experienced the eo-Alpine HP-LT metamorphism, could represent fragments of the Tethyan oceanic lithosphere.

2. Geological Setting

The Austroalpine Mt. Mary nappe is located between the Aosta, Valpelline and Valtournanche valleys. In the earlier literature, the Mt. Mary nappe was defined as a single nappe consisting of the Arolla and Valpelline series (ARGAND, 1906) or else as a composite nappe formed by two sliding tectonic units ("Gleitbretter"; DIEHL et al., 1952). The Mt. Mary nappe tectonically overlies the Piedmont ophiolite system and in turn is overthrust by the Dent Blanche composite nappe. The Mesozoic

¹Dipartimento di Mineralogia e Petrologia, Università di Padova, Corso Garibaldi, 37, I-35122 Padova.

²Dipartimento di Geologia, Paleontologia e Geofisica, Università di Padova, via Giotto, 1, I-35137 Padova.

³C.N.R. Istituto di Geologia applicata, Corso Stati Uniti, 4, I-35100 Padova.




Fig. 1 Tectonic sketch-map of the Italian North-Western Alps.

1) Austroalpine Dent Blanche (DB) and Sesia-Lanzo (SL) composite nappe system, including Mt. Emilius (E), Glacier-Rafay (GR), Tour Ponton (TP), Pillonet (P), Santanel (S) and Verres (VE) Klippen: a) Upper element, b) Lower element. 2) Austroalpine Mt. Mary composite nappe: a) Lower unit; b) Intermediate kinzigitic unit; c) Upper unit. 3) Tethyan ophiolites. 4) Upper Penninic Monte Rosa (MR), Gran Paradiso (GP) and Arcesa-Brusson (AB) nappes; 5) Middle Penninic Gr. St. Bernard nappe. 6) Location of Mt. Mary peridotite body. ARF: Aosta-Ranzola fault.

sequence of the Roisan zone, located between the Mt. Mary and Dent Blanche basement units, was interpreted as the sedimentary cover of the Mt. Mary unit (MASSON, 1938; DIEHL et al., 1952; ELTER, 1960).

Recent studies in this area (CESARE, 1987; ZAGGIA, 1987; CANEPA et al., 1987; in press), reveal that the internal structural setting of Mt. Mary needs to be revised by introducing an uppermost third unit. This unit is composed of a pervasively sheared cover and basement series (Fig. 1). Therefore, the composite nappe of Mt. Mary consists of three tectonic units which are distinguished by different lithological compositions and are separated by meter-size mylonitic horizons.

From bottom to top the three units are:

i) The Lower unit (ARGAND's Arolla Series), which is composed of high-grade paragneisses of unknown age (Hercynian or older) and of Permo-Carboniferous granitoids; they suffered a pervasive Alpine metamorphic overprint under glaucophane-bearing greenschist facies conditions;

ii) The Intermediate unit (ARGAND's Valpel-line Series) which consists of high-grade kinzigitic paragneisses, metabasites and marbles, locally overprinted by an Alpine retrogression of undefined age. The high temperature granulite/amphibolite metamorphism is supposed to be of Hercynian (or older) age by comparison with the Dent Blanche nappe and the Ivrea zone (DAL PIAZ et al., 1972; HUNZIKER, 1974; HUNZIKER & ZINGG, 1980).

iii) The Upper unit, which overthrust both the underlying units and is predominantly composed of mylonites derived from locally preserved Triassic-Cretaceous cover series (Roisan Zone) and from granitoid rocks. This unit exhibits an intense Alpine metamorphism under greenschist facies conditions.

3. Spinel peridotite of Mont Mary

3.1. PETROGRAPHIC DESCRIPTION

The spinel - amphibole - harzburgite of Mt. Mary displays a high-grade metamorphic fabric and is composed of olivine, orthopyroxene, amphibole, green spinel and minor magnetite. Olivine, orthopyroxene (locally with clinopyroxene exsolution lamellae) and amphibole are the most abundant phases; green spinel and magnetite occur as inclusions in orthopyroxene and as interstitial grains. The rock is fine grained, although the prophyroblastic orthopyroxene, often including olivine and amphibole, may reach 4-5 mm in length. The microstructural relationships indicate an equilibrium texture between olivine, orthopyroxene,

spinel and amphibole (the latter possibly derived from a former clinopyroxene); reaction rims or pseudomorphic replacements have not been observed.

The local development of chlorite-rims around interstitial spinel is attributed to a subsequent reequilibration during which amphibole remained stable. Finally all the major mineral phases are strongly deformed (undulose extinction) and anastomosing micro-shear zones developed. The latter are marked by the transformation of olivine to serpentine and magnetite, which has also been observed inside microcracks. The serpentinization postdates the chlorite growth.

3.2. MINERAL CHEMISTRY

Microprobe analyses were obtained from all major mineral phases using the ARL-SEM microprobe of C.N.R. at the Department of Earth Sciences of Milano University; they are given in Tab.

Tab. 1 Electron microprobe analyses of pyroxenes, olivine and amphiboles from harzburgite M157 of Mont Mary

	Opx		Olivine		Amphibole	
SiO ₂	53.76	54.18	39.03	39.21	46.75	47.01
TiO ₂	n.d.	n.d.	n.m.	n.m.	0.06	0.09
Al ₂ O ₃	2.94	2.82	n.d.	n.d.	11.02	11.27
Cr ₂ O ₃	0.08	0.08	n.m.	n.m.	0.23	0.22
FeO*	10.55	10.14	14.94	15.18	5.45	5.31
MnO	0.35	0.30	0.28	0.28	0.16	0.11
MgO	31.93	32.19	45.79	45.27	19.17	19.44
CaO	0.38	0.29	n.d.	0.02	12.68	13.20
Na ₂ O	n.d.	n.d.	n.m.	n.m.	1.31	1.47
K ₂ O	n.m.	n.m.	n.m.	n.m.	0.25	0.22
Total	99.99	100.00	100.04	99.96	97.08	98.34
Cations						
Si	1.898	1.906	0.982	0.987	6.652	6.610
Ti	0.000	0.117	0.000	0.000	0.006	0.009
Al	0.122	0.117	0.000	0.000	1.848	1.867
Cr	0.002	0.002	0.000	0.000	0.025	0.024
Fe	0.311	0.298	0.315	0.320	0.648	0.624
Mn	0.010	0.008	0.006	0.006	0.019	0.013
Mg	1.680	1.688	1.714	1.699	4.066	4.075
Ca	0.014	0.010	0.000	0.001	1.933	1.988
Na	0.000	0.000	0.000	0.000	0.361	0.400
K	0.000	0.000	0.000	0.000	0.045	0.039
Total	4.037	4.029	3.017	3.013	15.603	15.649
x _{Mg} **	0.843	0.849	0.845	0.841	0.862	0.867

*Total Fe as FeO

**x_{Mg} = Mg/(Mg+Fe), where Fe is total iron

Tab. 2 Electron microprobe analyses of spinels, chlorites and serpentines from harzburgite M157 of Mont Mary.

	Spinel		Chlorite		Serpentine	
SiO ₂	n.d.	n.d.	31.97	32.57	40.46	44.32
Al ₂ O ₃	57.95	57.96	17.46	15.79	2.34	0.33
Cr ₂ O ₃	5.11	4.29	n.m.	n.m.	n.m.	n.m.
FeO*	19.38	19.54	3.15	3.82	1.45	3.86
MnO	0.24	0.22	n.d.	0.05	0.01	0.07
MgO	16.86	16.93	36.22	36.91	43.65	41.99
CaO	n.m.	n.m.	n.d.	n.d.	n.d.	n.d.
Total	99.54	99.94	88.81	89.14	87.91	90.40
Cations						
Si	0.000	0.000	5.911	6.027	1.872	1.999
Al	1.799	1.806	3.808	3.445	0.128	0.018
Cr	0.106	0.089	0.000	0.000	0.000	0.000
Fe ^{3+***}	0.095	0.104	0.000	0.000	0.000	0.000
Fe ²⁺	0.333	0.330	0.487	0.591	0.056	0.146
Mn	0.005	0.005	0.000	0.008	0.000	0.000
Mg	0.662	0.667	9.980	10.180	3.009	2.829
Ca	0.000	0.000	0.000	0.000	0.000	0.000
Total	3.000	3.001	20.186	20.251	5.065	4.992
x _{Mg} ^{***}	0.607	0.606	0.953	0.945	0.982	0.951

*Total Fe as FeO

**Fe³⁺ in the spinel formula was recalculated with the charge balance method

***x_{Mg} = Mg/Mg+Fe, where Fe is total iron

1 and 2. The applied correction program was MAGIC IV of COLBY (1972).

Olivine exhibits a uniform composition of Fo 0.84 (chrysolite).

Orthopyroxene is a Mg-rich bronzite (En 0.84÷0.85) with moderate XAl₂O₃ (2.9 wt%) and low Cr₂O₃ (0.08 wt%) and CaO (0.3 wt%) contents. TiO₂ and Na₂O contents have not been detected.

Green spinel is aluminous and contains less Cr₂O₃ (4.3÷5.1 wt%) than the ones reported by ERNST (1978) from the peridotites of the Western Alps (often around 9.1÷13.1. wt%).

Amphibole is a Mg-hornblende according to the scheme proposed by LEAKE (1978) and is characterized by very low TiO₂ contents. The XMg (XMg = Mg/Mg+Fe, where Fe is total iron) of the amphibole is 0.87; it is similar but slightly higher than the corresponding values for olivine and orthopyroxene (≈ 0.85). This relationship is in agreement with the

$$Kd_{\text{Fe-Mg}}^{\text{ol-amph}} - \text{values}$$

reported by ROBINSON et al. (1982). On a Na+K (A-site) versus Si diagram, the Mg-hornblendes of the Mt. Mary harzburgite plot at the upper end of

ROBINSON et al.'s (1982, Fig. 46, p. 101) data points from olivine-orthopyroxene-Al-spinel ± chlorite bearing ultramafites, close to the array of chlorite-free assemblages.

Chlorite occurring as overgrowth rims on Al-spinel is Mg-rich (clinocllore).

3.3. BULK ROCK GEOCHEMISTRY

Major and trace element concentrations of sample M157 have been obtained by X-ray fluorescence analysis at the C.N.R. laboratory of Department of Mineralogy and Petrology of Padova University and are reported in Tab. 3. Rare Earth element (REE) concentrations, analysed by ICP spectrometry at the Laboratoire de Spectrochimie of Nancy, are listed in Tab. 3.

The major element composition of the spinel-amphibole-harzburgite from the Mt. Mary nappe reveals some remarkable differences with the other peridotites from the Western Alps: the Mt. Mary harzburgite shows anomalously high Al₂O₃, Fe₂O₃, CaO and Na₂O contents, and rather low MgO content with respect to the mean values of the peridotites from the Western Alps (ERNST, 1978). These data may be explained by the large abundance of green spinel and amphibole in the investigated samples.

The minor element concentrations, in particular Ni and Cr, are lower than those reported from the Western Alps' peridotites (OTTONELLO et al., 1984). They are, however, very similar to the values

Tab. 3 Bulk rock analysis of major elements*, transitional elements** and REE*** of harzburgite M157 of Mont Mary.

Oxide	wt.%	Element	ppm	Element	ppm
SiO ₂	42.35	Y	5	La	0.50
TiO ₂	0.29	Zr	13	Ce	3.30
Al ₂ O ₃	6.91	Nb	-	Nd	1.68
Fe ₂ O ₃	11.35	Zn	52	Sm	1.04
MnO	0.18	Cr	1540	Eu	0.26
MgO	29.57	Sr	8	Gd	0.99
CaO	4.00	Rb	-	Dy	1.32
Na ₂ O	0.27	Ni	1219	Er	0.79
K ₂ O	0.11	Ba	-	Yb	0.89
P ₂ O ₅	0.02			Lu	0.12
L.O.I.	4.23				
Total	99.28				

Analytical methods:

*Atomic absorption, wet-method

**X-ray fluorescence spectrometry

***Inductively coupled plasma-atomic spectrometry




Fig. 2 Chondrite-normalized (normalizing values used were those of EVENSEN et al., 1978) REE composition of Mt. Mary peridotite (1), in comparison with the composition patterns of subcontinental peridotites from Baldissero (2), Finero (3) and Alpe Arami (4) (OTTONELLO et al., 1984 and references therein.).

obtained for the pyroxenite body of Mt. Emilius Klippe (BENCIOLINI, 1988).

The chondrite-normalized REE concentrations of the Mt. Mary harzburgite (Fig. 2) show a flat REE-distribution pattern with an average mean value of approximately 5 times chondritic. They differ considerably from the trends observed for similar lithologies from the Alpe Arami, Baldissero, Finero and Balmuccia peridotite bodies, which show lower concentrations and marked light REE depletion (OTTONELLO et al., 1984). The depletion of the light REE relative to the heavier REE has been interpreted as the result of a previous partial melting and melt extraction events.

4. Metamorphic evolution

Temperature estimates of (peak?) metamorphic conditions of the Mt. Mary harzburgite were obtained from different calibrations of the olivine-spinel Fe-Mg partition geothermometer. The calculations gave the following results: FABRIES (1979) 800°C, O'NEILL (1981) 700°C, SACK (1982) 900°C and ROEDER et al. (1979) 950°C. The widely scattered data from the same olivine-spinel pairs are mainly due to the large extrapolation from the experimental data obtained at very high temperatures and probably to some reequilibration effects. Indeed, the studies of LEHMANN (1983) and OZAWA (1984) have shown the importance of Fe-Mg diffusion between olivine and spinel and its influence on

thermometric calibrations. The presented estimates, however, give a reasonable order of magnitude of the metamorphic temperature suffered by the harzburgite of Mt. Mary.

The mineralogy, textures and estimated temperature ranges are consistent with a complete reequilibration of the ultramafic body under high-temperature amphibolite/granulite facies conditions. JENKINS (1983) proposed a simplified petrogenetic grid for hydrous ultramafites in the CNMASH system. Based on this grid, the observed amphibole-spinel and amphibole-spinel-chlorite assemblages are consistent with the inferred metamorphic conditions. Similar assemblages have also been reported by ROBINSON et al. (1982, Tab. 6, page 100).

The evolution of the Mt. Mary harzburgite may be traced, starting from a former spinel lherzolite stage (any traces of plagioclase or garnet are missing). A lherzolitic composition of the former protholith may be inferred from the high CaO content of the rock, which is now retained in the abundant amphibole.

The lack of tremolite-rich amphiboles and the Al-rich composition of the hornblende indicate that the peridotite underwent metamorphic reequilibration above 10 kbars in the chlorite - amphibole field, i.e., above the [Gt] invariant point of JENKINS (1981; 1983).

These relationships have been obtained from Fe-free CMASH and NCMASH assemblages. The mineral phases in the present study show considerable Fe contents ($X_{Mg}(\text{bulk rock}) = 0.82$), which affects the location of the univariant curves in the P-T space. CAIRONI & TROMMSDORFF (1988, Fig. 25, page 46) indicate qualitatively that the univariant curves for the tremolite and chlorite stability limits are displaced to lower temperatures.

The latest tectono-metamorphic event recorded by the ultramafite is the serpentinization of olivine along micro-shear zones under lower temperature and pressure conditions. This last event is most probably of Alpine age.

5. Conclusions

The presented structural, petrographic and geochemical data of the Mont Mary peridotite are consistent with a metamorphic reequilibration of a former spinel lherzolite under HT amphibolite/granulite facies conditions, at pressures possibly exceeding 10 kbars. A lherzolitic protholith is inferred by the high CaO content of the rock, now retained in the amphiboles, but probably derived from former clinopyroxenes.

The association of the ultramafic slice with continental lithologies, as well as the geochemical features, suggests a non-ophiolitic origin of the rocks. Consequently, they may be interpreted as a fragment of the subcontinental mantle, which has been emplaced and reequilibrated at lower crustal levels (granulite-facies conditions) and subsequently underwent exhumation and deformation during the Alpine orogenesis.

Another possible interpretation of the ultramafic lens of Mont Mary is a cumulate origin of the rock body, resulting from fractional crystallization of calc-alkaline or tholeiitic magmas. Such an interpretation has been suggested by one of the reviewers, basing on the low MgO, high Al_2O_3 and FeO_{tot} concentrations of the bulk rock sample and the relatively low XMg of the mineral phases of the hornblende-harzburgite.

We object to this interpretation because of the absence of clearly related magmatic rocks of basic and intermediate composition. Such rocks do, in general, accompany cumulate-type ultrabasic rocks (e.g. Klamath Mountains, California, SNOKE et al., 1981).

Acknowledgements

Prof. G.V. Dal Piaz thoroughly reviewed the manuscript, and the useful comments of Dr. Peter Ulmer improved it in the last version. We wish to thank them, and also Dr. Paola Tartarotti for the help given with the chemical analyses. Part of the analytical work was performed at the C.N.R. electron microprobe laboratory of the Earth Sciences Department, Università di Milano.

This study was carried out with the financial support of M.P.I. (40%, 60% grant to G. V. DAL PIAZ).

References

- ARGAND, E. (1934): La zone penninque. *Guide Géol. Suisse*, 3, 145-191.
- BECCALUVA, L., DAL PIAZ, G.V., MACCIOTTA, G. and ZEDA, O. (1978): The Austroalpine harzburgite body of the Artogna valley (Italian Western Alps) *Mem. Ist. Geol. Min. Univ. Padova*, 33, 173-181.
- BENCIOLINI, L. (1988): A new ultramafic body discovery in the Mt. Emilius Klippe (Austroalpine crust, Western Alps). In progress.
- CAIRONI, V. and TROMMSDORFF, V. (1988): *Lezioni di petrologia delle rocce metamorfiche*. Clup, Milano.
- CANEPA, M., CASTELLETTO, M., CESARE, B., DAL PIAZ, G.V., MARTIN, S., POLINO, R. and ZAGGIA, L. (1987): The Mont Mary Klippe: a composite nappe of the Austroalpine system. (Poster), V Réunion Esquisse Tectonique Suisse, Fribourg, 4-5 December, 1987.
- CANEPA, M., CASTELLETTO, M., CESARE, B., MARTIN, S., POLINO, R. and ZAGGIA, L. (in press): The Austroalpine Mont Mary composite nappe, Italian Western Alps. *Mem. Sc. Geol., Padova*.
- CESARE, B. (1987): Studio geologico e strutturale del lembo del Mont Mary (settore occidentale). Unpublished thesis, *Inst. Geol. Univ. Padova*, 202 p.
- COLBY, J.W. (1972): MAGIC IV: a computer program for quantitative electronmicroprobe analysis. Bell Telephone Laboratories Inc., Pennsylvania.
- DAL PIAZ, G.V., HUNZIKER, J.C. and MARTINOTTI, G. (1972): La zona Sesia-Lanzo e l'evoluzione tettonico-metamorfica delle Alpi nordoccidentali interne. *Mem. Soc. Geol. It.*, 11, 433-466.
- DIEHL, E.A., MASSON, R. and STUTZ, A.H. (1952): Contributo alla conoscenza del ricoprimento Dent Blanche. *Mem. Ist. Geol. Min. Univ. Padova*, 17, 5-52.
- ELTER, G. (1960): La zona pennidica dell'alta e media Valle d'Aosta e le unità limitrofe. *Mem. Ist. Geol. Min. Univ. Padova*, 22, 1-113.
- ERNST, W.G. (1978): Petrochemical study of lherzolitic rocks from the Western Alps. *J. Petrol.*, 19, 341-392.
- EVENSEN, N.M., HAMILTON, P.T. and O'NIONS, R.K. (1978): Rare-earth abundances in chondritic meteorites. *Geoch. Cosmoch. Acta*, 31, 1637-1665.
- FABRIES, J. (1979): Spinel-olivine geothermometry in peridotites from ultramafic complexes. *Contrib. Mineral. Petrol.*, 69, 329-336.
- HUNZIKER, J.C. (1974): Rb-Sr and K-Ar age determination and the Alpine tectonic history of the Western Alps. *Mem. Ist. Geol. Min. Univ. Padova*, 31, 54 pp.
- HUNZIKER, J.C. and ZINGG, A. (1980): Lower Paleozoic Amphibolite to Granulite facies metamorphism in the Ivrea Zone (Southern Alps, Northern Italy). *Schweiz. Min. Petr. Mitt.*, 60, 181-213.
- JENKINS, D.M. (1981): Experimental phase relations of hydrous peridotites modelled in the system H_2O -CaO-MgO- Al_2O_3 - SiO_2 . *Contrib. Mineral. Petrol.*, 77, 166-176.
- JENKINS, D.M. (1983): Stability and composition relations of calcic amphiboles in ultramafic rocks. *Contrib. Mineral. Petrol.*, 83, 375-384.
- LEAKE, B.E. (1978): Nomenclature of amphiboles. *Can. Min.*, 16, 501-520.
- LEHMANN, J. (1983): Diffusion between olivine and spinel: applications to geothermometry. *Earth Planet. Sci. Lett.*, 64, 123-138.
- MASSON, R. (1938): *Geologisch-Petrographische Untersuchungen im unteren Valpeltine*. *Schweiz. Mineral. Petrogr. Mitt.*, 18, 54-212.
- NICOT, E. (1977): Les roches mèse- et catazonales de la Valpeltine (nappe de la Dent Blanche, Alpes Italiennes). Thèse doctorat 3ème cycle, Univ. Paris VI, unpublished.
- O'NEILL, H.S.C. (1981): The transition between spinel lherzolite and garnet lherzolite and its use as a geobarometer. *Contrib. Mineral. Petrol.*, 77, 185-194.
- OTTONELLO, G., ERNST, W.G. and JORON, J.L. (1984): Rare earth and 3d transition element geochemistry of peridotitic rocks: I. Peridotites from the Western Alps. *J. Petrol.*, 25, 343-372.
- OZAWA, K. (1984): Olivine-spinel geospeedometry: analysis of diffusion controlled Mg-Fe²⁺ exchange. *Geoch. Cosmoch. Acta*, 48, 2597-2611.
- POGNANTE, U., RÖSLI, U. and TOSCANI, L. (1985): Petrology of ultramafic and mafic rocks from the Lanzo peridotite body (Western Alps). *Lithos*, 18, 201-214.
- ROBINSON, P., SPEAR, F.S., SCHUMACHER, J.C., LAIRD, J., KLEIN, C., EVANS, B.V. and DOOLAND, B.L. (1982): Phase relations of metamorphic amphiboles: natural occurrence and theory. Vol. 9b. *Petrology and experimental phase relations*. D.R. Veblen & H. Ribbe editors, *Miner. Soc. of America*, 1-227.

- ROEDER, P., CAMPBELL, I.H. and JAMIESON, H.E. (1979): A re-evaluation of the olivine-spinel geothermometer. *Contrib. Mineral. Petrol.*, 68, 325-334.
- SACK, R.O. (1982): Spinel as petrogenetic indicators: activity-composition relations at low pressures. *Contrib. Mineral. Petrol.*, 79, 169-186.
- SANDRONE, R. and COMPAGNONI, R. (1986): Relics of pargasite-bearing peridotite in the antigorite serpentinite of Balangero, near Lanzo (Western Alps). *Ophioliti*, 11, 35-47.
- SNOKE, A.W., QUICK, J.E. and BOWMAN, H.R. (1981): Bear Mountain igneous complex, Klamath Mountains, California: an ultrabasic to silicic calc-alkaline suite. *J. Petrol.*, 22, 501-552.
- STUTZ, A.H. and MASSON, R. (1938): Zur Tektonik der Dent Blanche Decke. *Schweiz. Mineral. Petrogr. Mitt.*, 18, 40-53.
- VOSHAGE, H., HUNZIKER, J.C., HOFFMANN, A.W. and ZINGG, A. (1987): A Nd and Sr isotopic study of the Ivrea zone, Southern Alps, N-Italy. *Contrib. Mineral. Petrol.*, 97, 31-42.
- ZAGGIA, L. (1987): Studio geologico e strutturale del lembo del Mont Mary (settore centrale) - Unpublished thesis, Inst. Geol. Univ. Padova, 181 pp.

Manuscript received November 11, 1988; revised manuscript accepted February 2, 1989