

**Zeitschrift:** Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie  
**Band:** 70 (1990)  
**Heft:** 3  
  
**Artikel:** Fission track and nannofossil ages from a Paleocene bentonite in the Schlieren Flysch (Central Alps, Switzerland)  
**Autor:** Winkler, Wilfried / Hurford, Anthony J. / Salis Perch-Nielsen, Katharina von  
**DOI:** <https://doi.org/10.5169/seals-53629>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Fission track and nannofossil ages from a Paleocene bentonite in the Schlieren Flysch (Central Alps, Switzerland)

by Wilfried Winkler<sup>1</sup>, Anthony J. Hurford<sup>2</sup>, Katharina von Salis Perch-Nielsen<sup>1</sup> and Gilles S. Odin<sup>3</sup>

### Abstract

A bentonite layer contained in the Schlieren Flysch series has been dated by calcareous nannofossils and by radiometric fission track analysis of volcanic zircon grains. The nannofossil assemblage belongs to NP5 zone and the mean fission track age is  $57.8 \pm 2.7$  Ma. This new age constrains the poorly documented Paleocene part of the geological time scale.

**Keywords:** Bentonite, zircon, fission track, calcareous nannofossils, Schlieren Flysch, Central Alps, Switzerland.

### Introduction

The presence of bentonite layers in the Voirons, Gurnigel, Schlieren and Wägital Flysch was reported by WINKLER (1983) and WINKLER et al. (1985 a, b). The volcanic origin of these thin, yellow and white altered layers was deduced from the clay mineral and heavy mineral composition, and by the presence of recrystallized pyroclastic debris in some layers. Chemical analysis and theoretical correction for diagenetic modifications point generally to partly andesitic and partly alkali basaltic (trachy-basaltic) sources for these bentonites.

The record of synorogenic volcanism is generally very poor in the Alps. The occurrence of the altered ash layers thus may be due to distant volcanic activity not directly related to the Alpine orogeny. However, palaeogeographic, sedimentologic and textural considerations make it probable that they were derived from subduction related arc or back-arc volcanic centers situated to the south of the Alpine orogen (WINKLER, 1983; WINKLER et al., 1985b).

The host rock of the bentonites is a 1500–1800 m thick turbiditic and hemipelagic Late Cretaceous to Eocene flysch series forming a more or

less continuous belt from Lake Geneva to Central Switzerland. From sedimentologic, petrographic and structural evidence it appears that the 300 km long flysch sequence was deposited in a remnant South Penninic, periodically convergent deep-sea trench basin with a source area composed of both basement and sedimentary cover rocks (WINKLER, 1983, 1984b). The lack of carbonate material in the hemipelagic layers points to a depositional environment below the local CCD (e.g. WINKLER, 1984a).

We have observed 14 nearly uncontaminated bentonite layers in Late Maastrichtian, Paleocene and Early Eocene flysch series and, particularly in the Tertiary, they are closely related to the Tonstein-Schichten facies representing diachronous abyssal plain/oceanic slope deposits, comparatively rich in lime-free hemipelagic shales (WINKLER, 1983, 1984b). Their occurrence is also closely linked to the presence of iron- and manganese-bearing layers (WINKLER et al., 1985b).

The bentonite layers consist of a few millimeter to 2 centimeter thin yellow and white clay layers with high contents of montmorillonite in free phase (up to 100%) or in the mixed-layer phase montmorillonite/illite (up to 80%). As an important test of

<sup>1</sup> Department of Earth Sciences, Swiss Federal Institute of Technology, ETH-Zentrum, CH-8092 Zürich, Switzerland.

<sup>2</sup> Department of Geological Sciences, University and Birkbeck Colleges, Gower Street, London WC1E 6BT, United Kingdom.

<sup>3</sup> Laboratoire de Géochronologie et Sédimentologie Océanique, Université Pierre et Marie Curie, 4, place Jussieu – Tour 26, 75252 Paris, France.

purity, in individual layers the total heavy mineral content was recorded. The assemblages consist nearly exclusively of euhedral apatite, zircon and a few pseudo-hexagonal biotites and thus represent uncontaminated volcanic ash layers. Additional tourmaline, rutile, anatase, brookite and garnet were considered as terrigenous impurities (WINKLER et al., 1985b).

One of these layers (sample WW 1948) of early Late Paleocene age revealed a very rich heavy mineral content with abundant and well preserved euhedral zircon grains. Together with the generally good calcareous nannofossil control in the flysch series (e.g. WINKLER, 1983) it was considered valuable to combine fission track dating of zircon with current nannofossil zonation. In spite of the occurrence of the ash layer WW 1948 in a very lime-

poor facies, by repeated sampling we were successful in finding good nannofossil assemblages allowing a first hand correlation of biostratigraphy with the fission track dates.

#### Specification of bentonite WW 1948

The analysed ash layer occurs in the Änggenlouenen section in the Entlebuch area ca. 30 km SW of Lucerne (section locality, Swiss topographic map 1: 25.000 "Schüpfheim", coord. 647.760 / 194.020). The samples were taken from the Lower Tonstein-Schichten Fm. at the foot of the Fürstein Mt. chain (see Figs 1 and 2).

The Lower Tonstein-Schichten Fm. at this location belongs to the outer facies belt of the Schlieren flysch basin, characterized by thick abyssal




Fig. 1 Simplified geological map of Central Switzerland with location of the Fürstein mountain.




Fig. 2 View of the northern slope of the Fürstein with Paleocene and Lower Eocene formations. The box indicates the position of section A in Fig. 3 with the bentonite in the lowermost part (detailed section B).

plain/oceanic slope deposits, reaching a younger age (earliest Late Paleocene) in its top than in the inner facies belt (NP4 or latest Early Paleocene, WINKLER, 1983, 1984b). They consist of a generally thin-bedded repetition of fine-grained turbiditic sandstones and shales intercalated with comparatively thick and well developed greyish-green hemipelagic claystones (see Fig. 3). In contrast to other formations of the Schlieren Flysch series, the turbiditic shales in the Tonstein-Schichten facies are always poor in carbonate material. In addition to this, the Lower Tonstein-Schichten Fm. is characterized by the occurrence of greenish, well sorted, quartzitic turbidite beds, referred to as "Öl-quarzite" by the pioneers of Swiss geology.

The clay fraction of the bentonite is composed of 67% montmorillonite and 29% illite in the mixed-layer phase; kaolinite amounts to 4% (semi-quantitative estimates). From statistically counted heavy mineral grains (enriched by bromoform separation technique) 68% are apatite, 30% zircon, 2% titanite, 1% chromite and 3% contaminating non-volcanic minerals (2% tourmaline, 1% brookite). The typology of zircon (PUPIN, 1980) was evaluated on 150 individual grains and the homogeneous distribution in the diagram indicates a primary calc-alkaline magma source. The observed

homogeneous typology of the zircon grains proves that the layer is the product of a single and not reworked volcanic event. The assumed diagenesis corrected chemical composition of the bentonite points to the presence of a basaltic or phonolitic volcanic ash, depending on which correction factors are applied (see WINKLER et al., 1985b).

There is good control on the maximum temperature reached during burial of the bentonites. From illite crystallinity measurements, clay mineral assemblage and feldspar preservation it is assumed that the series was affected by low to medium grade diagenetic overprint of approximately 70–100 °C (WINKLER, 1983). Therefore apatite fission track dating is not recommended.

#### Calcareous nannofossil biostratigraphy

The nannofossil samples were mostly taken at the turbiditic sand-shale transition, because this is in the present formation the most favourable place to obtain carbonate-bearing and sufficiently soft samples. Six samples were studied for calcareous nannofossils in simple smear-slides and examined with an optical microscope. Five of them contained quite rich assemblages (Tab. 1). The preservation is




Fig. 3 Composite (A) and detailed (B) section of the Änggenlouenen outcrop in the Lower Tonstein-Schichten Formation.

Tab. 1 Calcareous nannofossil biostratigraphy below and above the bentonite layer in the Schlieren Flysch. Abbreviations: vR, + = very rare, present, R = rare, Rf = rare to few, C = common. Preservation: vP = very poor, P = poor, Pm = poor to moderate, M = moderate. For complete citations of authors of species see PERCH-NIELSEN (1985 a, b).

Sample	WW 1954	WW 3037	WW 1957	WW3039	WW 3271	WW 794
Nannofossil abundance & preservation	C, Pm	vR, vP	R, Pm	R, Pm	R, M	C, Pm
<b>Palaeocene nannofossils</b>						
<i>Chiasmolithus</i> sp.	+					
<i>C. edentulus</i>	+		+			
<i>Coccolithus pelagicus</i> s.a.	+	+	Rf	+	R	R
<i>Fasciculithus billii</i>	?	+	?	?	?	
<i>F. janii</i>	+		+	+		
<i>F. tympaniformis</i>	+	+	+	+	+	Rf
<i>F. ulii</i>	+		+			
<i>F. sp. cf. F. magnus</i>	+			+		
<i>Markalius inversus</i>	+		+			
<i>Neochiastozygus</i> sp.	+					
<i>Neocrepidolithus</i> sp.	+					
<i>Placozygus sigmoides</i>	+	+	+	+	+	+
<i>Prinsius martinii</i>	+		+	+	R	
<i>Sphenolithus primus</i>	C	+	+	+	+	Rf
<i>Thoracosphaera</i> sp.	+			+	R	+
<i>Toweius craticulus</i>	+		+	+	+	+
<i>T. eminens</i>	+		+	+	+	R
<i>Chiasmolithus danicus</i>			+			
<i>C. inconspicuus</i>			+	+		
<i>Crucioplacolithus tenuis</i> s.a.			+			
<i>Ericsonia subpertusa</i>				+		
<i>Ericsonia</i> sp. cf. <i>E. robusta</i>			+			
<i>Fasciculithus pileatus</i>			+	+		
<i>Neochiastozygus perfectus</i>			+	+	+	
<i>Prinsius bisulcus</i>			+			+
<i>Chiasmolithus bidens</i>				+	+	+
<i>C. frequens</i>				+		
<i>Braarudosphaera bigelowii</i>				+		
<i>Ellipsolithus distichus</i>				+		
<i>Neochiastozygus digitosus</i>				+		
<i>Lanternites duocavus</i>				+		
<i>Ellipsolithus macellus</i>					+	
<i>Pontosphaera</i> sp.					+	
<i>Heliolithus cantabriae</i>					+	+
<i>H. kleinpellii</i>					+	Rf
<i>Sphenolithus anarrhopus</i>						+
Nannofossil - Zone	NP 5				NP 6	
<b>Mesozoic nannofossils (reworked)</b>						
<i>Arkhangelskiella cymbiformis</i>	+		+	+	+	
<i>Ceratolithus aculeus</i>					+	
<i>Chiastozygus</i> sp.					+	
<i>Chiastozygus fessus</i>				+		
<i>Eiffelithus eximius</i>			+			+
<i>E. turriseiffelii</i>					+	
<i>Ellipsagelosphaera communis</i>				+	+	
<i>Erolithus floralis</i>						+
<i>Lucianorhabdus</i> sp.	+		+			
<i>Micula decussata</i>	+	+	+	+	R	+
<i>Nannoconus steinmannii</i>					+	
<i>Prediscosphaera cretacea</i>			+			
<i>Reinhardtites anthophorus</i>				+		
<i>Stephanolithon bigotii</i>					+	
<i>Stradneria crenulata</i>			+			
<i>Watznaueria barnesae</i>	+	+	+	+	R	+



very poor in one sample and poor to moderate in the others. The poorly preserved assemblages are affected heavily by dissolution. This has affected the diversity of the assemblages, in that fragile species have been removed or broken and are no longer determinable. The biostratigraphic marker species of the Late Paleocene are, however, relatively solution resistant and thus the zonal assignments are judged to be realistic.

The two samples below and the two samples above the bentonite layer (see Fig. 3, Tab. 1) feature assemblages typical of NP5, the *Fasciculithus tympaniformis* Zone of MARTINI (1971), including *F. tympaniformis* but no *Heliolithus kleinpellii*. The latter appears very rare in sample WW 3271. It reaches rare to few in the uppermost sample studied and assigns these two samples to NP6, the *Heliolithus kleinpellii* Zone, since the marker of NP7, *Discoaster mohleri*, was not found. No calcareous nannofossils indicative of younger zones were observed.

VAROL (1989) subdivided both NP5 and NP6, a subdivision that was not feasible in our material. He noticed an overlap of *H. kleinpellii* with *Fasciculithus jani* and *F. pileatus*, an interval that should fall between samples WW 3039 and WW 3271. Its absence might indicate slow sedimentation or a hiatus between those two samples.

Reworked Mesozoic coccoliths were found in all samples. Most species have a Late Cretaceous range but sample WW 3271 also contains species with a Late Jurassic range (*Stephanolithion bigotii*) and some with an early Cretaceous distribution (*Nannoconus steinmannii*) – an expression of deeper erosion during the regression around the NP 5/6 boundary (HAQ et al., 1988)?

### Fission track ages

The clayey bentonite samples (totally approx. 1.5 kg) were disaggregated by repeated soaking in cold Désogéne (CIBA-GEIGY, Basel) and short treatments in the ultra-sound bath. Heavy minerals (including zircons) were separated from the sandy residue by conventional heavy liquid (bromofom) techniques. Fission track ages were measured for two independent zircon samples (B-20 and WW 1948) using the external detector method approach (GLEADOW, 1981).  $^{238}\text{U}$  spontaneous fission tracks were etched in the polished zircon crystals using a eutectic of KOH–NaOH at 220 °C for 15 hrs (GLEADOW et al., 1976).  $^{235}\text{U}$  induced fission tracks were recorded in an external detector of low-uranium mica, held against the zircons during irradiation with thermal neutrons. These micas were subsequently etched using 40% HF at 20 °C for 45 mins.

Sample B-20 (see preliminary presentation by HURFORD et al., 1987) was irradiated in the J1 thermal facility of the HERALD reactor, Aldermaston, UK, and sample WW 1948 in the thermal column of the PLUTO reactor, Harwell, UK. Track counting utilised a ZEISS Axioplan microscope (100× oil objective, total magnification 1250×) equipped with a Stagemover computerised stage system to locate the zircon crystals and their mirror image mica impressions. Neutron fluences were monitored by including wafers of uranium dosimeter glass CN1 (HURFORD and GREEN, 1983) at either end of the sample stack and counting the induced tracks recorded in a mica detector held against the glass.

Sample ages (Tab. 2) were calculated using the zeta calibration approach (FLEISCHER and HART, 1972) using a value of  $113.0 \pm 2.6$  for  $\zeta_{\text{CN1}}$  determined by repeated analysis of age standards (HURFORD and GREEN, 1983). As a control of calibration, aliquots of zircon age standards from the Fish Canyon tuff (San Juan, Colorado, see HURFORD and HAMMERSCHMIDT, 1985) and the Buluk Member tuff (northern Kenya, see HURFORD and WATKINS, 1987) were included in each irradiation. Analytical results are given in Tab. 2. Analyses were subjected to a  $\chi^2$ -test (GALBRAITH, 1981) to determine whether any extra-Poissonian error might be present. Each data set passed the test at > 5% level, indicating the Poissonian error (the conventional error of GREEN, 1981) derived from counting statistics to be a reasonable estimator of experimental error.

The results for the two samples are identical within their errors with a mean of  $57.8 \pm 2.7$  (1  $\sigma$ ) Ma (or a weighted mean of  $57.8 \pm 1.9$  [1  $\sigma$ ] Ma).

### Discussion and conclusions

Our data indicate that the bentonite, embedded in sediments biostratigraphically correlated with NP5 zone (early Late Paleocene) contains volcanic zircons which yield a mean fission track age of  $57.8 \pm 2.7$  Ma. Biostratigraphic dating had to be carried out on turbiditic shales. The first occurrences of nannofossil marker species were considered for zonal assignments. Both the reworked nature and possible postdepositional dissolution could represent a probable source of error. However, the zonal markers for NP5 (*Fasciculithus tympaniformis*) and NP6 (*Heliolithus kleinpellii*) are rather solution resistant forms and they occur in our section in an obviously undisturbed succession. In addition, this biostratigraphic correlation fits well with other nannofossil dates in adjacent areas (WINKLER, 1983).

Tab. 2 Zircon fission track data measured on the bentonite samples B-20 and WW 1948 and standards.

Samples and Standards	Mineral and No. Crystals	Spontaneous $\rho_s$ ( $N_s$ )	Induced $\rho_i$ ( $N_i$ )	$P_X^2$	Dosimeter $\rho_d$ ( $N_d$ )	Age Ma $\pm 1 \sigma$
<b>Schlieren Flysch</b>						
B-20	zircon 20	6.728 (2122)	4.305 (1358)	70%	0.66 (4700)	58.0 $\pm$ 2.5
Wi 1948	zircon 10	8.181 (1167)	4.893 (698)	20%	0.611 (3880)	57.5 $\pm$ 2.9
<b>Age Standards</b>						
Fish Canyon	zircon 20	5.535 (2778)	8.395 (4213)	55%	0.749 (4700)	27.8 $\pm$ 1.1
Fish Canyon	zircon 14	6.31 (2910)	8.127 (3748)	45%	0.611 (3880)	26.8 $\pm$ 0.8
Buluk Member	zircon 20	1.149 (1433)	2.869 (3579)	65%	0.742 (4700)	16.8 $\pm$ 0.7
<b>Notes:</b> (i). track densities ( $\rho$ ) are as measured and are ( $\times 10^6 \text{ tr cm}^{-2}$ ); numbers of tracks counted ( $N$ ) shown in brackets; (ii). analyses by external detector method using 0.5 for the $4\pi/2\pi$ geometry correction factor; (iii). ages calculated using dosimeter glass CN-1 with $\zeta_{\text{CN1}} = 113 \pm 3$ ; other constants as defined in HURFORD and GREEN (1983); (iv). $P_X^2$ is probability of obtaining $\chi^2$ value for $v$ degrees of freedom, where $v = \text{no. crystals} - 1$ ; (v). independent ages of standards: Fish Canyon Tuff $27.8 \pm 0.7$ Ma; Buluk Member Tuff $16.2 \pm 0.2$ Ma (see HURFORD and GREEN, 1983; HURFORD and WATKINS, 1987).						

Preliminary radiometric  $^{238}\text{U}/^{206}\text{Pb}$  dating of two zircon samples from the same bentonite layer was provided by FISCHER (1988) revealing ages of  $52.57 \pm 0.22$  and  $54.78 \pm 0.50$  ( $2 \sigma$ ) Ma. These quite similar and young dates certainly confirm the syn-sedimentary volcanic derivation of the zircon grains. However, in comparison with current chronostratigraphic calibrations as CURRY and ODIN (1982) these radiometric ages correlate with NP zones 8–10 at about the Paleocene-Eocene boundary. From palaeontologic evidence the fission track ages are preferred to the U–Pb ages since the transition from Late Paleocene to Early Eocene corresponding with the U–Pb ages occurs in the same section higher up in the series at the boundary between NP 9 and NP 10 which coincides with the

transition from the Guber Sandstone to the Upper Tonstein-Schichten Formation in Fig. 2.

The present fission track age of  $57.8 \pm 2.7$  Ma corresponds better with most existing chronostratigraphic and biostratigraphic correlations. It coincides with NP5 in CURRY and ODIN (1982), approximately with the transition from NP5 to NP6 in HARLAND et al. (1982; but not with HARLAND et al., 1990) and lowermost NP6 in HAQ et al. (1988). Because of the poor outcrop conditions at the base of the Fürstein mountain chain the position of the bentonite inside the NP5 zone cannot be more precisely assessed at the present time.

The mean fission track age of  $57.8 \pm 2.7$  Ma is a new constraint for calibrating the Palaeogene time scale at a place where very few results are available.



### Acknowledgements

Fission track dating in London is supported by NERC Grant GR3/7068 and by a BP research award. In early stages of the present work one of us (WW) was supported by the Swiss National Science Foundation, projects 2.486-0.82 and 2.083-0.83. Thanks go also to Diane Seward, Daniel Bernoulli and Walter Wildi for the critical review of the earlier version of the manuscript.

### References

- CURRY, D. and ODIN, G.S. (1982): Dating of the Palaeogene. In: G.S. ODIN (ed.): Numerical dating in stratigraphy. John Wiley and Sons, Ltd., 607-630.
- FISCHER, H. (1988): Isotopenchemische Untersuchungen und Datierungen an Mineralien und Fossilien aus Sedimentgesteinen: 1. Glaukonite aus dem Jura, der Molasse und dem Helvetikum (K-Ar, Rb-Sr), 2.  $^{87}\text{Sr}/^{86}\text{Sr}$ -Isotopenstratigraphie an marinen und limnischen Mikro- und Makrofossilien, 3. Primärminerale aus tertiären Bentoniten und Tuffen (U-Pb, K-Ar). Unpubl. PhD-thesis, ETH-Zürich.
- FLEISCHER, R.L. and HART, H.R. (1972): Fission track dating: technique and problems. In: BISHOP, W.W., MILLER, J.A. and COLE, S. (eds): Calibration of Hominid evolution. Scottish Academic Press, Edinburgh, 135-170.
- GALBRAITH, R.F. (1981): On statistical models for fission track counts. *Mathematical Geology* 13, 471-488.
- GLEADOW, A.J.W. (1981): Fission track dating methods: what are the real alternatives? *Nuclear Tracks* 5, 3-14.
- GLEADOW, A.J.W., HURFORD, A.J. and QUARF, R.D. (1976): Fission track dating of zircon: improved etching conditions. *Earth and Planetary Science Letters* 33, 273-276.
- GREEN, P.F. (1981): A new look at statistics in fission track dating. *Nuclear Tracks*, 5, 77-86.
- HAO, B.U., HARDENBOL, J. and VAIL, P.R. (1988): Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level change. *Soc. Econom. Paleont. Mineral. Spec. Publ.* 42, 71-108.
- HARLAND, W.B., COX, A.V., LLEWELLYN, P.G., PICKTON, C.A.G., SMITH, A.G. and WALTERS, R. (1982): A geologic time scale. Cambridge University Press, 131 p.
- HARLAND, W.B., ARMSTRONG, R.L., COX, A.V., CRAIG, L.E., SMITH, A.G. and SMITH, D.G. (1990): A geologic time scale. Cambridge University Press, 263 p.
- HURFORD, A.J. and GREEN, P.F. (1983): The zeta age calibration of fission track dating. *Isotope Geoscience* 1, 285-317.
- HURFORD, A.J. and HAMMERSCHMIDT, K. (1985):  $^{40}\text{Ar}/^{39}\text{Ar}$  and K/Ar dating of the Fish Canyon and Bishop Tuff: calibration ages for fission track dating standards. *Chemical Geology (Isotope Geoscience Section)* 58, 23-32.
- HURFORD, A.J. and WATKINS, R.T. (1987): Fission track age of the Buluk Member, Bakate Formation, northern Kenya: a suitable fission track standard. *Chemical Geology (Isotope Geoscience Section)* 66, 209-216.
- HURFORD, A.J., ODIN, G.S., BARBIN, V., WINKLER, W. and ZELVELDER, M. (1987): Fission track dating of Palaeogene pyroclastic zircons and apatites: The Priabonian of Priabona (stratotype area), the earliest Thanetian in the northern Alps. *EUG IV - Meet. Strasbourg, Abstracts*, 251.
- MARTINI, E. (1971): Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: FARINACCI, A. (ed.): Proc. II Plankt. Conf., Roma 1970, 2, Tecnoscienza, Roma, 739-785.
- PERCH-NIELSEN, K. (1985a,b): Mesozoic calcareous nanofossils. Cenozoic calcareous nanofossils. In: BOLLI, H.M., SAUNDERS, J.B. and PERCH-NIELSEN, K. (eds): *Plankton Stratigraphy*, Cambridge University Press, Cambridge, 329-426 and 427-554.
- PUPIN, J.P. (1980): Zircon and Granite Petrology. *Contr. Mineral. Petrol.* 73, 207-220.
- VAROL, O. (1989): Palaeocene calcareous nanofossil biostratigraphy. In: CRUX, J.A. and VAN HECK, S.E. (eds): *Nannofossils and their applications*, Ellis Horwood Ltd, Chichester, 267-310.
- WINKLER, W. (1983): Stratigraphie, Sedimentologie und Sedimentpetrographie des Schlieren-Flysches (Zentralschweiz). *Beitr. geol. Karte Schweiz [N.F.]* 158.
- WINKLER, W. (1984a): Rhabdammina or "Flysch-type agglutinated" fauna - what relation to turbidites?. In: OERTLI, H.J. (ed.): *Benthos '83*, 2nd int. Symp. on Benthic Foraminifera (Pau, April 1983. Elf Aquitaine, Pau and Bordeaux, 611-617.
- WINKLER, W. (1984b): Palaeocurrents, and petrography of the Gurnigel-Schlieren flysch: a basin analysis. *Sediment. Geol.* 40, 169-189.
- WINKLER, W., WILDI, W., VAN STUIJVENBERG, J. and CARON, C. (1985a): Wägital-Flysch et autres flysch peniques en Suisse Centrale: Stratigraphie, sédimentologie et comparaisons. *Eclogae geol. Helv.* 78, 1-22.
- WINKLER, W., GALETTI, G. and MAGGETTI, M. (1985b): Bentonite im Gurnigel-, Schlieren- und Wägital-Flysch: Mineralogie, Chemismus, Herkunft. *Eclogae geol. Helv.* 78, 545-564.

Manuscript received August 30, 1990; revised manuscript accepted September 24, 1990.