

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 70 (1990)

Heft: 3

Artikel: Amphibolites from Lake Emosson/Aiguilles Rouges, Switzerland :
tholeiitic basalts of a Paleozoic continental rift zone

Autor: Raumer, J.F. von / Galetti, G. / Pfeifer, H.R.

DOI: <https://doi.org/10.5169/seals-53632>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 29.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Amphibolites from Lake Emosson/Aiguilles Rouges, Switzerland: Tholeiitic basalts of a Paleozoic continental rift zone

by J.F. von Raumer¹, G. Galetti¹, H.R. Pfeifer² and R. Oberhänsli³

Abstract

In the northern part of the external Aiguilles Rouges massif, two 1 to 5 m thick zones of amphibolite interlayered in micaschists can be followed over a distance of several kilometers. The whole series is strongly folded and shows polyphase metamorphic mineral assemblages grading from eclogite to upper amphibolite facies. The mineralogy is quite variable, but two major types of amphibolites can be distinguished: garnet-plagioclase amphibolites and plagioclase-amphibolites with rare garnet-pseudomorphs. The amphibole compositions vary from magnesio-hornblende to rare actinolitic amphibole and garnets are rich in almandine and grossular. The whole-rock composition of both types of amphibolites is tholeiitic, but vary considerably in the Mg-Fe-ratio (Mg-number) and some trace elements: the plagioclase-amphibolites with high Mg-numbers, rich in Cr and Ni, but poor in Ti, Zr and Y correspond to little evolved and the garnet-amphibolites to more fractionated tholeiitic magmas similar to present-day mid ocean ridge basalts (MORB). Chondrite-normalized rare earth element patterns are flat or slightly enriched in light rare earths ranging between 10 and 40 times chondrite, but do not differ significantly from one type to the other, corresponding to transitional type MOR-basalts. However, K-, Rb- and Ba-contents approach values typical for plume-type, rather enriched MOR-basalts. Such compositions are typical for continental rift zones with strong extension, leading at the end to the formation of oceanic crust, like the Jurassic flood basalt provinces of southern Africa and Brasil or the Tertiary opening of the Red Sea. The large compositional variability and the small volume suggest that the protoliths of these amphibolites were subvolcanic sills or dikes.

Keywords: Lake Emosson, Aiguilles Rouges massif, Switzerland, amphibolite, dike, geochemistry, metamorphism.

Introduction

The Emosson area lies in the pre-Carboniferous crystalline basement of the Helvetic realm of the Western Alps in the Swiss part of the Aiguilles Rouges massif. This Variscan basement area is part of the northernmost lying, external massifs and has undergone only minor thrusting during the Alpine orogeny. The accompanying metamorphism was of very low grade (max. 2 kb, 270 °C, VON RAUMER, 1984a), thus preserving relics of older events. The external massifs represent, regardless of their present outcrop area, small pieces of pre-Permian polymetamorphic crust (VON RAUMER, 1976, 1981, 1984a). Among the metamorphic rock suites that outcrop near the lake of Emosson, amphibolites

are, thanks to their possibly preserved geochemical signatures, particularly suitable to unravel the paleotectonic environment of this complex rock unit.

Geological setting and petrography

The first geologic outline of the Emosson area was presented by COLLET et al. (1952). Actual field work reveals a strongly folded, complex lithology (Fig. 1). A metasedimentary zone up to 1.5 km large is intruded by granitoid rocks limited in the east by a major Carboniferous shear zone. The metasediments derived from detrital sediments of Late Precambrian to Early Paleozoic age (VON RAUMER, 1988) build up a reduced series of inter-

¹ Institut de Minéralogie et de Pétrographie, Université, Pérolles, CH-1700 Fribourg, Switzerland.

² Centre d'Analyse Minérale, Université de Lausanne, BFSH2, CH-1015 Lausanne, Switzerland.

³ Institut für Geowissenschaften, Universität Mainz, Saarstrasse 21, D-6500 Mainz, Federal Republic of Germany.

layers of metagreywackes, quartzites and metapelites, where just one important micaschist layer can be recognized. These rock series suffered deformation and metamorphism of Caledonian- or Variscan age, summarized as follows (cf. VON RAUMER, 1984b, 1987): The earliest recognizable metamorphic event of the Aiguilles Rouges massif is preserved as meta-eclogites, which occur 15 km to the southwest of the study area. Their minimum P-T conditions have been estimated at 11 kb, 780 °C by LIÉGOIS and DUCHESNE (1980). According to PAQUETTE et al. (1989), this event is Caledonian in age

(425–395 Ma). Within the study area eclogite relics are only rarely found (see below), but the subsequent history can be reconstructed from the metapelitic rocks, which are characterized by a Barrovian type of metamorphism. Kyanite-garnet assemblages, estimated to have formed at 8–10 kb, 550 °C, are associated with large scale folds (F_2 , F_3) of kilometer scale and are thought to be of Silurian age (VON RAUMER, 1984 a, b). Sillimanite overgrowing kyanite, together with local anatexis, indicate a later increase in temperature and decrease in pressure (4–5 kb, 600–700 °C). This event is proba-




Fig. 1 Geological map of the Emosson area with the main lithologic units and the isoclinal fold pattern of amphibolites (black). In the southeast a series of Permian faults dissects all rock units.

bly of Devonian or Early Carboniferous age and is associated with open folds (F_4) in a NE-SW direction (JOYE, 1989). The late stage of the evolution is documented by Late Carboniferous shear zones (Fig. 1) accompanied by fluid migration and retrograde transformations. The presence of small quantities of pumpellyite and prehnite are attributed to the weak Tertiary metamorphic overprint (VON RAUMER, 1974).

The amphibolites are relatively scarce in comparison to the granitic and pelitic country rocks; they can be mapped over several kilometers and display a complex fold pattern (Fig. 1). They are mostly limited to the micaschist layer mentioned above and only locally found in the adjacent metagreywackes. At one locality a discordant contact with metaquartzites has been observed. Within a given layer, the amphibolites appear as lens shaped boudin-like bodies of 5 to 20 m in length. Such huge lenses with internal fold structures are interpreted as the hinge zones of older fold structures.

The amphibolites can be divided up into five types (Tab. 1) well distinguishable in the field which may occur together in one same lens; a garnet-plagioclase type (group 1) and a plagioclase type (group 2), both groups representing the volumetrically most significant rock types (Fig. 2) forming parallel layers. Of less importance are the clinzoisite bearing type (group 3), an amphibole-garnet schist (group 4) and a biotite-actinolite-garnet-plagioclase schist (group 5). Each of these is described below. Most types show a more or less lepidoblastic texture and are finegrained (0.5–1 mm). The schistosity, which is sometimes of mylonitic origin, is underlain by fine, plagioclase bearing layers, which may contain small garnets (type 1B).

The *garnet-plagioclase-amphibolites of group 1* contain garnet, amphibole and plagioclase as main phases. Three subtypes can be distinguished, based on grainsize (c.f. Tab. 1). The coarse grained type amphibolite (type 1A, e.g. sample 1604) always occurs at the limit between the amphibolite bodies




Fig. 2 Photomicrographs of the two major types of amphibolite of the Lake Emosson area.

A) Garnet (G) in plagioclase-garnet amphibolite with amphibole (grey) and plagioclase (white, rock type 1A, sample no. 1544).

B) Garnet (white field in the center) replaced by plagioclase (white) and amphibole prisms (grey) in a matrix of amphibole and plagioclase (rock type 2B, sample no. 1538). Diameter of garnet 2 mm.

Tab. 1 Typical modal compositions of the five types of amphibolites distinguished (c.f. chapt. petrography) determined by point counting.

Mineral	Rock type and sample number										
	1A 1604	1B 1507	1C 1513	1C 1527	2A 1515	2B 1523	2B 1538	2C 1554	3 1548	4 1536	5 1542
Amphibole	57	71	68	55	61	74	70	79	71	41	8
Plagioclase	32	28	27	30	32	22	30	21	7	35	15
Garnet	6	<1	1	0	0	0	0	0	0	22	18
Quartz	3	<1	3	15	0	1	<1	<1	15	2	14
Sphene	2	0	1	0	7	3	0	0	0	0	0
Ilmenite	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Biotite	0	0	0	<1	0	0	0	0	0	0	45
Clinozoisite	0	0	0	0	0	0	0	0	6	0	0

and the surrounding metapelite. The nearly euhedral garnets of about 5–8 mm diameter are surrounded by amphibole and plagioclase (Fig. 2A). In the blastomylonitic type amphibolites (type 1B, sample 1507) small garnets appear in thin plagioclase rich layers. In the main type garnet-plagioclase amphibolites (type 1C) garnets are lenticular in shape, and are very often partially replaced by plagioclase.

The majority of the *plagioclase-amphibolites of group 2* contain essentially plagioclase and amphibole in a granular texture (type 2B, sample 1523; Fig. 2B). The rare presence of textural relicts of garnet, consisting of granular plagioclase including some small prisms of amphibole, ilmenite, sphene and clinozoisite, show that a few of the plagioclase amphibolites samples represent the final stage of a transformation of former garnet amphibolites (type 2A, sample 1515). A few samples are characterized by large amphiboles that sometimes overgrow other phases (type 2C, sample 1554).

The *clinozoisite-bearing amphibolites of group 3* are a special feature found locally in the outer rim of the amphibolite bodies. White needles of cm-size consisting of clinozoisite, plagioclase, some quartz and calcite are interpreted as pseudomorphs after zoisite.

The *amphibole-garnet-schists of group 4* (sample 1544) occur in the center of some larger boudins. The large garnets contain inclusions of green or blue-green amphibole, sphene, and ilmenite and are surrounded by large prisms of actinolite and some plagioclase. In analogy to similar rocks from the nearby Lake Cornu area (LIÉGOIS and DUCHESNE, 1981) we interpret these rocks as relic eclogites. As they are strongly altered (plagioclase replaced by white mica, amphibole by calcite and chlorite) they have not been used for a more detailed investigation.

Finally the *biotite-plagioclase-actinolite-schists of group 5* appear as irregular lumps in a feldspar-rich zone in the border zone of larger amphibolite

boudins. Euhedral, zoned garnets or garnet relicts occur in a matrix of large biotites with few prisms of actinolite. Plagioclase generally shows an equigranular growth. It obviously represents a reaction rim between the amphibolites and their country rock. Because of intense replacement by sericitic white mica of all minerals no further investigation was undertaken.

Mineral chemistry

The composition of the main mineral phases has been analyzed on an ARL-SEMQ microprobe of the university of Bern, using natural and synthetic minerals as standards. The operating conditions were 15 kV accelerating voltage and 20 μ A sample current. The corrections were made with a Bence-Albee type of program.

AMPHIBOLES

As the microprobe only permits the determination of total iron, the chemical data were normalized after LAIRD (1978) and LAIRD and ALBEE (1981a) estimating Fe^{3+}/Fe^{2+} from charge balance and assigning endmembers from the formula proportions (Tab. 2A, Fig. 3a). In the nomenclature of LEAKE (1978) all amphiboles correspond to calcic amphiboles, most of them to magnesiohornblende, and only few to actinolites. Their composition is mainly characterized by Tschermak substitution (THOMPSON et al., 1981; Fig. 3a), whereas the edenite- and glaucophane-substitutions are less important. The few observed zoned amphiboles reveal an evolution from early actinolite composition to late magnesiohornblende by Tschermak substitution. Our amphiboles plot on a field corresponding to those from amphibolites accompanied by sillimanite-bearing assemblages in metapelitic rocks of LAIRD and ALBEE (1981b), Fig. 3b. In an ACF-diagram, the amphiboles plot close to the bulk rock composition.

Tab. 2 Mineral compositions (explanation of sample numbers: see text)

A: Chemical composition of amphiboles

Sample	1554 ¹⁾		1523		1515 ¹⁾		1507 ¹⁾	
	1604 ¹⁾	rim ¹⁾	1523	Core ²⁾	1515	1507	1523	1515
SiO ₂	46.58	44.78	45.30	51.50	45.89	47.52	55.88	60.50
Al ₂ O ₃	13.61	12.40	10.89	4.74	11.06	8.02	10.25	7.33
TiO ₂	0.00	0.75	0.00	0.00	0.70	0.00	28.24	25.93
Cr ₂ O ₃	0.13	0.08	0.00	0.00	0.00	0.00	6.32	7.28
FeO	12.30	15.79	14.32	14.10	15.01	15.69	0.08	0.03
MgO	11.96	10.12	10.90	12.87	11.13	11.61	100.78	101.07
MnO	0.09	0.15	0.35	0.31	0.29	0.27		
CaO	10.81	11.16	11.46	11.56	11.71	11.82		
Na ₂ O	1.36	1.26	1.15	0.50	1.43	1.16		
K ₂ O	0.35	0.21	0.18	0.09	0.10	0.12		
H ₂ O stoic.	2.30	2.30	2.30	2.30	2.30	2.30		
Total	99.49	99.00	96.85	97.97	98.92	99.21		

C: Chemical composition of garnets

Sample	1604		1544		1507	
	core	rim	core	rim	core	rim
SiO ₂	38.35	38.10	38.27	38.28	38.86	38.86
Al ₂ O ₃	20.64	20.94	20.82	20.86	21.40	21.40
Cr ₂ O ₃	0.10	0.00	0.14	0.05	0.05	0.05
FeO	27.65	30.08	26.51	22.34	22.72	22.72
MnO	1.35	0.81	1.46	2.72	3.11	3.11
MgO	1.70	3.06	1.83	2.34	2.21	2.21
CaO	10.89	8.17	12.10	13.56	13.39	13.39
Total	100.68	101.16	101.13	100.15	101.74	101.74

B: Chemical composition of plagioclases

Sample	1554		1604		1523		1515		1507	
	SiO ₂	56.57	59.82	59.82	55.88	60.50	53.19			
CaO	9.90	7.88	7.88	10.25	7.33	12.40				
Al ₂ O ₃	27.02	25.80	25.80	28.24	25.93	29.84				
Na ₂ O	5.93	7.30	7.30	6.32	7.28	4.64				
K ₂ O	0.04	0.11	0.11	0.08	0.03	0.04				
Total	99.46	100.91	100.78	101.07	100.11					

Formula¹⁾:

Si	3.030	2.990	3.000	3.005	3.006
Al ^{IV}	0.000	0.010	0.000	0.000	0.000
Sum T	3.030	3.000	3.000	3.005	3.006
Al ^{VI}	1.953	1.927	1.925	1.935	1.958
Fe ³⁺ ²⁾	0.040	0.073	0.066	0.062	0.039
Sum	1.993	2.000	1.991	1.997	1.997
Fe ²⁺	1.787	1.901	1.673	1.405	1.431
Mn	0.090	0.054	0.097	0.181	0.204
Mg	0.200	0.359	0.214	0.274	0.255
Ca	0.922	0.689	1.017	1.140	1.110
Sum	2.999	3.003	3.001	3.000	3.000
Mol% Almandin	58.2	61.1	53.7	45.4	46.7
Mol% Pyrope	6.5	11.5	6.9	8.8	8.3
Mol% Gross. ³⁾	30.0	22.1	32.7	36.8	36.2
Mol% Spessar.	2.9	1.7	3.1	5.8	6.7
Mol% Andrad.	2.0	3.5	3.2	3.0	1.9
Mol% Uvarov.	0.3	0.0	0.4	0.2	0.2

Formula¹⁾:

Si	2.55	2.65	2.50	2.66	2.40
Al	1.44	1.35	1.49	1.35	1.59
Sum	3.99	4.00	3.99	3.01	3.99
Ca	0.48	0.37	0.49	0.35	0.60
Na	0.52	0.63	0.55	0.62	0.41
K	0.00	0.01	0.00	0.00	0.00
Sum	1.00	1.01	1.04	0.97	1.01
%An	48	37	45	33	60

¹⁾: normalized to 8 oxygen atoms.

²⁾: normalized to Σ cations-Ca-Na-K = 13 (Na can be on A- and M4-site);

³⁾: normalized to Σ cations-K = 15 (only K on A-site);

⁴⁾: calculated using charge balance (LAIRD and ALBEE, 1981a).

Analyst: R. Oberhänsli

Analyst: R. Oberhänsli

Analyst: R. Oberhänsli

PLAGIOCLASE

Plagioclase is the most abundant phase after amphibole with an overall range of An₃₀ to An₆₀ in which andesine compositions dominate. The variation of individual minerals is more restricted (Fig. 3b). Based on only few analyses (Tab. IIB), it seems that plagioclases from the plagioclase amphibolites have slightly higher An-contents than those from the garnet-amphibolites.

GARNET

Two generations of garnet are observed in type 1 amphibolites. Their garnets display a chemical zonation where the earlier generation preserved in the center of large garnets (e.g. sample 1604, Tab. IIC) corresponds to a grossularite-rich almandine with the general formula of $\text{Alm}_{58}\text{Py}_7\text{Gr}_{30}\text{Sp}_3\text{Andr}_2$. It has a similar composition to the garnets from the amphibole-garnet-schists (type [4], sample 1544). The rims of the large garnets of rock type (1A) have a general composition of $\text{Alm}_{61}\text{Py}_{12}\text{Gr}_{22}\text{Sp}_2\text{Andr}_3$.

Rock chemistry

DATA

A set of 55 samples has been analyzed by X-ray fluorescence for 30 major and trace elements. Tab. 3 contains representative analyses for the main rock

types and Fig. 4 shows selected elements plotted against the Mg-number ($\text{Mg}/(\text{Mg} + \text{Fe}^{2+})$). On the basis of the latter, three fairly well distinguishable chemical groups can be recognized, corresponding respectively to the plagioclase-amphibolites of group 2 with high Mg-numbers, to the garnet-amphibolites with relictic garnets of group 1C with medium Mg-numbers, and to the coarse-grained garnet-amphibolites of group 1A with low Mg-numbers. The garnet-amphibolites are in general poorer in Ni but richer in Fe, V and most incompatible elements like P, Ti, Zr and Y than the plagioclase-amphibolites. The garnet-amphibolite sample 1527 (belonging to group 1C and collected from a small amphibolite boudin of only 50 cm, surrounded by granitoid gneisses) shows a compositional trend approaching the garnet-amphibolites of group 1A.

Rare earth elements (REE) of 20 samples have been analyzed by instrumental neutron activation (Tab. 4, Fig. 5), including samples that show considerable replacement of plagioclase by sericitic white mica. The fresh, non-sericitized samples show, aside from mainly negative Eu-anomalies, flat patterns at an average enrichment level of 15 to 20 in comparison to chondrites (Fig. 5 a, b). Some of the garnet-amphibolites are slightly more enriched (20 to 30 times chondrites). The garnet-amphibolite sample no. 1527 displays a rather horizontal pattern with a general enrichment of about 40 for all REE.




Fig. 3 Mineral composition of Amphiboles.

A: Plot of the formula proportions of Al^{VI} , Fe^{3+} , Ti and Cr against Al^{IV} . The dotted line links the composition of core and rim of a zoned amphibole in sample 1523. Abbreviations: Act: actinolite, Ts: tschermakite, Par: pargasite, Gl: glaucophane, Ed: edenite, Win: winchite, Bar: barrosite.

B: Major element trends of amphiboles in a diagram proposed by LAIRD and ALBEE (1981b) for mafic schist in Vermont, with compositional fields of their amphiboles occurring with different minerals of mafic and pelitic rocks: oligoclase in mafic rocks (shaded area); sillimanite (limited by dashed lines) and garnet (solid lines) in pelitic rocks. Numbers indicate plagioclase composition (% anorthite) for a given rock. The amphiboles plot in the sillimanite field, confirming their upper amphibolite facies character. The dotted line links the composition of core and rim of a zoned amphibole in sample 1523. Sample-numbers: see text.

Tab. 3 Bulk rock composition of the analyzed amphibolites (% for oxides, ppm for traces).

Element	Type 1A	Type 1B	Type 1C		Type 1C	Type 2	
	1604	1507	Fresh (14)*	repl. (7)*	contam. 1527	fresh (13)*	repl. (9)*
SiO ₂	50.58	49.98	49.17	50.11	52.67	48.62	48.07
TiO ₂	2.83	1.25	1.31	1.41	2.13	0.99	1.00
Al ₂ O ₃	13.62	15.11	15.63	16.09	13.28	16.11	16.65
Fe ₂ O ₃	2.88	1.98	1.85	1.61	3.31	1.76	1.47
FeO	11.80	8.43	8.72	8.80	9.65	7.70	7.74
MnO	0.28	0.20	0.20	0.23	0.19	0.19	0.19
MgO	5.99	7.99	8.24	7.50	6.74	9.51	9.53
CaO	8.69	10.64	10.16	8.47	6.70	10.07	9.82
Na ₂ O	1.74	2.43	2.18	1.21	2.24	2.13	1.41
K ₂ O	0.19	0.38	0.47	1.32	0.97	0.56	1.13
P ₂ O ₅	0.29	0.14	0.16	0.17	0.30	0.09	0.10
H ₂ O ⁺	0.58	1.27	1.52	2.40	1.80	2.05	2.56
Total	99.47	99.79	—	—	99.98	—	—
Zr	173.	90.	86.	95.	293.	62.	61.
Y	46.	21.	22.	25.	77.	17.	17.
Sr	123.	300.	256.	231.	113.	217.	241.
Rb	3.	18.	18.	81.	29.	23.	49.
Pb	11.	11.	10.	10.	11.	12.	11.
Ga	23.	15.	16.	17.	21.	14.	13.
Zn	150.	90.	89.	93.	217.	102.	102.
Cu	70.	99.	64.	41.	60.	39.	58.
Ni	45.	52.	61.	50.	56.	106.	120.
Co	50.	43.	45.	44.	45.	44.	43.
Cr	135.	134.	240.	156.	179.	438.	411.
V	326.	217.	216.	205.	237.	180.	167.
Ce	54.	40.	16.	22.	55.	16.	17.
Nd	53.	7.	23.	18.	56.	19.	9.
Ba	110.	164.	178.	619.	216.	209.	401.
La	29.	6.	9.	11.	21.	7.	4.
Sc	62.	51.	51.	52.	54.	48.	44.
S	508.	103.	388.	503.	2325.	14.	107.

* mean values, in parenthesis: number of averaged samples. "repl.": extensive replacement of plagioclase by sericitic white mica.

Methods used: Major elements analyzed on fused beads (sample : Li₂BO₃ = 1 : 5) and corrected for matrix effects, absorption etc. with fundamental alpha-factors (DE JONGH, 1979). Traces analyzed on pressed powder pellets according to NISBET et al. (1979). FeO determined with colorimetric methods (WILSON, 1960). H₂O calculated from ignition loss.

Analyst: G. Galetti (major elements), J.C. Lavanchy (traces).

The complete data set of 55 samples is available from the Artemise data bank of the Centre de Recherches Pétrographiques et Géochimiques du CNRS at Nancy, France.

METASOMATIC CHANGES

Before trying to determine possible protoliths and their paleotectonic environment, evident metasomatic changes during the certainly complex history have to be considered and discussed.

The most obvious alteration is the replacement of plagioclase grains by fine grained white mica (sericite). The amphibolites with sericitized plagioclases ("repl." in Tab. 3) show a loss of CaO, Sr

and Na₂O and the enrichment in K₂O, Rb, Ba and S relative to the corresponding amphibolites with well-preserved plagioclases ("fresh" in Tab. 3). V, Nd and Cr are slightly diminished and Tb is strongly enriched in the sericitized samples. Two explanations are offered: either the protoliths underwent an early postdepositional transformation or the amphibolites suffered a later (Permian or Tertiary-alpine) alteration. HUMPHRIS (1984) describes enrichment in Tb and La from basalts that have been




Fig. 4 Bulk rock composition of amphibolites from the Emosson area as a function of the Mg-number ($Mg/(Mg+Fe^{2+})$), calculated with the assumption that $Fe_2O_3/FeO = 0.1$, the mean ratio for unaltered basalts). Diamonds: garnet-plagioclase-amphibolites (type 1A). Open circles: garnet-plagioclase-amphibolites with relic garnets (type 1C). Triangles: plagioclase-amphibolites (type 2). Filled circle: garnet-amphibolite no. 1527 from a small amphibolite boudin.

Tab. 4 Rare earth analyses of selected 20 amphibolite samples (second line: rock types, c.f. chapter petrography of the text; r = plagioclase replaced by sericitic white mica; cont.: contaminated by country rock).

	1505 1c	1514 1c	1515 1c	1519 1c	1520 1c	1521 1c	1523 1c	1527 cont	1509 1c-r	1512 1c-r
La	11.21	12.24	10.89	7.10	6.16	6.97	3.17	12.93	10.52	10.90
Ce	16.48	17.71	16.95	15.50	16.58	15.81	10.02	31.52	16.44	14.48
Nd	12.35	14.08	13.07	9.15	9.29	10.38	8.48	18.01	6.35	11.94
Sm	5.02	4.84	4.90	3.03	3.20	3.01	3.13	8.19	4.58	4.63
Eu	1.27	1.18	1.13	0.18	1.08	1.12	0.94	2.27	1.12	1.06
Tb	1.19	2.84	1.31	0.55	1.10	1.19	1.19	2.06	2.94	1.17
Yb	3.80	3.60	4.25	2.10	2.49	2.34	3.98	8.82	3.41	4.46
Lu	0.62	0.49	0.53	0.29	0.30	0.35	0.48	1.13	0.49	0.59
SREE	51.94	56.98	53.03	37.90	40.20	41.37	31.41	84.93	50.33	49.23
La/Sm	2.23	2.53	2.22	2.34	1.93	2.32	1.02	1.58	2.30	2.35
La/Yb	2.95	3.40	2.56	3.38	2.47	2.98	0.80	1.47	3.09	2.44

	1502 2	1511 2	1518 2	1526 2	1532 2	1540 2	1545 2	1547 2	1503 2-r	1504 2-r
La	4.34	4.22	11.64	2.95	4.52	3.62	3.14	3.77	4.61	4.52
Ce	13.26	6.74	19.86	7.48	9.85	8.48	9.08	7.91	13.18	9.03
Nd	8.75	6.35	14.17	7.94	9.42	9.69	8.47	6.94	11.11	9.63
Sm	2.48	2.65	4.92	1.89	2.38	1.98	2.40	2.14	2.50	3.56
Eu	1.13	0.66	0.83	0.86	1.39	0.83	0.18	0.83	1.00	0.94
Tb	1.19	0.24	1.47	1.49	1.09	0.81	0.58	1.06	0.77	1.87
Yb	2.10	2.75	4.61	2.15	2.33	2.07	2.17	1.87	2.24	3.37
Lu	0.24	0.37	0.50	0.27	0.29	0.24	0.32	0.30	0.28	0.49
SREE	33.49	23.98	58.00	25.03	31.27	27.72	26.34	24.82	35.69	33.41
La/Sm	1.75	1.59	2.37	1.56	1.90	1.83	1.31	1.76	1.84	1.27
La/Yb	2.07	1.53	2.52	1.37	1.94	1.75	1.45	2.02	2.06	1.34

R. A. Oliver and G. Vivier (Grenoble) analysts; INA analysis; uncertainties: La = $\pm 5\%$; Ce = $\pm 5\%$; Nd = $\pm 15\%$; Sm = $\pm 5\%$; Eu = $\pm 10\%$; Tb = $\pm 10\%$; Yb = $\pm 5\%$; Lu = $\pm 10\%$.

either weathered or zeolitized by oceanic metamorphism. As the trend of Tb-enrichment is also visible in the other samples, most samples have probably been slightly influenced by circulating fluids and/or weathering. However the most probable cause of the formation of sericite are the low grade hydrothermal alterations of Permian age, during which sulfides were precipitated along local fault zones in the whole Aiguilles Rouges massif (Fig. 1, cf. also VON RAUMER, 1987).

The direct chemical exchange with the adjacent metapelitic country rock is best illustrated with two samples from the marginal parts of the amphibolite boudins (no. 1527 and 1604). The garnet amphibolite 1527 had been collected in a small boudin surrounded by granitoid gneisses. It contains biotite in small amounts, while the normal garnet-amphibolites have none. Consequently, it is enriched in SiO₂, TiO₂, total Fe, K₂O, Rb, P₂O₅, Zn, Ce, Ba, S, Y, Zr and REE, whereas Al₂O₃, MgO, CaO, Sr and Cr





were depleted. It has obviously interacted with the adjacent metagreywackes and has been contaminated, including the formation of biotite. Similar tendencies are found for all 3 samples of the garnet-amphibolite group 1A (example: sample 1604), all collected from the limit between amphibolite and surrounding high grade metapelites. The strong zonation of garnets of the same samples, described above, are interpreted as a metasomatic change as well. Most likely these compositional changes occurred during the decompressional phase of the metamorphism, during which fluids may have had partial access to the rocks in question.

Taking into account all these secondary changes discussed above, there are 27 samples left that show no evidence for metasomatic changes. They all have been sampled in the central part of large amphibolite bodies. Only the non-sericitized samples have been plotted on diagrams of Figs 6 to 8 and have been retained for a discussion of their origin.

Origin of the amphibolites

POSSIBLE PROTOLITHS

Despite several attempts to chemically define metasedimentary amphibolites (FRÖHLICH, 1960;



LEAKE, 1964; MOINE, 1969) based on major and common trace elements, no satisfactory scheme exists to discriminate metagabbros or metabasalt rocks from metamarls. Studies of magmatic amphibolite-ultramafic rock series lying in probable metasedimentary amphibolite-bearing series (SCHLÄPFER, 1979; EVANS et al. 1981; SCHMIDT, 1989), show a very limited trace element variation close to modern oceanic basalts for the supposedly magmatic amphibolites and a irregular trace element distribution for the probably metasedimentary amphibolites. As can be seen from Figs 4 to 8, the Emosson amphibolites are quite well grouped in most diagrams. According to FRÖHLICH (1960), who found contents larger than 150 ppm of Cr and Cr/Ni ratios higher than 1 for typical magmatic mafic rocks, 80% of our samples are of magmatic





Fig. 5 REE abundance patterns of selected samples normalized to chondrite of NAKAMURA (1974), Tb: HASKIN et al. (1968) as reported in HENDERSON (1984), Tab. 1.5.

A: garnet-amphibolites. B: plagioclase-amphibolites. C: Amphibolites showing extensive replacement of plagioclase by sericitic white mica. D: Reference patterns of different tholeiitic basalts. 1) normal mid ocean ridge basalt (MORB) after PEARCE (1982). 2) transitional (T)-type MORB and plume (P)-type MORB after WOOD et al. (1979b, sample 411 and 413-1). 3) Continental tholeiite after DOSTAL et al. (1983, sample 718).




Fig. 6 Major element composition of the amphibolites (possibly metasomatized samples excluded). A: AFM-diagram, A = Na₂O + K₂O; F = FcO_{tot}; M = MgO; field 1: garnet amphibolites (types 1B, 1C), field 2: plagioclase amphibolites (type 2), diamonds: group 1A with large garnets. Dashed line: Skaergaard evolution-trend after WAGER (1960). B: TiO₂-Al₂O₃-diagram after PEARCE (1984), rectangle: field of basaltic liquids. C: R1-R2 diagram after DE LA ROCHE et al. (1980), modified by BELLIENI et al. (1984), ob = olivine-basalts; th = tholeiites; ab = andesitic basalt. D: CIPW normative composition of the amphibolites plotted in the projection diopside-olivine-hypersthene-quartz. Diamonds: garnet-amphibolites with large garnets (type 1A), open circles: garnet-plagioclase-amphibolites with relic garnets (type 1C), triangles: plagioclase-amphibolites (type 2), full circle: garnet-amphibolite sample no. 1527.

origin. The remaining samples plot in the ambiguous domain between 20 and 150 ppm of Cr. The following additional arguments are all consistent with a magmatic origin: most samples lie within the basaltic liquid field of PEARCE (1984), Fig. 6b. The two main groups (1C and 2) are tholeiitic (Figs 6c, 7, 8). Altogether, our samples exhibit a compositional trend strongly suggesting a magmatic differentiation series (cf. Fig. 4; see discussion below).

PFEIFER *et al.* (1989) demonstrated that the original grain size (fine-grained basalts, coarse-grained gabbros), at least for Mesozoic rocks, was largely preserved even during polyphase high grade metamorphism. The fine grain size of our




Fig. 7 Amphibolites plotted in the discrimination diagram of FLOYD and WINCHESTER (1975). alc: alkaline basalts, thol: tholeiitic basalts. Meaning of symbols: as in Fig. 6.

samples therefore suggests volcanic or subvolcanic rather than plutonic protoliths. In addition, the TiO_2 contents ($> 0.5\%$, Fig. 6b) and the V-contents (> 100 ppm) exclude a gabbroic origin (PFEIFER *et al.*, 1989).

When comparing common trace element behavior in tholeiitic rocks (PRINZ, 1967; CARMICHAEL *et al.*, 1974) to the amphibolites from Emosson, we can draw the following conclusions (Tab. 3): elevated values of Cr and Ni are observed in the plagioclase amphibolites, which could represent early differentiates. The garnet-plagioclase-amphibolites are enriched in V and P suggesting that they are more evolved than the plagioclase-amphibolites, which consequently must have formed prior to crystallization of magnetite and apatite. Also the elevated concentrations of Zr and

Ti indicates that garnet amphibolites are more evolved. Thus, the indications of the Mg-number are confirmed by the trace elements.

Compared to modern tholeiitic lavas, it appears as if the plagioclase-amphibolites were relatively undifferentiated normal spinel-olivine-tholeiites and the garnet-plagioclase-amphibolites more evolved hypersthene- or even quartz-tholeiites, a tendency which appears again in the CIPW-norm (Fig. 8), where most rocks are olivine-diopside-hypersthene-normative and only few have norms containing quartz.

PALEOTECTONIC ENVIRONMENT

An attempt to determine the paleotectonic environment can be made based mainly on the least mobile trace elements, most REE, Ti, Zr, Y, V and possibly Ni and Cr. Fig. 8 contains the usually employed classical trace element diagrams (cf. PEARCE, 1984). In most of them (Fig. 8 a, b, d, f) the two major groups of amphibolite fall into the field of mid ocean ridge basalts (MORB), but in some they fall onto the limit between MORB and island arc tholeiites (Fig. 8 c, e). Especially the plagioclase-amphibolites, with their more primitive character, have unusually low Zr and Y values and plot outside the MORB field. A similar tendency has been reported from amphibolites from the Belle-donne massif in France (MÉNOT, 1987). Since there is no indication of a metasomatic event, we interpret this as a primitive character, typical for subvolcanic rocks with a trace element composition between gabbroic and basaltic volcanic rocks (e.g. two samples from a sill at Wright Valley, Antarctica reported by THOMPSON *et al.*, 1983; PAPEZYK and HODYCH, 1980; or ophiolitic dikes in PFEIFER *et al.*, 1989). Fig. 9 (normalization to normal MORB compositions) contains a comparison with different types of MOR-basalts and continental tholeiites. They show once more the great compositional span of certain elements within the same group (especially Rb, Ba, K), but otherwise indicate a close similarity to enriched plume-type MORB and continental tholeiites. For the latter, DUPUY and DOSTAL (1984) and WEAVER and TARNEY (1983) report highly variable element ratios, leading to the typical humps in the MORB-normalized diagrams and this is exactly what is observed in the Emosson amphibolites (Figs 9 b, c). However, the chondrite-normalized diagram indicates, with its rather flat patterns, a closer affinity to less enriched transitional MOR basalts (Fig. 5d). Such apparently contradictory results have been reported from continental rift zones with strong extension leading to the formation of oceanic crust (BER-




Fig. 8 Classical trace element diagrams to determine the paleotectonic environment of basalts applied to the amphibolites. 1: garnet-amphibolites, 2: plagioclase amphibolites. Symbols see Fig. 6.

Abbreviations: WPB: within plate basalts; MORB: mid ocean ridge basalts; IAT: island arc tholeiites; OIB: ocean island basalts; LKT: low potassium tholeiites; CAB: calc-alkaline basalts.

(A): PEARCE and CANN (1973): Most samples (black area) plot into the field B (ocean floor basalts, low potassium tholeiites); dot: contaminated amphibolite 1527. A = LKT; B = CAB + LKT; C = CAB; D = WPB. (B): PEARCE (1980): The amphibolites plot into the field of MORB, but they are clearly separated into the three main rock types: 1: plagioclase amphibolites, 2: normal garnet-amphibolites and garnet-amphibolites with large garnets (group 1A; diamonds). (C): PEARCE and NORRY (1979): the amphibolites plot into the field of MORB, but they are clearly separated into the three main rock types. (D): SHERVAIS (1982): the amphibolites follow a line parallel to the Ti/V ratio of 50, that separates the field of MORB and the field of OIB-WPB. Plagioclase-amphibolites (group 2) plot closer to the origin of the diagram than the group of garnet amphibolites. (E): PEARCE (1980): the different amphibolites are more or less separated, but no special attribution is possible. (F): BECCALUVA et al. (1983): the plagioclase-amphibolites (group 2) plot between the island arc tholeiites (IAT) and ocean floor tholeiites (OFT), whereas most garnet amphibolites (type 1C and also type 1A, diamonds) plot into the area of ocean floor tholeiites.




Fig. 9 Trace elements of the different amphibolites compared to typical oceanic and continental tholeiites.

A-C: Normalized with the primordial mantle composition of WOOD et al. (1979a).

D-F: Normalized with the N-MORB composition of PEARCE (1980). References: 1) normal mid-ocean ridge basalt (MORB) after PEARCE (1980). 2) transitional (T-)type MORB and plume (P-) type MORB after WOOD et al. (1979b, sample 411 and 413-1). 3) Continental tholeiite after DOSTAL et al. (1983, sample 718).

TRAND et al., 1982; COX, 1983; BELLINI et al., 1984; UPTON et al., 1984; FODOR and VETTER, 1984; COLEMAN and MCGUIRE, 1988).

Conclusions

The studied amphibolites have suffered a poly-phase high-grade metamorphism of Caledonian and Variscan age (VON RAUMER 1984b, 1987, 1988). After the early formation of eclogites the metabasic rocks followed the Barrovian evolution trend involving decompression and a temperature increase as indicated by the petrology of the accompanying metapelites. During this second metamor-

phic event, limited chemical exchange took place between the metapelitic country rocks and the small amphibolite boudins and the marginal zones of larger ones. During hydrothermal activity of Permian age, plagioclases were partly replaced by sericitic white mica. In contrast the Tertiary overprint was almost negligible (local pumpellyite-prehnite).

Despite this complex metamorphic history, the described amphibolites have preserved a chemical pattern interpreted as an igneous differentiation series of transitional MORB characteristics as is typical for continental rift zones at the onset of ocean floor spreading. As the volume of the mafic rocks is small and occurs as several hundred-meter-

long thin layers, it is very likely that the studied mafic rocks were sills or dikes which intruded detrital sediments of continental character (today metapsammitic and metapelitic gneisses and schists). Their unusual high contents in K, Rb and Ba can be explained by enrichment from the surrounding wall (FRATTA and SHAW, 1974; WEAVER and TARNEY, 1981; GILL and BRIDGWATER, 1979; GOLDBERG et al., 1986). Magmatic differentiation from olivine-tholeiite to hypersthene-tholeiite is well known from diabase dikes (PAPEZYK and HODYCH, 1980). Magma chilling at the contact and later crystallization of the central parts may explain the differences in composition.

The overall chemical variation of the described amphibolites with its transitional MORB trend, but enriched in K, Rb and Ba is typical for continental rift zones characterized by sub-alkalic basalts (WILSON, 1988) and has been hypothesized for the Aiguilles Rouges massif by PAQUETTE et al. (1989). A model of the inferred paleotectonic environment

during the formation of the basaltic rocks of Lake Emosson is presented in Fig. 10. It corresponds to the tentative model of the External Massifs in general by VON RAUMER (1987) and is also in good agreement with the general picture of the Paleozoic Central Europe as proposed by ZWART and DORN-SIEPEN (1980).

Acknowledgements

Several people helped us with their analytical skills and their experience to operate computers. We are especially indebted to J.C. Lavanchy (Lausanne) who provided the XRF trace elements and patiently helped to draw and redraw several generations of diagrams. We thank A. Colombi for having let us use his mineral normalization programs. REE analyses were performed thanks to O. Richards and G. Vivier (Grenoble). A. Ploquin (Nancy) has kindly incorporated our data in the Artemise data bank. With his constructive remarks on an early version, J. Pearce contributed a lot to this manuscript. In addi-




Fig. 10 Diagrammatic sections illustrating the hypothetical paleotectonic situation of the Emosson amphibolites as former tholeiitic eruptions in an extending continental crust, following the models of FODOR and VETTER (1984) and UPTON et al. (1984) and the paleotectonic interpretations (inset) of VON RAUMER (1987).

tion, we profited from the critical remarks and suggestions of Z. Sharp (Lausanne), B. Messiga and R. Tribuzio (Pavia). This research has been supported by the Swiss National Science Foundation Project no. 2.610-0.85.

References

- BECCALUVA, L., DI GIROLAMO, P., MACCIOTTA, G. and MORRA, V. (1983): Magma affinities and fractionation trends in ophiolites. *Ophioliti* 8, 307-324.
- BELLIENI, G., COMIN CHIARAMONTI, P., MARQUES, L.S., MEILI, A.J., PICIRILLO, P.M., NARDY, A.J.R. and ROISENBERG, A. (1984): High- and low-TiO₂ flood basalts from the Parana plateau (Brazil): petrology and geochemical aspects bearing on their mantle origin. *N. Jahrb. Mineral. Abhandl.* 150, 273-306.
- BERTRAND, H., DOSTAL, J. and DUPUY, C. (1982): Geochemistry of early Mesozoic tholeiites from Morocco. *Earth. Planet. Sci. Lett.* 58, 225-239.
- CARMICHAEL, I.S.E., TURNER, F.J. and VERHOOGEN, E.J. (1974): *Igneous Petrology*, p. 480, McGraw Hill, New York.
- COLEMAN, R.G. and MCGUIRE, A.V. (1988): Magma systems related to the Red Sea rift system. *Tectonophysics* 150, 77-101.
- COLLET, L.W., OULIANOFF, N. and REINHARD, M. (1952): Note explicative de la feuille 24 de l'Atlas, Finhaut. *Geologischer Atlas der Schweiz* 1 : 25000, Kümmerly & Frey, Bern.
- COLOMBI, A. (1990): Métamorphisme et géochimie des roches mafiques des alpes ouest-centrales (Géoprofil Viège-Domodossola-Locarno). *Mémoires de Géologie (Lausanne)* 4, 216 p.
- COX, G.C. (1983): The Karoo Province of Southern Africa: origin of trace element enrichment patterns. In: HAWKESWORTH, C.J. and NORRY, M.J. (eds), *Continental basalts and mantle xenoliths*, 139-157, Shiva, Nantwich.
- DE JONGH, W.K. (1979): The atomic number $z = 0$: loss and gain on ignition in XRF analysis treated by the IN-equation. *X-ray spectrometry* 8, 52-56.
- DE LA ROCHE, H., LETERRIER, P., GRANDCLAUDE, P. and MARCHAL, M. (1980): A classification of volcanic and plutonic rocks using R1-R2 diagrams and major-element analyses. Its relationship with current nomenclature. *Chem. Geol.* 29, 183-210.
- DOSTAL, J., DUPUY, C. and BARAGER, W.R.A. (1983): Geochemistry and petrogenesis of basaltic rocks from Coppermine River area, Northwest Territories. *Can. J. Earth. Sci.* 20, 684-698.
- DUPUY, C. and DOSTAL, J. (1984): Trace elements geochemistry of some continental tholeiites. *Earth and Planet. Sci. Lett.* 67, 61-69.
- EVANS, B.W., TROMMSDORFF, V. and GOLES, G.G. (1981): Geochemistry of high-grade eclogites and metaroddingites from the Central Alps. *Contrib. Mineral. Petrol.* 76, 301-311.
- FLOYD, P.A. and WINCHESTER, J.A. (1975): Magma type and tectonic setting discrimination using immobile elements. *Earth and Planet. Sci. Lett.* 27, 211-218.
- FODOR, R.V. and VETTER, S.K. (1984): Rift-zone magmatism: Petrology of basaltic rocks transitional from CFB to MORB, southeastern Brazil margin. *Contr. Mineral. Petrol.* 88, 307-321.
- FRATTA, M. and SHAW, D.M. (1974): "Residence" contamination of K, Rb, Li and Tl in diabase dikes. *Canad. J. Earth. Sci.* 11, 422-429.
- FRÖHLICH, F. (1960): Ein Beitrag zur Geochemie des Chroms. *Geochim. Cosmochim. Acta* 20, 215-240.
- GILL, R.C.O. and BRIDGWATER, D. (1979): Early archean basic magmatism in West Greenland: the geochemistry of the Ameralik dykes. *J. Petrol.* 20, 695-726.
- GOLDBERG, S.A., BUTLER, J.R. and FULLAGAR, P.D. (1986): The Bakersville dike swarm: geochronology and petrogenesis of late Proterozoic basaltic magmatism in the Southern Appalachian Blue Ridge. *Amer. J. Sci.* 286, 403-430.
- HASKIN, L.A., HASKIN, M.A., FREY, F.A. and WILDEMAN, T.R. (1968): Relative and absolute terrestrial abundances of the rare earths. In: AHRENS, L.H. (Ed.), *Origin and distribution of the elements*. Pergamon Oxford, pp. 889-911.
- HENDERSON, P. (1984): General geochemical properties and abundances of the rare earth elements. In: Henderson, P. (Ed.), *Rare earth element geochemistry*, 1-32. Elsevier Amsterdam.
- HUMPHRIS, S.E. (1984): The mobility of the rare earth elements in the crust. In: HENDERSON, P. (ed.), *Rare earth element geochemistry*, 317-342. Elsevier Amsterdam, 410 p.
- JOYE, J.B. (1989): L'évolution pression - température - déformation dans le massif des Aiguilles Rouges, massif externe alpin. Thèse Université Fribourg, 1-135.
- LAIRD, J. (1978): AMPH3, a FORTRAN IV program for the normalization of amphiboles and pyroxenes. Manuscript Program libr. Petrology, ETH-Zürich.
- LAIRD, J. and ALBEE, A.L. (1981a): High pressure metamorphism in mafic schist from Northern Vermont. *Amer. J. Sci.* 281, 97-126.
- LAIRD, J. and ALBEE, A.L. (1981b): Pressure, temperature, and time indicators in mafic schist: their application to reconstructing the polymetamorphic history of Vermont. *Amer. J. Sci.* 281, 127-175.
- LEAKE, B. (1964): The chemical distinction between ortho- and paraamphibolites. *J. petrol.* 5, 238-254.
- LEAKE, B.E. (1978): Nomenclature of amphiboles. *Amer. Mineral.* 63, 1023-1052.
- LIÉGOIS, J.P. and DUCHESNE, J.C. (1981): The Lac Cornu retrograded eclogites (Aiguilles Rouges Massif, Western Alps, France): evidence of crustal origin and metasomatic alteration. *Lithos* 14, 35-48.
- MÉNOT, R.-P. (1987): Magmatismes paléozoïques et structuration carbonifère du massif de Belledonne (Alpes Françaises). Contraintes nouvelles pour les schémas d'évolution de la chaîne varisque ouest-européenne. Ph.D. diss. Univ. Claude Bernard of Lyon I.
- MOINE, B. (1969): Orthoamphibolites et formations métasédimentaires calco-magnésiennes de la région d'Ambatofinandrahana (Madagascar). Contribution au problème de l'origine des amphibolites. *Sci. de la Terre* XIV/2, 111-138.
- NAKAMURA, N. (1974): Determination of REE, Ba, Fe, Mg, Na and K in carbonaceous and ordinary chondrites. *Geochim. Cosmochimica Acta* 38, 757-775.
- NISBET, E., DIETRICH, V. and ESENWEIN, A. (1979): Routine trace element determination in silicate minerals and rocks by X-ray fluorescence. *Fortschr. Mineral.* 57, 264-279.
- PAPEZYK, V.S. and HODYCH, J.P. (1980): Early Mesozoic diabase dikes of the Avalon Peninsula, Newfoundland: Petrochemistry, mineralogy and origin. *Canad. J. Earth. Sci.* 17, 1417-1430.
- PAQUETTE, J.L., MÉNOT, R.P. and PEUCAT, J.J. (1989): REE, Sm-Nd and U-Pb zircon study of eclogites from the Alpine External Massifs (Western Alps):

- Evidence for crustal contamination. *Earth and Planet. Sci. Lett.* 96, 181–198.
- PEARCE, J.A. (1980): Geochemical evidence for the genesis and eruptive setting of lavas from Tethyan ophiolites. *Proc. int. Ophiolite Conference, Nicosia, Cyprus*. PANAYIATOU, N. (ed.), pp. 261–272.
- PEARCE, J.A. (1982): Trace element characteristics of lavas from destructive plate boundaries. In: THORPE, R.S., ed., *Andesites*, 525–548, Wiley, New York.
- PEARCE, J.A. (1984): A "users guide" to basalt discrimination diagrams. Unpublished manuscript. The Open University, Milton Keynes, 37 p.
- PEARCE, J.A. and CANN, J.R. (1973): Tectonic setting of basic volcanic rocks determined using trace element analyses. *Earth and Planet. Sci. Lett.* 19, 290–300.
- PEARCE, J.A. and NORRIS, M.J. (1979): Petrogenetic implications of Ti, Zr, Y and Nb variations in volcanic rocks. *Contrib. Mineral. Petrol.* 69, 33–47.
- PEIFER, H.-R., COLOMBI, A. and GANGUIN, J. (1989): Zermatt-Saas and Antrona zone: a petrographic and geochemical comparison of polymetamorphic ophiolites of the West-central Alps. *Schweiz. Mineral. Petrogr. Mitt.* 69, 217–236.
- PRINZ, M. (1967): Geochemistry of basaltic rocks: Trace elements. In: HESS, H.H. and POLDERVAART, A., eds, *Basalts: the Poldervaart treatise on rocks of basaltic composition*, 271–323, Interscience Publishers.
- VON RAUMER, J.F. (1974): Zur Metamorphose amphibolischer Gesteine im Altkristallin des Mont-Blanc und des Aiguilles-Rouges-Massivs. *Schweiz. Mineral. Petrogr. Mitt.* 54, 471–488.
- VON RAUMER (1976): Variszikum in den Zentral- und Westalpen. *Nova Acta Leopoldina N.F.* 224, 45, 119–144.
- VON RAUMER (1981): Variscan events in the Alpine region. *Geologie en Mijnbouw* 60, 67–80.
- VON RAUMER (1984a): The external massifs, relics of Variscan basement in the Alps. *Geol. Rdsch.* 73, 1–31.
- VON RAUMER (1984b): The Hercynian basement in the Helvetic Realm (Western and Central Alps). In: DAL PIAZ, G.V., ed., *Alpi Occidentali*, 55–67, *Mem. Soc. Geol. It.* 29.
- VON RAUMER (1987): Les massifs du Mont Blanc et des Aiguilles Rouges, témoins de la formation de croûte varisque dans les Alpes Occidentales. *Géologie Alpine* 63, 7–24.
- VON RAUMER (1988): Caledonian-Variscan structures in the Alps. *Schweiz. Mineral. Petrogr. Mitt.* 68, 3, 291–299.
- VON RAUMER, J.F. and SCHWANDER, H.W. (1985): Garnet evolution in pre-Variscan pelitic rocks from the Lake Emossion area, Aiguilles Rouges Massif, Western Alps. *J. Metamorphic Geol.* 3, 467–479.
- SCHLÄPFER, E. (1979): *Metamorphose und Strukturen der Cima-Lunga-Serie, IV Val Efra-Val Gagnone*. Unpubl. diploma thesis, ETH-Zurich, 54 p.
- SCHMIDT, M. (1989): Petrography, geochemistry and structural evolution of the mafic-ultramafic complex of Gana Rossa (Bellinzona zone, southern steep belt, Switzerland/Northern Italy). *Schweiz. Mineral. Petrogr. Mitt.* 69, 393–406.
- SHERVAIS, J.W. (1982): Ti–V plots and the petrogenesis of modern and ophiolitic lavas. *Earth and Planet. Sci. Lett.* 59, 101–118.
- THOMPSON, J.R., LAIRD, J. and THOMPSON, A.B. (1981): Reactions in amphibolites, greenschist and blueschist. *J. Petrol.* 23, 1–27.
- THOMPSON, R.N., MORRISON, M.A., DICKIN, A.P. and HENDRY, G.L. (1983): Continental flood basalts, Arachnids rule OK? In: HAWKESWORTH, C.J. and NORRIS, M.J., eds, *Continental basalts and mantle xenoliths*, 158–185. Shiva, Nantwich.
- UPTON, B.G.J., EMELEUS, C.H. and BECKINSALE, R.D. (1984): Petrology of the northern east Greenland Tertiary flood basalts: evidence from Hold with Hope and Wollaston Forland. *J. Petrol.* 25, 151–184.
- WAGER, L.R. (1960): The major element variation of the layered series of the Skaergaard intrusion and a re-estimation of the average composition of the hidden layered series and of the successive residual magmas. *J. Petrol.* 1, 364–398.
- WEAVER, B.L. and TARNEY, J. (1981): Chemical changes during dyke metamorphism in high-grade basement terrains. *Nature* 289, 47–49.
- WEAVER, B.L. and TARNEY, J. (1983): Chemistry of the subcontinental mantle: inferences from Archean and Proterozoic dykes and continental flood basalts. In: HAWKESWORTH, C.J. and NORRIS, M.J., eds, *Continental basalts and mantle xenoliths*, 209–229. Shiva, Nantwich.
- WILSON, A. (1960): The microdetermination of ferrous iron in silicate minerals by a volumetric and colorimetric method. *Analyst* 85, 823–827.
- WILSON, M. (1988): *Igneous petrogenesis. A global tectonic approach*. Unwin Hyman, London. 466 p.
- WOOD, D.A., JORON, J.L., TREUIL, M., NORRIS, M. and TARNEY, J. (1979a): Geochemistry of basic lavas from Iceland and the surrounding ocean floor. *Contr. Mineral. Petrol.* 70, 319–339.
- WOOD, D.A., TARNEY, J., VARET, J., SAUNDERS, A.D., BOUGOULT, H., JORON, J., TREUIL, M. and CANN, J.R. (1979b): Geochemistry of basalts drilled in the North Atlantic by IPOD 49: implications for mantle heterogeneity. *Earth Planet. Sci. Lett.* 42, 77–97.
- ZWART, H.J. and DORNSIEPEN, U.F. (1980): The Variscan and pre-Variscan tectonic evolution of Central and Western Europe, a tentative model. 26. *Intern. Geol. Congr. Paris, Colloque C6, Géologie de l'Europe*, 226–232.

Manuscript received February 2, 1990; revised manuscript accepted June 18, 1990.