Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =

Bulletin suisse de minéralogie et pétrographie

Band: 72 (1992)

Heft: 1

Artikel: Preliminary results of geochemistry, Sm-Nd and Rb-Sr studies of post-

Karoo carbonatite complexes in Southern Africa

Autor: Ziegler, U.R.F.

DOI: https://doi.org/10.5169/seals-54903

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 21.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Preliminary results of geochemistry, Sm—Nd and Rb—Sr studies of post-Karoo carbonatite complexes in Southern Africa*

by U.R.F. Ziegler 1

Abstract

The preliminary results presented in this article show that the major and trace element composition of most of the analysed post-Karoo carbonatite complexes of Namibia (Kalkfeld), Zambia (Kaluwe, Chasweta), Malawi (Kangankunde) and Tanzania (Mbeya) is in good agreement with the expected composition for the respective carbonatite types. The chondrite normalized REE patterns of all the analysed carbonatites show a strong LREE/HREE enrichment and the absence of significant anomalies. K-Ar age determinations of the Eisenberg carbonatite (Kalkfeld complex) yielded ages between 153.6 and 172.8 Ma. Rb-Sr and Sm-Nd studies indicate depleted mantle reservoirs for the Kalkfeld, Kaluwe and Kangankunde carbonatites while the Mbeya complex seems to have a Rb-depleted, Nd-enriched reservoir. The study of the Chasweta is not conclusive with respect to its mantle source as the results show a bimodal sample distribution which either indicates different mantle sources or severe contamination/mixing for the Chasweta magmas.

Keywords: Geochemistry, isotope geochemistry, carbonatites, post-Karoo complex, southern Africa.

1. Introduction

During the late Triassic the initial rifting of the South Atlantic triggered the separation of Africa from Antarctica and South America. Intracontinental stretching accompanying the opening of the South Atlantic and the South Indian Ocean led to a multitude of anorogenic magmatic events in southern Africa. During the early to late Jurassic the magmatism of the Karoo igneous province (ERLANK, 1984) was the characteristic and volumetrically most important type of magmatism in southern Africa. Old lines of weakness near the eastern end of Atlantic-related transform fault zones extending far inland (e.g. in Angola and Namibia) and intracontinental regions such as the Chilwa province in Malawi (e.g. WOOLLEY, 1991), the Rufunsa province in Zambia (Bailey, 1966; Turner, 1988) and the Mbeya area

in southern Tanzania (Van Straaten, 1989) where lineaments and/or fault systems occur and partly transsect each other were the place of subsequent late Jurassic to early Cretaceous (ca. 165 to 100 Ma) ultra-alkalic and carbonatitic magmatism in southern Africa (Kampunzu and Popoff, 1991). Within the last few years several studies on African carbonatites (e.g. Bell and Blenkinsop, 1989; Kinnaird and Bowden, 1991) have shown that the majority of African carbonatites have relatively smooth and steep REE patterns and that the source regions of the so far analysed African carbonatite complexes lie within the subcontinental mantle which must be inhomogeneous with respect to its Sr and Nd isotopic composition.

As no Rb-Sr, Sm-Nd, REE and age data exist on many of the major carbonatite complexes of southern Africa (e.g. the Kalkfeld complex in Namibia and the Rufunsa carbonatites of Zam-

^{*} Extended abstract of a contribution presented at the annual meeting of the Swiss Society of Mineralogy and Petrology, Chur, October 10 and 11, 1991.

¹ Ittigenstrasse 11, CH-3063 Ittigen.

bia) while the isotope and trace element characteristics of the carbonatites of eastern Africa and the Republic of South Africa are partly well known there exists a major gap in the knowledge on the mechanisms and sources of the post-Karoo magmatism in these areas. The aim of the present study is thus to gain more information on these subjects on a regional scale by the analysis of the REE- and isotope systematics of various post-Karoo carbonatite complexes of southern Africa. This report summarises the first preliminary results of this ongoing study.

2. Sample description

The analysed samples cover all the major structural and tectonic settings in which Mesozoic carbonatites occur in southern Africa. They were collected in the following areas:

The Rufunsa province is located at the triple junction of the NE-SW Luangwa rift with the middle and lower Zambesi rifts (Fig. 1A). This province formed close to the notheastward extension of the (reactivated) Proterozoic Mwembeshi shear zone (DALY, 1988). Carbonatites occur in a group of four complexes emplaced into a sequence of Upper Karoo sandstones (BAILEY, 1966; Turner and Rex, in press). Detailed descriptions of the Rufunsa carbonatites which are not associated with any silicic magmatic rocks may be found in Bailey (1960 and 1966), Turner (1988). K-Ar age determinations of Turner and Rex (in press) yielded ages between 101 and 113 Ma for the Kaluwe complex. The samples which were analysed during this study derive from the northernmost Kaluwe and the southernmost Chasweta complexes. The Kaluwe samples Z19, Z20 and Z21 are dark grey, fine-grained sövites. The Chasweta samples Z1, Z4, Z12 are dark brown, fine-grained ferrocarbonatites while Z17 is a brownish, fine-grained sövite.

The Kalkfeld province (Fig. 1B) of central western Namibia is situated on the southwest-northeast trending lineament of the Damaraland province (Prins, 1981) which represents an Atlantic related transform fault zone. The three carbonatite complexes of the Kalkfeld province (Kalkfeld, Ondurakorume, Osongombo) which have been described in detail by Verwoerd (1967) intruded the country rock of pan-African age amidst (on a regional scale) a large number of alkaline ring complexes. The analysed fine-grained biotite sövite samples N15 and N33 were collected on the central Eisenberg plug of the Kalkfeld complex. The large (up to 3 cm in diameter), partly chloritized biotites (N36 I – N36 IV)

used for the K-Ar age determinations of this study derive from the biotite sövites of the summit area of Eisenberg.

The Chilwa alkaline province (Fig. 1C) of southern Malawi is situated at the junction of the north-south trending East African Rift and the east-west trending Zambezi rift. Seventeen carbonatite centres of a wide compositional range are known from this area (Garson, 1965). K-Ar age determinations of Snelling (1965) and Cahen and SNELLING (1966) on the Kangankunde, Tundulu and Chilwa island complexes yielded ages of 123 ± 6 Ma, 133 ± 7 Ma and 136 ± 7 Ma, respectively. The analysed samples M5, M7 and M8 are dark brown, fine-grained, monazite rich magnesiocarbonatites from Kangankunde Hill which is (besides some alnöitic dykes and some carbonated nephelinite) free of associated silicate rocks.

The Mbeya (Panda Hill) province is located in southwestern Tanzania south of the Tanzania craton along strike of the East African Rift (Fig. 1D). K-Ar age determinations of Snelling (1965) and Miller and Brown (1963) yielded ages between 96 ± 9 Ma and 113 ± 6 Ma for the four Mesozoic carbonatites of southwestern Tanzania occurring in this area. The analysed samples EH12 and EH13 are creamy white to grey, finegrained sövites from the Panda Hill complex close to the city of Mbeya.

3. Methods

The major and REE elements of the samples N15, N33, Z1, Z4, Z12, Z17, Z19, Z20, Z21, M5 and M7 were determined at CRPG, Nancy, France using inductively coupled plasma atomic emission spectrometry while the REE composition of the samples EH12 and EH13 was determined by neutron activation at Bondar Clegg Ltd., Ottawa, Canada. CO₂ was coulometrically determined at CRPG in Nancy. All the isotopic measurements were carried out at the Laboratory for Isotope Geology of the University of Bern. The Rb-Sr concentrations and the Sr isotopic compositions were determined using the isotope dilution method as described by JAEGER (1979). The Sm and Nd concentrations as well as the Nd isotopic composition were determined according to the method described by Stoessel and Ziegler (1989) using a two column element separation procedure. The Rb analyses were carried out on an "Ion Instruments" solid source mass spectromenter. Elemental Sr, Sm, and Nd measurements were carried out on a "VG Sector" thermal ionisation mass spectrometer by single and triple filament




Fig. 1. Location map of the analysed complexes: A) Rufunsa area (Zambia) after Balley (1989); B) Kalkfeld province (Namibia) after Kinnair and Bowden (1991); C) Chilwa alkaline province (Malawi) after Woolley (1991); D) Carbonatite areas of Tanzania after Van Straaten (1989).

138

Tab. 1 Major, trace and rare earth element concentrations of the analysed samples.

Complex Kalkfeld Namibia			Chasweta Zambia			Kaluwe Zambia			Kangankunde Malawi		Mbeya Tanzania		
Sample	พ15	и33	Z1	Z4	Z12	Z 17	Z19	220	Z21	м5	M 7	EH12	EH13
Major E	Elements	(wt%):							·				
SiO2 Al2O3 Fe2O3 MnO MgO CaO Na2O K2O TiO2 P2O5 CO2	0.79 0.13 3.72 0.91 0.21 51.82 0.07 tr 2.22 37.44	25.67 9.31 9.71 0.56 5.75 21.2 0.61 3.47 2.38 1.20 13.93	1.53 0.71 11.69 1.67 8.21 31.43 0.03 tr 0.11 1.18 34.93	3.11 1.45 11.94 1.01 4.25 38.37 0.03 0.11 0.07 1.29 32.39	1.63 0.78 11.35 2.02 5.05 37.26 0.04 tr 0.06 3.00 31.76	1.51 0.44 3.69 0.51 0.48 51.61 0.03 0.03 0.11 1.35 37.21	53.75 0.06 0.03 0.02 3.45	5.74 1.58 2.15 0.51 0.88 48.76 0.06 0.89 0.08 0.86 35.27	0.0 0.4 0.2 55.7 tr tr tr 1.4	4 5.99 7 1.56 8 4.76 8 1.64 7 12.03 5 21.66 0.05 0.14 0.02 1 2.84 1 33.93	4.99 1.41 10.64 2.51 11.71 26.87 0.02 tr 0.02 1.08 35.58		
Trace Elements (ppm):													
Ba Be	434 2	>5000	>5000	>5000 10	>5000 10	>5000 5	1119 1	>5000 5	1278 2	3586 8	>5000	650	1500
Co Cr	19 13	31 362	42 27	69 26	53 23	21 18	19 12	18 18	11 8	32 17	22 21	14 46	17 23
Cu Ga	49 54	53 16	155 35	57 32	96 41	13 20	26 19	22 21	19 19	42 163	31 79		
Nb Ni	189 16	170 86	>1000 19	460 157	609 40	546 15	962 14	379 15	362 10	170 25	23 13		
Rb Sc Sr	1 26 8064	66 32 5844	<1 24 1830	6 26 1663	<1 42 1737	4 35 723	2 16 5272	22 17 992	1 20 748	2 39 90258	1 28 27260	6 5 13278	<1 4 3910
Th V	252 25	301 148	216 151	329 369	433 201	109 159	71 75	99 168	55 104	822 43	326 49	4	214
Y Zn Zr	333 86 37	86 127 275	141 510 105	266 395 375	197 202 123	90 142 104	62 26 53	96 168 76	74 45 32	80 >1000 153	46 >1000 66		
Rare earth elements (ppm):													
La Ce Nd Sm Eu Gd Dy Er	2967 5193 2897 335 75.8 212 113 47.6	183 285 180 35.7 9.7 23.7 13.2	38.2	1825 2945 1267 161 41.7 127 77.5 40.3	2892 4195 1507 177 46.9 146 85.0 38.8	513 876 374 57.1 19.1 63.3 43.0	243 496 239 36.8 8.31 23.6 11.3 4.71	60.8 45.8	402 724 294 43.9 10.0 28.3 15.0	90 318 111	5625 9012 3129 216 53 144 65 58	612 1120 391 46.9 10	306 610 222 31.9 7
Yb Lu	23.9 3.9	6.1 1.1	17.0 2.9	29.7 5.5	26.7 5.0	15.9 3.0	1.93 0.33	17.2	2.8	23	20 4	9 0.9	7 <0.2

modes, respectively. The ⁸⁷Sr/⁸⁶Sr ratios are normalized to ⁸⁸Sr/⁸⁶Sr = 0.1194 while the ¹⁴³Nd/¹⁴⁴Nd ratios are normalized to a ¹⁴⁶Nd/¹⁴⁴Nd ratio of 0.72190. Total blanks were < 1 ng for Sm, 0.01–0.1 ng for Nd and < 10 ng for Sr. K was analysed by flame photometry from an aliquot of the sample. The isotopic composition and the concentration of Ar were determined on a VG MM1200 solid source mass spectrometer following the extraction and purification in a Pyrex glass line and the addition of 99.997% pure ³⁸Ar spike.

4. Results and discussion

The results of the major, trace and rare earth element analyses are listed in table 1 while those

of the isotopic analyses are compiled in tables 2-4.

The K-Ar ages of the analysed Eisenberg samples range between 153.6 and 172.8 Ma and thus confirm the classification of the Kalkfeld carbonatites as post-Karoo complexes.

Figure 2 shows a general classification of the analysed samples according to the propositions of Woolley and Kempe (1988). A comparison with the data in Woolley and Kempe (1988) shows, with the exception of N33, that the major and trace element geochemistry of the samples is (although certain trace elements are in cases slightly enriched compared to the literature data) generally in good agreement with the expected range for the respective carbonatite types. The abnormal composition of N33 is due to the presence of biotite, chlorite, vermiculite, augite and some

Tab. 2 Tabulation of the Rb and Sr results. Constants used are: 87 Rb decay constant = $1.42 \cdot ^{-11}a^{-1}$ (Steiger and Jaeger, 1977), 85 Rb/ 87 Rb = 2.59265, 86 Sr/ 88 Sr = 0.1194, 84 Sr/ 86 Sr = 0.056584 (Steiger and Jaeger, 1977), 87 Sr/ 86 Sr $_{\text{Uniform Reservoir UR}}$ = 0.7045 (e.g. De Paolo, 1988), 87 Rb/ 86 Sr UR = 0.0827 (e.g. De Paolo, 1988).

Sample	Rb ppm	Sr ppm	87Sr/86Sr	+/-	87Rb/86Sr	+/-	Epsilon Sr(T)
Kalkfeld Namibia							
N15 N33	0.57 65.53	8064 5844	0.703669 0.703918	10 12	0.000206 0.032432	0.000002 0.000324	-9.1 -6.6
Chasweta Zambia						, 	
Z1 Z4 Z12 Z17	0.32 6.19 0.41 3.68	1830 1663 1737 723	0.707739 0.705736 0.706338 0.703836	9 7 11 8	0.000498 0.010769 0.000683 0.014702	0.000005 0.000108 0.000007 0.000147	47.9 19.2 28.0 -7.9
Kaluwe Zambia				*			
Z19 Z20 Z21	2.18 22.89 1.08	5272 992 748	0.703286 0.704240 0.703747	9 11 8	0.001195 0.066747 0.004169	0.000012 0.000667 0.000042	-15.4 -3.3 -8.9
Kangankun Malawi	de					n 1	,
M5 M7 M8	1.82 0.81 0.59	90258 27260 39367	0.703016 0.703067 0.703033	5 7 9	0.000059 0.000086 0.000043	0.000001 0.000001 0.000001	-19.0 -18.3 -18.7
Mbeya Tanzania							
EH12 EH13	5.69 0.61	13279 3910	0.703876 0.703407	7 8	0.001238 0.000449	0.000012 0.000004	-7.2 -13.9

K-feldspar besides carbonates and limonite. Figure 3 shows the results of the REE analyses in chondrite normalized abundance patterns. All the REE abundance patterns are very steep and smooth due to very high light REE abundances and the absence of pronounced anomalies both of which are typical features of carbonatites. The $\varepsilon Nd_{(T)}$ values calculated for the respective intrusion ages of the analysed complexes from the measured ¹⁴³Nd/¹⁴⁴Nd data range between –5.3 and +5.5. The Kangankunde, Kaluwe and Eisenberg samples all have positive $\varepsilon Nd_{(T)}$ values while those of the Mbeya and Chasweta samples are at least partially negative. With the exception of Chasweta all the $\varepsilon Sr_{(T)}$ values range between –19.0 and –3.3. The $\varepsilon Sr_{(T)}$ values of the Chasweta samples are between –7.9 and +47.9.

Figure 4 shows the results of the isotopic measurements in an $\varepsilon Nd_{(T)}$ v. $\varepsilon Sr_{(T)}$ diagram. All the Kaluwe, Eisenberg and Kangankunde samples as well as one Chasweta data point scatter within the upper left (depleted) quadrant. This indicates that




Fig. 2 CaO-MgO-Total Iron Oxide + MnO plot of the analysed carbonatites (same symbols as in Fig. 4).

Tab. 3 Tabulation of the Sm and Nd results. Constants used are: 147 Sm decay constant = 0.00654 b.y. - (Lugmair and Marti, 1978), 143 Nd/ 144 Nd_{Chondritic Uniform Reservoir CHUR (t = 0)} = 0.51264, 143 Nd/ 144 Nd_{Depleted Mantle DM(0)} = 0.51316, 147 Sm/ 144 Nd_{CHUR} = 0.1967, 147 Sm/ 144 Nd DM = 0.2136.

Sample	Sm ppm	Nd ppm	143Nd/144Nd	+/-	147Sm/144	Nd +/-	Epsilon Nd(T)
Kalkfeld Namibia					it.		ė.
N15 N33	334.65 35.74	2897.8 180.06	0.512575 0.512580	2 3	0.0698 0.1200	0.0007 0.0012	-1.3 0.4
Chasweta Zambia							
Z1 Z4	177.00	1582.1 1267.0	0.512442	2	0.0676	0.0007	-2.1 -2.8
Z12 Z17	176.95 57.12	1506.9 374.48	0.512280 0.512849	5 16	0.0710 0.0922	0.0007 0.0009	-5.3 -5.5
Kaluwe Zambia							
Z19 Z20 Z21	36.83 63.07 43.95	238.88 439.67 293.51	0.512737 0.512694 0.512750	18 3 3	0.0932 0.0867 0.0905	0.0009 0.0009 0.0009	3.3 2.6 3.6
Kangankun Malawi	de						
M5 M7 M8	477.18 216.33 189.05	8129.6 3129.3 2950.3	0.512679 0.512694 0.512669	2 3 2	0.0355 0.0418 0.0387	0.0004 0.0004 0.0004	3.3 3.5 3.0
Mbeya Tanzania		e					
EH12 EH13	46.88 31.85	390.76 222.00	0.512511 0.512506	3 5	0.072523 0.086734	0.0007 0.0009	-1.0 -1.3

Tab. 4 Tabulation of the K–Ar results. Constants used are: ${}^{40}\text{K} = 5.543 \cdot 10^{-10} \text{a}^{-1}$ (total ${}^{40}\text{K}$ decay constant), ${}^{40}\text{K}^{\text{e}} = 0.581 \cdot 10^{-10} \text{a}^{-1}$ (Steiger and Jaeger, 1977), ${}^{40}\text{Ar}/{}^{36}\text{Ar}_{\text{atm}} = 295.5, {}^{38}\text{Ar}/{}^{36}\text{Ar}_{\text{atm}} = 0.1869$ (Steiger and Jaeger, 1977).

Sample		K(wt%)	40Ar/36Ar	40K/36Ar	Age	(Ma
N36	I	1.41	1243.8	94898	164.	3
N36	II	2.44	1069.5	78984	161.	3
N36	III	3.37	3620.4	315562	172.	8
N36	IV	2.58	6144.8	627735	153.	6

these carbonatites derive from reservoirs depleted in Large Ion Lithophile Elements (LILE) relative to bulk Earth while the Mbeya samples seem to derive from a mantle source with lower Sm/Nd and Rb/Sr ratios relative to the bulk Earth. Three of the four Chasweta samples plot within the enriched quadrant. A comparison with literature data taken from a compilation of Bell and Blenkinsop (1989) shows that the majority of the samples plots in the areas of known carbonatite compositions although some of them do not follow the East African carbonatite line of Bell

and BLENKINSOP (1989). The high 87Sr/86Sr and low ¹⁴³Nd/¹⁴⁴Nd ratios of the three enriched Chasweta samples are somewhat puzzling as they plot far away from known carbonatite compositions. The fact that the fourth Chasweta sample (Z17) as well as all the Kaluwe samples do not show such aberrant isotopic compositions raises even more questions on the genesis of these Rufunsa carbonatites. The only conclusions that can be drawn from the present data are the following: The Kaluwe samples as well as Z17 seem to have their origins in a mantle reservoir depleted in LILE and their isotopic composition remained more or less unchanged during formation, ascent and emplacement of the carbonatite magmas. If the sövitic Z17 and the three enriched ferrocarbonatitic Chasweta samples which show a certain tendency towards Group II kimberlites (data from Gurney et al., 1991) do not have the same origins then the aberrant samples must either derive from a comparatively enriched mantle source or they must have had a strong crustal component in their




Fig. 3 Chondrite normalized REE plot for the analysed carbonatites.




Fig. 4 Plot of $\varepsilon Nd_{(T)}$ versus $\varepsilon Sr_{(\Gamma)}$ for the analysed samples. Data for comparison are taken from the compilation of Bell and Blenkinsop (1989).

source area while Z17 derives from a depleted source. If, however, they have common origins then certain parts of the Chasweta magma must have been subject to mixing processes including the assimilation of continental crust material. The first interpretation is supported up to a certain degree by the considerations of Bailey (1989) who identified magnesio-chromite-bearing pyroclasts in part of the Chasweta carbonatites which thus indicate a certain kimberlite affinity of these pyroclastic carbonatites. As magnesio-chromite is only present in the pyroclastic carbonatites and is absent in the coarse grained intrusive carbonatites Bailey (1989) also stated that the Rufunsa carbonatites must derive from various origins; a conclusion which would also support the first interpretation. However, only further detailed field and laboratory work can help to find a solution to this Rufunsa problem.

Acknowledgements

The author would like to thank Prof. E. Jäger for making this study possible. The geological surveys of Namibia, Zambia and Malawi are thanked for their continuous support during my visits to the respective countries. I am specially indebted to K. Lyungu who introduced me to the Rufunsa carbonatites. The study was partly financed by the "Schweizerischer Nationalfonds".

References

- BAILEY, D.K. (1989): Carbonate melt from the mantle in the volcanoes of south-east Zambia. Nature, 338/ 6214, 415-418.
- Bailey, D.K. (1966): Carbonatite volcanoes and shallow intrusions in Zambia. In: TUTTLE, O.F. and GITTINS, J. (editors): Carbonatites. Interscience, New York,
- BAILEY, D.K. (1960): Carbonatites of the Rufunsa valley, Feira District. Bulletin of the geological Survey of Northern Rhodesia, 5.
- Bell, K. and Blenkinsop, J. (1989): Neodymium and strontium isotope geochemistry of carbonatites. In: Bell, K. (editor): Carbonatites, Genesis and Evolution. Unwin Hyman, London, 278-300.
- CAHEN, L. and SNELLING, N.J. (1966): The geochronology of equatorial Africa. North-Holland, Amsterdam, 195 p.

 DALY, M.C. (1988): Crustal shear zones in Central Afri-
- ca: a kinematić approach. Episodes 11/1, 5–11. DE PAOLO, D.J. (1988): Neodymium Isotope Geochemistry. An Introduction. Springer, Berlin, 187 p.
- ERLANK, A.J. (1984): Petrogenesis of the volcanic rocks of the Karoo province. Geol. Soc. South Afr. Spec.
- Publ. 13, 395 p.
 Garson, M.S. (1965): Carbonatites in southern Malawi. Bull. geol. Surv. Malawi, 15, 128 p.
- GURNEY, J.J., MOORE, R.O., OTTER, M.L., KIRKLEY, M.B., HOPS, J.J. and McCandless, T.E. (1991):

- Southern African kimberlites and their xenoliths. In: KAMPUNZU, A.B. and LUBALA, R.T. (editors): Magmatism in Extensional Structural Settings, The Phanerozoic African Plate. Springer, Berlin, 495-
- JAEGER, E. (1979): The Rb-Sr method, 13-26. In: JAEGER, E. and HUNZIKER, J.C. (editors): Lectures in Isotope Geology. Springer, Berlin, 329 p.
- KAMPUNZU, A.B. and POPOFF, M. (1991): Distribution of the Main Phanerozoic African Rifts and Associated Magmatism: Introductory Notes. In: KAMPUNZU, A.B. and LUBALA, R.T. (editors): Magmatism in Extensional Structural Settings, The Phanerozoic African Plate. Springer, Berlin, 2-10.
- KINNAIRD, J.A. and BOWDEN, P. (1991): Magmatism and Mineralization Associated with Phanerozoic Anorogenic Plutonic Complexes of the African Plate. In: Kampunzu, A.B. and Lubala, R.T. (editors): Magmatism in Extensional Structural Settings, The Phanerozoic African Plate. Springer, Berlin,
- LUGMAIR, G.W. and MARTI, K. (1978): Lunar initial ¹⁴³Nd/¹⁴⁴Nd; differential evolution of the lunar crust and mantle. Earth Planet. Sci. Lett. 39, 349-357.
- MILLER, J.A. and Brown, P.E. (1963): The age of some carbonatite activity in south-west Tanganyika. Geological Magazine 100, 276-9.
- Prins, P. (1981): The geochemical evolution of the alkaline and carbonatite complexes of the Damaraland igneous province, South West Africa. Ann. Univ. Stellenbosch Ser A1 (Geol.) 3, 145-278.
- Snelling, N.J. (1965): Age determination on three African carbonatites. Nature 205, 491
- STEIGER, R.H. and JAEGER, E. (1977): Subcommission on geochronology: convention on the use of decay constants in geo- and cosmochronology. Earth.
- Planet. Sci. Lett., 36, 358–362. Stoessel, G.F.U. and Ziegler, U.R.F. (1989): Age determinations in the Rehoboth Basement Inlier, SWA/Namibia. Unpubl. doctoral thesis. University of Bern, Switzerland, 250 p.
- TURNER, D.C. (1988): Volcanic carbonatites of the Kaluwe complex, Zambia. J. Geol. Soc. London, 145,
- TURNER, D.C. and REX, D.C. (in press): Volcaniclastic carbonatite at Kaluwe, Zambia: Age and relations
- to sedimentary rocks in the Zambezi rift valley.

 Van Straaten, P. (1989): Nature and structural relationships of carbonatites from Southwest and West Tanzania. In: Bell, K. (editor): Carbonatites, Genesis and Evolution. Unwin Hyman, London, 177-
- Verwoerd, W.J. (1967): The carbonatites of South Africa and South West Africa. Geol. Surv. South Afr. Handb., 6, 452 p. OOLLEY, A.R. (1991): The Chilwa Alkaline Igneous
- Province of Malawi: A Review. In: KAMPUNZU, A.B. and LUBALA, R.T. (editors): Magmatism in Extensional Structural Settings, The Phanerozoic African Plate. Springer, Berlin, 637 p.
- Woolley, A.R. and Kempe, D.R.C. (1989): Carbonatites: Nomenclature, average chemical compositions, and element distribution. In: Bell, K. (editor): Carbonatites, Genesis and Evolution. Unwin, Hyman, London, 1–14.