

<b>Zeitschrift:</b>	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
<b>Band:</b>	72 (1992)
<b>Heft:</b>	3
<b>Rubrik:</b>	New minerals recently approved by the Commission on New Minerals and Mineral Names, International Mineralogical Association

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 31.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**New minerals recently approved  
by the  
Commission on New Minerals and Mineral Names  
International Mineralogical Association**

The information given here is provided by the Commission on New Minerals and Mineral Names, I.M.A. for comparative purposes and as a service to mineralogists working on new species.

Each mineral is described in the following format:

IMA No.

(any relationship to other minerals)

Chemical Formula

Crystal system, space group

unit cell parameters

Colour; lustre; diaphaneity.

Optical properties

Strongest lines in the X-ray powder diffraction pattern

The names of these approved species are considered confidential information until the authors have published their descriptions or released information themselves.

**No other information will be released by the commission**

J. A. Mandarino, Chairman  
Commission on New Minerals and Mineral Names  
International Mineralogical Association

**1991 Proposals**

IMA No. 91-001

$Hg_2^{+1}Hg_3^{+2}Cr^{+6}O_5S_2^{-2}$

Triclinic:  $P\bar{1}$

a 8.116, b 9.501, c 6.891 Å,  $\alpha$  100.43°,  
 $\beta$  110.24°,  $\gamma$  82.80°

Orange-red; adamantine; transparent.

Biaxial (sign unknown), all indices of refraction are greater than 2.

5.72 (90), 3.373 (60), 3.008 (100), 2.864 (50b),  
2.774 (50), 2.536 (50), 2.486 (50), 2.425 (60).

IMA No. 91-003

The niobium analogue of bismutotantalite.

$Bi(Nb,Ta)O_4$

Orthorhombic: Pcmn

a 4.992, b 5.677, c 11.731 Å

Black; semi-metallic; transparent in small ( $< 0.03$  mm) fragments.

Biaxial (+),  $\alpha$  2.38,  $\beta$  2.42,  $\gamma$  2.47,  $2V$ (calc.) 85°.

3.164 (100), 2.934 (90), 2.842 (45), 2.495 (45),  
1.769 (45), 1.734 (80).

IMA No. 91-005

$(Zn,Co,Ni)_6(SO_4)_4(OH,Cl)_{10} \cdot 5 H_2O$

Hexagonal:  $P6_3$ ,  $P6_3/m$  or  $P6_322$   
a 8.344, c 21.59 Å

Bright to deep pink; vitreous to pearly;  
transparent.

Uniaxial (-),  $\omega$  1.584,  $\epsilon$  1.544

10.8 (100), 3.300 (90), 2.725 (60), 2.563 (50),  
2.351 (40), 1.575 (30).

IMA No. 91-007

$Mn_5(OH)_4(VO_4)_2$

Monoclinic: C2/m

a 9.604, b 9.558, c 5.393 Å,  $\beta$  98.45°

Orange-red; vitreous; transparent.  
Biaxial (-),  $\alpha'$  1.803,  $\gamma'$  1.810, 2V(meas.) large.  
4.76 (S), 3.00 (M), 2.680 (VS), 2.656 (M),  
2.155 (M), 1.565 (M), 1.510 (M).

## IMA No. 91-008

The Ba-dominant end-member of the alunite group.  
 $\text{Ba}_{0.5}\square_{0.5}\text{Al}_3(\text{SO}_4)_2(\text{OH})_6$   
Hexagonal:  $\text{R}\bar{3}\text{m}$   
a 6.992, c 17.22 Å

White to light yellowish; vitreous; transparent.  
Uniaxial (+),  $\omega$  1.588,  $\epsilon$  1.604.  
5.73 (50), 3.49 (55), 2.98 (100), 2.283 (80),  
1.909 (70), 1.747 (60).

## IMA No. 91-009

The Ca-dominant end-member of the alunite group.  
 $\text{Ca}_{0.5}\square_{0.5}\text{Al}_3(\text{SO}_4)_2(\text{OH})_6$   
Hexagonal:  $\text{R}\bar{3}\text{m}$   
a 6.983, c 16.759 Å

White to light yellowish; vitreous; transparent.  
Uniaxial (+), indices of refraction unknown.  
4.91 (69), 2.97 (100), 2.231 (51), 1.899 (43),  
1.745 (37), 1.375 (40).

## IMA No. 91-010

$\text{Ca}(\text{Fe},\text{Mg})_6(\text{SiO}_4)_2(\text{PO}_4)_2$   
Hexagonal:  $\text{R}\bar{3}\text{m}$   
a 6.240, c 26.784 Å  
Yellow-brown; vitreous; transparent.  
Uniaxial (-),  $\omega$  1.770,  $\epsilon$  1.759.  
5.00 (60), 3.119 (100), 2.689 (80), 2.558 (100),  
2.505 (80), 1.560 (80).

## IMA No. 91-012

$\text{Mn}_5\text{SnB}_2\text{Si}_5\text{O}_{20}$   
Monoclinic: P2/m  
a 28.77, b 7.01, c 13.72(2) Å,  $\beta$  96.6(2)°  
Orange-yellow; vitreous; transparent.  
Biaxial (-),  $\alpha$  1.696,  $\beta$  1.711,  $\gamma$  1.715,  
2V(meas.) 57°, 2V(calc.) 54°.  
3.41 (8), 3.22 (8), 2.83 (10), 2.81 (10), 2.24 (7),  
1.750 (6).

## IMA No. 91-013

$(\text{Na},\text{K})_{17}\text{Fe}^{+3}\text{TiSi}_{16}\text{O}_{29}(\text{OH})_{30} \cdot 2 \text{H}_2\text{O}$   
Orthorhombic: Cmc<sub>1</sub>, Cmc<sub>2</sub><sub>1</sub> or C2cm  
a 29.77, b 11.03, c 17.111(5) Å  
Colourless (white or grey in aggregates);  
vitreous; transparent.  
Biaxial (-),  $\alpha$  1.532,  $\beta$  1.548,  $\gamma$  1.559(2),  
2V(meas.) 79°, 2V(calc.) 79°.  
10.38 (100), 4.516 (75), 3.220 (65), 3.097 (80),  
2.972 (65), 2.773 (90).

## IMA No. 91-014

$\text{Na}_4\text{K}_3(\text{Fe},\text{Mn},\text{Ti})_2\text{Si}_8\text{O}_{20}(\text{OH})_4 \cdot 4 \text{H}_2\text{O}$   
Triclinic: P  $\bar{1}$   
a 10.244, b 11.924, c 5.276 Å,  $\alpha$  103.491°,  
 $\beta$  96.960°,  $\gamma$  91.945°

Olive-green with brownish or yellowish shades;  
vitreous; transparent.  
Biaxial (+),  $\alpha$  1.569,  $\beta$  1.574,  $\gamma$  1.590,  
2V(meas.) 58°, 2V(calc.) 59°.  
11.57 (100), 3.386 (19), 3.006 (21), 2.992 (28),  
2.716 (22), 2.598 (26).

## IMA No. 91-015

$\text{Na}_8\text{KSi}_9\text{O}_{18}(\text{OH})_9 \cdot 19 \text{H}_2\text{O}$   
Monoclinic: P2<sub>1</sub>/c  
a 24.91, b 11.94, c 14.92 Å,  $\beta$  94.47(9)°  
Colourless; vitreous; transparent.

Biaxial (-),  $\alpha$  1.460,  $\beta$  1.478,  $\gamma$  1.481,  
2V(meas.) 43°, 2V(calc.) 44°.  
4.26 (60), 3.08 (100), 2.938 (70B), 2.649 (60B),  
2.400 (35), 2.289 (35).

## IMA No. 91-016

A member of the adelite-descloizite group.  
 $\text{CaMn}(\text{OH})\text{SiO}_4$   
Orthorhombic: P2<sub>1</sub>2<sub>1</sub>2<sub>1</sub>  
a 5.838, b 7.224, c 8.690(1) Å  
Deep red; vitreous; transparent.  
Biaxial (+),  $\alpha$  1.840,  $\beta$  (calc.) 1.854,  $\gamma$  1.920,  
2V(meas.) 50°.  
5.558 (S), 3.070 (S), 2.687 (S), 2.584 (VS),  
1.565 (M).

## IMA No. 91-017

The ferric-analogue of philipsbornite.  
 $\text{PbFe}_3^{+3}\text{H}(\text{AsO}_4)_2(\text{OH})_6$   
Hexagonal:  $\text{R}\bar{3}\text{m}$   
a 7.359, c 17.113(8) Å  
Greenish-yellow; vitreous to adamantine;  
translucent to transparent.  
Uniaxial (-),  $\omega$  1.975,  $\epsilon$  1.955.  
5.966 (50), 3.678 (40), 3.092 (100), 2.283 (30),  
1.992 (30), 1.840 (25).

## IMA No. 91-018

The Mg-dominant analogue of concolite and  
the rhombohedral dimorph of boracite.  
 $(\text{Mg},\text{Fe})_3\text{B}_7\text{O}_{13}\text{Cl}$   
Hexagonal: R3c  
a 8.574, c 20.99 Å  
Colourless; vitreous; transparent.  
Uniaxial (-),  $\omega$  1.684,  $\epsilon$  1.668.  
3.497 (34), 3.028 (100), 2.711 (66), 2.144 (37),  
2.050 (73), 1.828 (25).

## IMA No. 91-019

$\text{Fe}_3^{+2}(\text{Fe}^{+3},\text{Ti})_3[\text{O}_2\text{As}_2\text{O}_5]_2$

**Monoclinic: P<sub>2</sub>/m**

a 10.625, b 3.264, c 8.990 Å, β 109.15°.  
 Dark brown to black; submetallic to metallic;  
 opaque (translucent in thin fragments).  
 In reflected light: creamy white (in oil, white  
 with a weak brown tint); no internal  
 reflections; anisotropy visible along grain  
 boundaries (in oil, clearly visible); bireflectance  
 not visible (in oil, very weak  
 along grain boundaries); nonpleochroic.  
 R-values: (15.5–15.9%) 470 nm, (15.0–  
 15.5%) 546 nm, (14.8–15.0%) 589 nm,  
 (14.2–14.5%) 650 nm.  
 2.985 (67), 2.811 (94), 2.749 (100), 2.391 (85),  
 1.779 (48), 1.709 (35).

**IMA No. 91-020**

$\text{Ba}(\text{Al,Mg})(\text{PO}_4,\text{CO}_3)(\text{OH})_2 \cdot \text{H}_2\text{O}$   
 Orthorhombic: Pnna or Pnnn  
 a 8.939, b 5.669, c 11.073(3) Å  
 Pale blue; silky; translucent.  
 Biaxial (−), α 1.616, β 1.629, γ 1.640,  
 2V(meas.) 70–90°, 2V(calc.) 85°.  
 5.54 (79), 3.479 (82), 3.345 (59), 2.768 (100),  
 2.543 (61), 2.072 (41).

**IMA No. 91-021**

A polymorph of mundrillaite.  
 $\text{Ca}(\text{NH}_4)_2(\text{HPO}_4)_2 \cdot \text{H}_2\text{O}$   
 Orthorhombic: space group unknown  
 a 20.959, b 7.403, c 6.478(5) Å  
 White; vitreous; transparent.  
 Biaxial (−), α 1.506, β 1.510, γ 1.512,  
 2V(meas.) 65°, 2V(calc.) 70°.  
 10.5 (57), 6.99 (100), 4.739 (36), 3.705 (89),  
 3.651 (39), 3.177 (55).

**IMA No. 91-022**

$\text{KZr}_2(\text{PO}_4)_3$   
 Hexagonal: R̄3c  
 a 8.687, c 23.877(7) Å  
 Pale blue to blue-green to nearly colourless;  
 vitreous; transparent.  
 Uniaxial (+), ω 1.656, ε 1.682.  
 6.41 (50), 4.679 (50), 4.329 (100), 3.806 (90),  
 2.928 (90), 2.502 (50).

**IMA No. 91-023**

The Cs-analogue of gainesite.  
 $\text{NaCs}(\text{Be,Li})\text{Zr}_2(\text{PO}_4)_4$   
 Tetragonal: I4<sub>1</sub>/amd  
 a 6.573, c 17.28 Å  
 White to colourless; vitreous; translucent to  
 transparent.  
 Uniaxial (+), ω 1.634, ε 1.645.  
 6.159 (90), 4.326 (80), 4.099 (40), 3.281 (80),  
 3.060 (100), 2.896 (30), 1.849 (30).

**IMA No. 91-024**

$\text{Sb}_6(\text{SO}_4)\text{O}_8 \cdot \text{H}_2\text{O}$   
 Triclinic: P1  
 a 11.434, b 29.77, c 11.314(4) Å, α 91.07°,  
 β 119.24°, γ 92.82°  
 Colourless to white; adamantine; transparent  
 to translucent.  
 Biaxial (+), mean n 2.08, birefringence low,  
 2V(meas.) >> 60°.  
 14.835 (50), 9.270 (41), 6.810 (67), 3.304 (93),  
 3.200 (39), 3.092 (100).

**IMA No. 91-025**

$\text{Cu}_4(\text{As,Sb})_2\text{S}_5$

Orthorhombic: space group unknown  
 a 14.51, b 13.30, c 17.96(1) Å

Silvery lead grey; metallic; opaque.

In reflected light: grey, weak anisotropism,  
 weak bireflectance, nonpleochroic. R<sub>min.</sub>  
 and R<sub>max.</sub>: (31.5, 32.5%) 470 nm, (31.1,  
 32.0%) 546 nm, (30.3, 31.15%) 589 nm,  
 (27.2, 23.4%) 650 nm.  
 3.36 (7), 2.999 (100), 2.594 (20), 2.238 (6),  
 1.833 (40), 1.564 (15b).

**IMA No. 91-026**

$(\text{Co,Zn,Ni})_6(\text{SO}_4)(\text{OH,Cl})_{10} \cdot 8 \text{ H}_2\text{O}$

Hexagonal: space group unknown  
 a 8.363, c 26.18(7) Å

Pink to light pink; pearly; transparent.

Uniaxial (−), ω 1.568, ε 1.542.  
 13.1 (100), 3.523 (30), 2.985 (30), 2.681 (40),  
 2.527 (90).

**IMA No. 91-027**

The fluorine-analogue of britholite-(Ce) of the  
 apatite group.

(REE, Ca)<sub>5</sub>(Si,P)<sub>3</sub>O<sub>12</sub>F, where Ce is the  
 dominant REE.

Hexagonal: P6<sub>3</sub>m  
 a 9.517, c 6.983(4) Å

Tan, reddish-brown; adamantine; opaque to  
 translucent.

Uniaxial (−), ω 1.792, ε 1.786.  
 2.845 (100), 2.822 (40), 2.747 (30), 1.970 (30),  
 1.870 (40).

**IMA No. 91-028**

A member of the amphibole group.

$(\text{Na,K})\text{Na}_2[\text{Mg}_2(\text{Fe}^{3+},\text{Mn}^{3+})_2\text{Li}]\text{Si}_8\text{O}_{22}(\text{OH,F})_2$   
 Monoclinic: C2/m

a 9.808, b 17.850, c 5.289(1) Å, β 104.22(2)°

Dark red; vitreous; translucent.

Biaxial (+), α 1.667, β 1.675, γ 1.691,  
 2V(meas.) 59–71°, 2V(calc.) 71°.  
 8.399 (56), 3.383 (18), 3.254 (20), 3.122 (100),  
 2.798 (48), 2.696 (15).

## IMA No. 91-029

$\text{NiSb}_2(\text{OH})_{12} \cdot 6 \text{H}_2\text{O}$   
 Hexagonal: P31m, P31m or P312  
 a 16.016, c 9.789(2) Å  
 Light-blue; vitreous; transparent.  
 Uniaxial (+),  $\omega$  1.600,  $\epsilon$  1.605.  
 4.6195 (100), 3.3537 (100), 2.3431 (80),  
 2.0909 (60), 1.8050 (70), 1.7496 (60).

## IMA No. 91-030

$(\text{Pb},\text{Mo},\square)_8\text{O}_8\text{Cl}_2$   
 Tetragonal: I4/mmm, I4̄2m, I4m2, I4mm  
 or I4̄22  
 a 3.9922, c 22.514(2) Å  
 Carmine; adamantine; translucent.  
 In reflected light: grey, weak to moderate  
 anisotropy, moderate bireflectance, weak  
 pleochroism, internal reflections abundant,  
 $R_1$  and  $R_2$  (19.6, 22.0%) 470 nm, (18.0,  
 20.5%) 546 nm, (17.4, 19.6%) 589 nm,  
 (16.95, 18.8%) 650 nm.  
 3.507 (32), 2.983 (100), 2.816 (78), 1.989 (75),  
 1.658 (51), 1.586 (33).

## IMA No. 91-031

$\text{Ca}_6\text{Cu}_3(\text{SO}_4)_3(\text{OH})_{12} \cdot 2 \text{H}_2\text{O}$   
 Monoclinic: P2<sub>1</sub>/c (pseudo C2/c)  
 a 15.122, b 14.358, c 22.063 Å,  $\beta$  108.68°.  
 Dark blue; vitreous; transparent.  
 Biaxial (-),  $\alpha$  1.590,  $\beta$  1.610,  $\gamma$  1.619,  
 2V(meas.) 65°, 2V(calc.) 67°.  
 3.393 (100), 3.368 (55), 3.200 (53), 3.188 (65),  
 3.120 (85), 3.098 (57).

## IMA No. 91-032

$\text{Fe(OH)}_3$   
 Orthorhombic Immm (pseudocubic)  
 a 7.544, b 7.558, c 7.560(4) Å  
 Dark bottle green; vitreous to adamantine;  
 transparent.  
 Biaxial (-), the indices of refraction are  
 between 1.92 and 1.94.  
 3.774 (100), 2.671 (35), 2.395 (30), 1.904 (15),  
 1.697 (60), 1.548 (40).

## IMA No. 91-033

$\text{Au}_2\text{Pb}$   
 Cubic: Fd3m  
 a 7.933(5) Å  
 Colour unknown because of the small grain  
 size; metallic; opaque.  
 In reflected light: silvery grey, dark grey when  
 highly oxidized; no anisotropy, bireflect-  
 ance, pleochroism or internal reflections;  
 R (56.0%) 470 nm, (59.5%) 546 nm,  
 (60.0%) 589 nm, (62.0%) 650 nm.  
 4.595 (21), 2.810 (30), 2.391 (100), 2.301 (25),  
 1.526 (23), 1.196 (26).

## IMA No. 91-034

$\text{Ca}(\text{UO}_2)_3(\text{CO}_3)_4 \cdot 3 \text{H}_2\text{O}$   
 Orthorhombic: Pmn<sub>1</sub>, Pmn<sub>2</sub>, or P2<sub>1</sub>nm  
 a 15.337, b 17.051, c 6.931 Å  
 Canary yellow; vitreous; transparent.  
 Biaxial (-),  $\alpha$  1.603(calc.),  $\beta$  1.690,  $\gamma$  1.710,  
 2V(meas.) 49°.  
 8.55 (100), 6.94 (50), 4.11 (60), 3.723 (60),  
 3.460 (50), 2.772 (70).

## IMA No. 91-037

$[\text{Ag}_5(\text{Pb},\text{Fe})\text{Bi}_7]_{13}(\text{Sb},\text{Bi})_2\text{S}_{17}$   
 Monoclinic: C2/m or Cm  
 a 13.515, b 4.098, c 26.000 Å,  $\beta$  93.00°  
 Grey; metallic; opaque.  
 In reflected light: white, distinct anisotropy,  
 very weak bireflectance, no pleochroism,  
 no internal reflections,  $R_{\max}$  and  $R_{\min}$ ,  
 (42.2, 39.7%) 470 nm, (41.4, 38.8%) 546 nm,  
 (40.8, 37.9%) 589 nm, (39.8, 36.9%) 650 nm.  
 3.49 (8), 3.37 (9), 3.24 (9), 2.82 (10), 2.01 (7),  
 1.992 (8), 1.967 (6).

## IMA No. 91-038

$\text{Pb}_2(\text{Mn},\text{Fe},\text{Mg})_3\text{Fe}^{3+}_{14}\text{O}_{26}$   
 Hexagonal: P6<sub>3</sub>/mmc, P6<sub>3</sub>mc or P6̄2c  
 a 5.951, c 33.358 Å  
 Black; submetallic; opaque.  
 In reflected light: grey with pale brownish tint,  
 moderate anisotropy, weak bireflectance,  
 no pleochroism, no internal reflections,  $R_O$   
 and  $R_E$  (23.6, 22.3%) 470 nm, (22.8, 21.9%)  
 546 nm, (22.2, 21.5%) 589 nm, (21.3, 21.0%)  
 650 nm.  
 4.168 (55), 3.011 (60), 2.9750 (70), 2.8017 (95),  
 2.6236 (100), 2.6125 (90).

## IMA No. 91-042

$\text{N}(\text{CH}_3)_4[\text{Si}_2(\text{Si}_{0.5}\text{Al}_{0.5})\text{O}_6]_2$   
 Orthorhombic: I222  
 a 8.984, b 8.937, c 8.927 Å  
 White, colourless, light yellow; vitreous;  
 transparent.  
 Biaxial (-),  $\alpha$  1.529,  $\beta$  (calc.) 1.530,  $\gamma$  1.531,  
 2V(meas.) 76°.  
 6.33 (8), 4.46 (8), 3.66 (10), 2.60 (8), 1.760 (8),  
 1.351 (8).

## IMA No. 91-043

The Sb-dominant member of the colusite  
 group.  
 $\text{Cu}_{26}\text{V}_2(\text{Sb},\text{Sn},\text{As})_6\text{S}_{32}$   
 Cubic: P43n  
 a 10.705 Å  
 Colour not observed because of the small size;  
 metallic; opaque.

- In reflected light: grey with a light-brown tint; no anisotropy, bireflectance, pleochroism or internal reflections; R (25.2%) 470 nm, (28.3%) 546 nm, (29.9%) 589 nm, (31.0%) 650 nm.  
3.10 (10), 1.892 (9), 1.614 (7), 1.226 (4), 1.094 (6), 1.030 (4).
- IMA No. 91-044**  
The Ge-dominant member of the colusite group.  
 $\text{Cu}_{26}\text{V}_2(\text{Ge},\text{As})_6\text{S}_{32}$   
Cubic:  $\bar{P}43n$   
a 10.568 Å  
Grey-black; metallic; opaque.
- In reflected light: greenish-yellow, olive-yellowish-cream; no internal reflections, anisotropy, bireflectance or pleochroism; R (23.8%) 470 nm, (27.3%) 546 nm, (27.9%) 589 nm, (27.9%) 650 nm.  
3.05 (10), 2.64 (4), 1.870 (5), 1.595 (3), 1.320 (3), 1.212 (3), 1.079 (3), 1.017 (5).
- IMA No. 91-045**  
 $(\text{Ca},\text{Mg})_{14}\text{Mn}_3\text{Zn}_3\text{Be}_6\text{Si}_{14}\text{O}_{50}(\text{OH},\text{F})_8$   
Monoclinic:  $P2_1/c$   
a 9.08, b 18.03, c 14.59(4) Å,  $\beta$  104.8°  
Colourless; vitreous; transparent.  
Biaxial (-),  $\alpha$  1.674,  $\beta$  1.680,  $\gamma$  1.681,  
2V(meas.) 29.0°, 2V(calc.) 44°.  
2.863 (100), 2.771 (40), 2.653 (50), 2.388 (50), 2.272 (30), 1.832 (30).
- IMA No. 91-046**  
The Cu-dominant analogue of geigerite and chudobaite.  
 $(\text{Cu},\text{Co})_5(\text{AsO}_4)_3(\text{AsO}_3\text{OH})_2 \cdot 10 \text{ H}_2\text{O}$   
Triclinic: P1 or  $\bar{P}1$   
a 8.033, b 10.374, c 6.446(5) Å,  $\alpha$  79.62°,  
 $\beta$  84.95°,  $\gamma$  86.21°  
Green; vitreous; transparent.  
Biaxial (+),  $\alpha$  1.634,  $\beta$  1.662,  $\gamma$  1.720,  
2V(meas.) 75°, 2V(calc.) 72°.  
10.2 (100), 8.01 (60), 4.001 (50), 3.667 (60), 3.151 (50), 3.063 (50).
- IMA No. 91-047**  
 $\text{Tl}_3\text{AsS}_4$   
Orthorhombic: Pnma  
a 8.894, b 10.855, c 9.079 Å  
Dark red; adamantine to submetallic; opaque to translucent.  
In reflected light: red, red internal reflections, strong anisotropy, strong bireflectance, no pleochroism.  $R_{\max}$  and  $R_{\min}$  are: (4.78, 3.93%) 481 nm, (4.64, 3.86%) 547 nm, (8.64, 7.81%) 591 nm, (13.72, 11.78%) 644 nm.
- 4.14 (M), 3.99 (S), 3.80 (M), 3.47 (MSb), 3.35 (M), 2.813 (VS), 2.537 (M), 2.264 (MSb).
- IMA No. 91-048**  
 $\text{Na}_{25}\text{Ba}(\text{Y},\text{Gd},\text{Dy})_2(\text{CO}_3)_{11}(\text{HCO}_3)_4(\text{SO}_4)_2\text{F}_2\text{Cl}$   
Hexagonal:  $P6_3/m$   
a 8.811, c 37.03(3) Å  
Light green to yellowish-green; vitreous; transparent.  
Uniaxial (-),  $\omega$  1.536,  $\epsilon$  1.510.  
4.79 (42), 3.32 (40), 2.829 (100), 2.659 (51b), 2.531 (71b), 2.270 (90).
- IMA No. 91-050**  
 $(\text{Ca},\text{REE},\text{Th})_{15}\text{As}^{+5}(\text{As}_{0.5}^{+3}\text{Na}_{0.5})\text{Fe}^{+3}\text{Si}_6\text{B}_4\text{O}_{40}\text{F}_7$   
Hexagonal:  $R\bar{3}m$   
a 10.795, c 27.336(4) Å  
Yellowish-green; vitreous; transparent.  
Uniaxial (-),  $\omega$  1.757,  $\epsilon$  1.722.  
2.993 (S), 2.950 (S), 1.839 (MS), 1.802 (MS), 1.686 (MS), 1.572 (MS).
- IMA No. 91-051**  
 $\text{Ag}_9\text{SbTe}_3(\text{S},\text{Se})_3$   
Monoclinic: P2, P2/m or Pm  
a 8.900, b 8.302, c 19.49 Å,  $\beta$  82.98°  
Colour unknown because of the small grain size; metallic; opaque.  
In reflected light: grey with faint green-blue hue, anisotropy present with brownish-grey tone, weak bireflectance, no pleochroism, no internal reflections,  $R_{\max}$  and  $R_{\min}$ , (38.0, 34.2%) 470 nm, (36.6, 32.2%) 546 nm, (35.7, 31.8%) 589 nm, (34.0, 30.2%) 650 nm.  
3.82 (6), 2.89 (4), 2.83 (4), 2.22 (10), 2.14 (3), 2.13 (4).
- IMA No. 91-052**  
The Sb-analogue of skutterudite.  
 $\text{CoSb}_3$   
Cubic: Im3  
a 9.0411 Å  
Tin-white; metallic; opaque.  
In reflected light: tin-white, isotropic, no bireflectance, nonpleochroic, no internal reflections. R (59.0%) 470 nm, (58.7%) 546 nm, (58.7%) 589 nm, (58.7%) 650 nm.  
2.85 (100), 2.01 (80), 1.92 (80), 1.84 (80), 1.50 (80), 1.185 (80), 1.147 (80), 0.780 (100).
- IMA No. 91-053**  
 $\text{Zn}_{12}(\text{CO}_3)_3(\text{SO}_4)(\text{OH})_{16}$   
Orthorhombic:  $P22_12$   
a 15.724, b 6.256, c 5.427(5) Å  
White; vitreous; translucent.  
Biaxial (probably +),  $\alpha$  1.635(3),  $\beta$  1.650(3),  $\gamma$  could not be measured, 2V about 60°.

15.44 (100), 7.88 (100), 5.25 (20), 2.714 (40),  
2.577 (20), 2.397 (20), 1.565 (30b).

**IMA No. 91-054**



Hexagonal:  $R\bar{3}$

a 16.025, c 19.773 Å


Colourless to pale brown; vitreous; transparent.

Uniaxial (-),  $\omega$  1.589,  $\epsilon$  1.586.

8.076 (80), 6.544 (90), 4.659 (75), 3.776 (90),  
3.159 (85), 2.683 (100).

**IMA No. 91-055**

A member of the epidote group, related to  
dolloseite-(Ce).



Monoclinic:  $P2_1/m$

a 8.903, b 5.748, c 10.107 Å,  $\beta$  113.41°

Dark greyish-brown; vitreous; transparent.

Biaxial (-),  $\alpha$  1.773,  $\beta$  1.790,  $\gamma$  1.803,

2V(meas.) 83°, 2V(calc.) 82°.

9.32 (2), 5.23 (2), 4.67 (2), 3.52 (4), 2.91 (10),  
2.73 (7), 2.63 (8), 1.437 (2).