

<b>Zeitschrift:</b>	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
<b>Band:</b>	75 (1995)
<b>Heft:</b>	2
<b>Artikel:</b>	Preliminary fission-track ages on zircons and apatites from the Sardona unit, Glarus Alps, eastern Switzerland : late Miocene-Pliocene exhumation rates
<b>Autor:</b>	Lihou, Joanne / Hurford, Anthony / Carter, Andrew
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-57149">https://doi.org/10.5169/seals-57149</a>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 28.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Preliminary fission-track ages on zircons and apatites from the Sardona unit, Glarus Alps, eastern Switzerland: late Miocene-Pliocene exhumation rates

by Joanne Lihou<sup>1</sup>, Anthony Hurford<sup>2</sup> and Andrew Carter<sup>2</sup>

## Abstract

Fission-track age determinations from a series of samples from the Infrahelvetic Sardona unit of eastern Switzerland yielded mixed Variscan and Alpine zircon fission-track ages with a mode of 160 Ma, and late Miocene-Pliocene apatite fission-track ages of  $3.5 \pm 0.1$  Ma and  $4.5 \pm 0.4$  Ma. These data suggest that there was an eastward younging in the onset of exhumation of material between the eastern Aar massif and the Rhine Valley. An exhumation rate of  $0.8 \pm 0.2$  mm/yr was calculated for the region, which may be associated with a Pliocene local surface uplift maximum centred around Chur that continues to the present day. Erosional unroofing is concluded to be the principal cause for the Miocene-Recent regional exhumation of material.

**Keywords:** fission-track dating, exhumation, Late Miocene-Pliocene, Sardona unit, Glarus overthrust, Switzerland.

## 1. Introduction

The cooling pathways for rocks whose peak paleotemperature was less than 300 °C cannot be determined from conventional Rb/Sr and K/Ar radiometric dating, whose systems have closure temperatures exceeding 300 °C (HURFORD, 1991). This is why the application of the zircon and apatite fission-track techniques, which have closure temperatures of 200–250 °C and 50–120 °C, respectively (HURFORD, 1991), has been so useful in the external part of the Alps. Reviews of fission-track (F-T) methodology and analytical methods were given by HURFORD et al. (1989) and HURFORD (1991), where the ways in which these data can be used were outlined:

- (i) Zircon and apatite cooling ages from the same sample can be used to derive an average cooling rate between the assumed closure temperatures for the two systems.
- (ii) The distribution of confined apatite lengths is diagnostic of the style of cooling.
- (iii) The present day altitude-dependence of apatite (or zircon) ages can be used to calculate

an average *exhumation/denudation rate*, due to the earlier passage of a sample at a higher elevation below the closure temperature for retention of tracks.

The last method avoids having to use an estimated paleo-geothermal gradient, but assumes that the 110 °C (or 240 °C) paleo-isotherm remained horizontal and was not influenced by topography or disturbed by rapid exhumation (PARRISH, 1983). STÜWE et al. (1994) have shown that perturbation of the steady-state paleo-isotherm mirrors surface topography in a damped fashion and is most pronounced for small wavelength, high amplitude topographic features (wavelengths of  $\leq 20$  km, amplitudes of  $\geq 3$  km). The critical isotherms also need to have resided at a constant depth relative to the earth's surface (PARRISH, 1983). If these conditions are not met, fictitious apparent exhumation rates can be produced by the downward relaxation of isotherms following a period of active tectonism, rapid uplift and erosion (PARRISH, 1983), or else the apparent exhumation rate is substantially overesti-

<sup>1</sup> Department of Earth Sciences, University of Oxford, Parks Road, Oxford OX1 3PR, U.K.

<sup>2</sup> London Fission Track Research Group, Department of Geological Sciences, University College, Gower Street, London WC1E 6BT, U.K.

mated (STÜWE et al., 1994). If these conditions are met, however, the apparent exhumation rate quantifies the approach of rocks to the earth's surface.

The most recent review and data compilation of F-T analyses in the French and Swiss Alps was made by HUNZIKER et al. (1992). The present paper adds to the data set and provides further constraints on the exhumation of the Glarus Alps and the recent updoming centred around Chur.

## 2. Geological setting

The *Sardona* unit of eastern Switzerland is located between Elm in Sernftal and Bad Ragaz in the Rhine valley (Fig. 1), an area of  $25 \times 10$  km, at elevations of 500 to 2700 m. It consists of Cenomanian-Campanian pelagic limestones and

marls, plus Maastrichtian to Bartonian flysch deposits (LIHOU, 1993). It is an allochthonous unit, being bounded by thrusts, which together with the underlying Blattengrat and North Helvetic Flysch (NHF) units, forms part of the *Infrahelvetic Complex*. The Infrahelvetic Complex also includes the autochthonous and paraautochthonous cover to the *Aar massif*, since it comprises all the tectonic units that structurally underlie the Helvetic nappes and are separated from them by the *Glarus Overthrust* (Fig. 2). The present topography of the Glarus Overthrust is a WSW-ENE trending culmination, with a southern flank sloping  $15-20^\circ$  and a northern flank sloping  $10-15^\circ$  (SCHMID, 1975), that plunges gently eastwards from Vättis to the Rhine valley (TRÜMPY and TROMMSDORFF, 1980). Hence, the *Sardona* unit crops out in tectonic windows through the culmination in the Overthrust (Fig. 1).




Fig. 1 Tectonic map of the Glarus Alps, eastern Switzerland, adapted from TRÜMPY (1967) and SPICHER (1980), also showing the line of cross-section for figure 2, fission-track sample localities and the anchizone/epizone boundary within the Infrahelvetic Complex (shaded area) compared with its location in the Verrucano of the Glarus (Helvetic) nappe (dashed line), taken from WANG et al. (1995); marginal numbers refer to the Swiss coordinate system and are marked at intervals of 10 km.




Fig. 2 Simplified true scale structural cross-section through the Glarus Alps, eastern Switzerland, modified from TRÜMPY (1980) and PFIFFNER (1986) showing the projected positions of fission-track sample localities; key as in figure 1.

Bl – Blattengrat unit; Dr – Drusberg nappe; Gl – Glarus nappe; M – Mürtschen nappe; NHF – North Helvetic Flysch unit; PAM – Parautochthonous Mesozoic; Sa – Sardona unit; V – Verrucano.

### 3. Regional review of Alpine metamorphism and exhumation history

The Helvetic zone and Infrahelvetic Complex were deformed during the Neo-alpine Orogeny (Oligocene-Miocene). The deformation was subdivided into four phases by MILNES and PFIFFNER (1977, 1980): the Pizol, Cavistrau, Calanda and Ruchi phases. Burial by Penninic and Austroalpine nappes led to *very low grade metamorphism* in the Infrahelvetic Complex and the Helvetic nappes (FREY et al., 1973, 1980; GROSHONG et al., 1984). A variety of methods have been used to quantify the degree of incipient metamorphism and thus unravel the tectonothermal evolution: critical mineral assemblages, illite "crystallinity", coal rank and fluid inclusion microthermometry. The techniques involved are reviewed in FREY (1986).

Early work using these techniques in the Helvetic Alps established that metamorphic zones ran parallel to the Alpine border, increasing in grade from north to south, from diagenetic grade at the Alpine border, to anchizone-epizone in the par-autochthonous cover to the Aar massif, and downwards in the nappe pile (FREY et al., 1973, 1980; GROSHONG et al., 1984). Moreover, the same work showed that metamorphic zones cross-cut tectonic boundaries, implying that the metamorphism was post-tectonic and was related to burial of the nappe pile and concomitant heating. In addition, a discontinuity in the metamorphic gradient was identified across the Glarus Overthrust, at the contact between the Helvetic nappes and the Infrahelvetic flysch units (FREY et al., 1980; FREY, 1988).

Recent, local metamorphic studies have focused on the Infrahelvetic flysch units (ERDELBROCK, 1994; RAHN et al., 1994, 1995; WANG et

al., 1995). By mapping iso-reflectance lines derived from vitrinite reflectance values, RAHN et al. (1995) and ERDELBROCK (1994) were able to estimate the amount of post-metamorphic horizontal offset across the Glarus Overthrust to be in the order of 5–10 km. WANG et al. (1995) also found that the offset of illite "crystallinity" zones was ~10 km (Fig. 1). Peak metamorphic temperatures were gauged directly from the temperature of homogenization of fluid inclusions, and also from vitrinite reflectance data using the calibration technique of BOSTICK et al. (1978), which depends on the effective heating time ( $t_{\text{eff}}$ ) (i.e., the duration of burial of a rock within 15 °C of its peak temperature). RAHN et al. (1994) found that paleotemperatures of 270–310 °C for the NHF unit in Linthal (Fig. 1) derived by the two methods, were concordant when the  $t_{\text{eff}}$  was assumed to be 1 Myr. ERDELBROCK (1994) also concluded that there had been a very short heating time of no more than 1 Myr, during which paleotemperatures reached 300–320 °C in the Sardona unit in Calfeisental (Fig. 1).

*Dating* of Alpine metamorphism has been constrained by Rb/Sr and K/Ar dating of radiogenic minerals, a review of which was made by HUNZIKER et al. (1992). Metamorphism in the Helvetic nappes was dated as early to middle Oligocene (30–35 Ma), from concordant K/Ar and Rb/Sr mineral ages, with a second, Miocene metamorphic phase at 20–25 Ma affecting the Infrahelvetic Complex (HUNZIKER et al., 1986). The timing of the earlier metamorphism, which post-dates Calanda phase emplacement of the Helvetic nappes onto the Infrahelvetic Complex (GROSHONG et al., 1984), is further constrained by the age of the youngest metamorphosed sediments found in the Infrahelvetic Complex (i.e., the Engi slates of the NHF unit) of early Oligo-

cene age, and the first appearance of diagenetic grade metamorphosed Helvetic pebbles in the Upper Marine Molasse of mid Miocene age (HOMEWOOD et al., 1986; HUNZIKER et al., 1986). Geochronological ages for the younger metamorphism may be as young as 14 Ma (HUNZIKER, in PFIFFNER, 1986) and would therefore indicate that post-metamorphic, Ruchi phase, movements on the Glarus Overthrust are post-mid Miocene.

MICHALSKI and SOOM (1990) reconstructed the *cooling and exhumation history* of the basement of the Helvetic zone, the Aar and Gotthard massifs, using the fission-track technique, incorporating new and previously-published data. They concluded that cooling in the east of the Aar massif began in the early Miocene, where zircon cooling ages ranged from 19–27 Ma. Confined track lengths in the apatites, of 13.5–14.5  $\mu\text{m}$ , were interpreted as indicating undisturbed steady cooling. Apatite ages gave an exhumation rate of 0.3–0.5  $\mu\text{m}/\text{yr}$  from 10–5 Ma (i.e., during the late Miocene).

RAHN (1994) measured apatite F-T ages for 14 samples from the NHF unit, the parautochthonous cover to the Aar massif, in a profile from Linthal (725 m) to the Ruchi mountain (3100 m) (Fig. 1). The apatite ages ranged between 5.4 and 8.6 Ma, i.e., mid-late Miocene, and showed a strong altitude-dependence, from which an exhumation rate of 0.65 mm/yr was calculated. An interesting result was the discontinuity in the apatite ages at the Glarus Overthrust, which may indicate that there has been movement along this thrust since the late Miocene.

#### 4. Aims and methods

A preliminary F-T study in the Sardona unit was undertaken in order to corroborate the results of MICHALSKI and SOOM (1990) and RAHN (1994). Three samples were collected from the *Sardona Quartzite Formation* at each of two localities in Calfeisental (Fig. 1). This Formation is a Paleocene to early Eocene arenitic flysch unit, that also includes a debris flow conglomerate, the *Crystalline Conglomerate*, containing many exotic sedimentary and crystalline clasts probably from the Austroalpine and Penninic realm (LIHOU, 1993). At each locality, samples of the Sardona Quartzite, the matrix to the Crystalline Conglomerate and a granitic clast from the Crystalline Conglomerate were collected. In this way, concordant ages from a particular locality would increase confidence in the results. It was also hoped that the granitic clasts would yield either Alpine zircon F-T cooling ages, or else pre-Alpine cooling

ages that would answer some petrological questions as to its origins.

In addition, one sample was collected from the *Guschakopf Sandstone*, of the Ragaz unit, at much lower elevations in the Rhine Valley. This sandstone is probably a lateral equivalent to the Sardona Quartzite.

Samples for *F-T analysis* were prepared from heavy mineral concentrates of apatite and zircon. Spontaneous fission-tracks were revealed in polished mounts of apatite using 5N  $\text{HNO}_3$  at 20 °C for 20 seconds, and of zircon using a binary eutectic of KOH : NaOH at 225 °C for 20–36 hours. Irradiation of the samples with thermal neutrons was made at the thermal facility of the Risø Reactor at the National Research Centre, Roskilde, Denmark. Induced tracks were recorded in mica external detectors and the fluence monitored using Corning glass dosimeters CN-5 (~ 11 ppm U) for apatite and CN-2 (~ 36 ppm U) for zircon irradiations. Mica detectors were etched in 40% HF at 20 °C for 40 minutes. Tracks were counted using Zeiss Axioplan microscopes with  $\times 100$  objectives and a maximum magnification of  $\times 1250$ . Central ages were calculated using the IUGS-recommended zeta calibration approach (HURFORD, 1990).

#### 5. Results

*F-T central ages* are presented in table 1. The granite clasts (FTA 1 and 6) gave essentially the same zircon age of about 165 Ma. Central zircon ages from the Sardona Quartzite and the Guschakopf Sandstone had a much wider range of ~ 99–208 Ma. Insufficient apatite crystals were retrieved from the Guschakopf Sandstone to yield an age. However, the samples at each locality in Calfeisental (FTA 1–6) yielded concordant apatite ages, increasing confidence in the results (Tab. 1). Taking an average age for each locality, the apatite ages from the Sardona unit are  $3.5 \pm 0.1$  Ma and  $4.5 \pm 0.4$  Ma (i.e., late Miocene–Pliocene), and are therefore much younger than the mid-late Miocene ages obtained by RAHN (1994) from the NHF unit, 20 km to the west, and by MICHALSKI and SOOM (1990) from the Aar and Gotthard massifs 65 km to the southwest. However, the apatite ages from the Sardona unit are similar to the young  $4.5 \pm 1.2$  Ma age obtained by MICHALSKI and SOOM (1990) from a nearby locality at Vättis (Fig. 1, Tab. 1). This suggests that there may be an eastward younging in the apatite ages that mirrors the westward younging identified by MICHALSKI and SOOM (1990) at the western end of the Aar massif.

Tab. 1. Fission-track apatite and zircon analytical data for the Calfeisental-Rhine valley region. Notes:

- (i) H (m) is the perpendicular distance to the Glarus Overthrust calculated from the structure contour map of SCHMID (1975);
- (ii) analyses were made by the external detector method using 0.5 for the  $4\pi/2\pi$  geometry correction factor;
- (iii) track densities ( $\rho$ ) are quoted as  $\times 10^6$  tracks  $\text{cm}^{-2}$  and N is the total number of tracks counted;
- (iv)  $P\chi^2$  is the probability of obtaining  $\chi^2$  value for v degrees of freedom, where v = no. crystals - 1;
- (v) ages of FTA samples were calculated using dosimeter glass CN-5 (apatite) with  $\zeta CN5 = 339 \pm 5$ , and CN-2 (zircon) with  $\zeta CN2 = 124 \pm 5$ , calibrated by multiple analyses of IUGS apatite and zircon age standards (see GALBRAITH, 1992);
- (vi) the central age is a modal age, weighted for different precision of individual crystals (see HURFORD, 1990);
- (vii) KAW samples are from MICHALSKI and SOOM (1990); these authors do not quote the number of crystals used to derive their zircon central age for sample KAW 3215.

Sample no. Rock type	Locality	Elevation (m) H (m)	mineral	No. of crystals	Dosimeter $\rho_a$	Spontaneous $\rho_s$	Induced $\rho_i$	Age Dispersion $\chi^2$ %	Age (Ma) $\pm 1\sigma$					
KAW 2608 Verrucano cgl	Schwarze Hömer (749.75/205.75)	2400 +300	apatite zircon	10 3	7.43 6.02	5744 5271	0.7 138.0	18 361	13.2 55.0	320 144	<2 15	- -	6.4 $\pm$ 1.7 96 $\pm$ 1.0	
FTA 1 Granite clast	Heubützlipass (741.5/200.7)	2400 -350	apatite zircon	10 12	1.347 0.475	9337 3289	0.022 24.76	9 2299	1.242 4.458	508 414	<1 <1	107 24.7	4.0 $\pm$ 1.4 161 $\pm$ 15	
FTA 2 Cgl matrix	Heubützlipass (741.5/200.7)	2400 -350	apatite zircon	10 -	1.347 -	9337 -	0.818 -	44 -	3.715 -	1998 -	70 -	0.03 -	5.0 $\pm$ 0.8 -	
FTA 3 Quartzite	Heubützlipass (741.5/200.7)	2400 -350	apatite zircon	9 16	1.199 0.475	8306 3289	0.065 27.10	13 3381	2.886 3.688	580 460	<1 <1	80 66.8	0.0 66.8	4.6 $\pm$ 1.3 208 $\pm$ 36
FTA 4 Quartzite	Troseggobel (741.8/198.0)	1830 -1270	apatite zircon	11 -	0.475 -	3289 -	17.49 -	1745 -	5.061 -	505 -	<1 -	- -	- -	
FTA 5 Cgl matrix	Troseggobel (741.8/198.0)	1830 -1270	apatite zircon	20 -	1.199 -	8306 -	0.050 -	43 -	3.089 -	2642 -	15 -	39.7 -	3.4 $\pm$ 0.6 -	
FTA 6 Granite clast	Troseggobel (741.8/198.0)	1830 -1270	apatite zircon	20 19	1.199 0.475	8306 3289	0.093 20.02	70 3302	5.273 3.480	3961 574	30 <1	18.9 52.4	3.6 $\pm$ 0.4 169 $\pm$ 22	
KAW 3215 mylonite	Vättis (751.30/197.70)	1060 -2140	apatite zircon	10 ?	7.81 9.24	4871 4461	0.58 1.0	14 29	15.7 24.4	374 82	7 69	- -	4.5 $\pm$ 1.2 19.1 $\pm$ 4.2	
R13 Sandstone	Bad Ragaz (756.3/208.1)	600	apatite zircon	20	0.475	3289	19.81	2746	3.723	516	<1	- -	80.5 147 $\pm$ 28	

No track length data from the apatite samples is available because of low spontaneous track densities.

## 6. Discussion and interpretation

### 6.1. ZIRCON AGES

The concordance of zircon central ages for the granite clasts implies that either they have the same provenance or the same thermal history. Examination of a radial plot for sample R13 from the Guschakopf Sandstone (Ragaz unit) revealed an equivalent mode, indicated by a cluster of zircons close to 160 Ma (Fig. 3), suggesting that the sand may have been derived from weathering of the same type of granitic terrane as the clasts in the Crystalline Conglomerate. Another, 50 Ma mode (i.e., early Eocene age) in the zircon ages was obtained from 20% of the crystals analyzed and is close to the depositional age of the sandstone (Fig. 3). This age may be a true Alpine cooling age for the zircons, or alternatively, it could date the cooling of newly-formed igneous zircons. The grains possessing a 50 Ma age exhibited euhedral to subhedral morphology, suggesting that they represent first cycle detritus. WINKLER

et al. (1990) obtained a central zircon age of  $57.8 \pm 2.7$  Ma from a Paleocene bentonite horizon in the Schlieren Flysch which WINKLER (1983) concluded was derived from a subduction-related magmatic arc. The zircons from the Guschakopf Sandstone possessing a 50 Ma age could therefore have been derived from weathering of a similar volcanic source.

Zircon ages in the Sardona unit are probably *mixed Variscan and Alpine ages*. MICHALSKI and SOOM (1990) also found that zircon ages north of the Aar massif were a mixture of Variscan and Alpine cooling ages, implying that the peak Alpine paleotemperature was below the closure temperature for zircon and that the tracks were therefore not completely annealed. They concluded that the temperature range for the zircon mixed age zone (or partial annealing zone) was 200–250 °C. However, ERDELBROCK (1994) determined peak paleotemperatures of 300–320 °C for this region based upon vitrinite reflectance measurements, which are well in excess of the accepted closure temperature for zircon. It must be born in mind that these vitrinite reflectance values are at the upper limit of applicability of the technique and the paleotemperatures may therefore be slightly overestimated. Nevertheless, paleotemperatures exceeding 250 °C probably were achieved since samples FTA 1–3 lie within the anchizone/epizone transition (270–310 °C, RAHN et al., 1994) and samples FTA 4–6 fall within the epizone defined by WANG et al. (1995) (Fig. 1), so the zircon fission tracks should have been reset. Hence, an alternative explanation for the mixed zircon ages in this area must be sought.

Closure temperature also depends on the cooling rate, faster cooling producing an elevated closure temperature (HUNZIKER et al., 1992). However, MICHALSKI and SOOM (1990, Fig. 11) determined a very slow initial cooling rate of 7 °C/Myr from the early Miocene to Pliocene (19.1–4.5 Ma) for a sample from Vättis (Fig. 1), although this calculation assumed that the zircon closure temperature was only 225 °C and the apatite closure temperature was 120 °C. One remaining possibility is that the *effective heating time* for the peak metamorphism was too short (~1 Myr – see above) for many of the zircons to have had enough time to be completely reset. Newly generated zircon annealing data at laboratory and geological timescales have revealed that for effective heating times of  $10^7$  yr, the zircon partial annealing zone lies between 210 °C and 350 °C, but if the heating time is reduced to  $10^6$  yr, the partial annealing zone is shifted to 230–370 °C (YAMADA et al., 1995; TAGAMI et al., 1995). In light of this information, minor levels of track an-




Fig. 3 Radial plot displaying individual zircon grain ages for sample R13 against a standardized error (left hand side); grains with increasing precision are closer to the radial axis on the right. This particular sample, from the early Eocene Guschakopf Sandstone, statistically comprises three age modes; (I) the oldest modal age comprises Paleozoic zircons; (II) the second modal age of ~160 Ma is similar to the central zircon ages of granite clasts from the Crystalline Conglomerate (FTA 1 and 6); and (III) the youngest age of ~50 Ma is within error of the stratigraphic age – see text for discussion.

nealing that would be expected for the Sardona samples if the effective heating time was close to  $10^6$  yr.

## 6.2. APATITE AGES

The apatite results from the Sardona unit were combined with two of MICHALSKI and SOOM's (1990) results, from the sites at Vättis and Schwarze Hörner at the east end of Calfeisental (Tab. 1 and Fig. 1). There appears to be poor cor-




Fig. 4 (a) Scatter plot of apatite age versus elevation in the region Calfeisental-Pizol, showing poor correspondence, with sites FTA 1-3 and KAW 2608 from equivalent elevations above mean sea level (AMSL) displaying a range of central ages, although these ages could be considered equivalent within statistical error. Treating the FTA and KAW data sets separately, to take into account the geographical separation of 3-5 km between these localities and the possible effects of perturbations in the 110 °C isotherm, would result in exhumation rates of 0.6 and 0.7 mm/yr, respectively. (b) Scatter plot of apatite age versus perpendicular distance from the Glarus Overthrust in the region Calfeisental-Pizol, showing better correspondence than graph (a), although samples KAW 3215 and FTA 1-3 give the same central age, despite the fact that they reside at different distances from the Glarus Overthrust – see text for discussion.

KAW data points are taken from MICHALSKI and SOOM (1990), error bars shown are  $1\sigma$  and regression lines were fitted to the data using a computer-aided graphics programme.

respondence between apatite age and elevation (Fig. 4a), but better correspondence between age and perpendicular distance from the Glarus Overthrust (Fig. 4b). Sample KAW 3215 from Vättis plots to the right of the best regression line for these data, i.e., towards higher ages (Fig. 4b); its earlier cooling may have been caused by local Pliocene valley incision, when the Paleo-Rhine Valley ran through Kunkelpass, Vättis and Raagaz, only later deviating to its current course via Chur and Landquart (Adrian Pfiffner, pers. comm.; Fig. 1).

RAHN (1994) also reached the conclusion that there was better correspondence for his data from the NHF unit when he plotted apatite age against perpendicular distance from the Glarus Overthrust; samples varying in their perpendicular distance from the Glarus Overthrust exhibited a range of ages that could not be ascribed to differences in altitude, since they were derived from a horizontal sampling profile within Linthal. Using the Glarus Overthrust as the reference plane, an apparent uplift rate of  $0.8 \pm 0.2$  mm/yr was generated for the data from Calfeisental-Pizol (Fig. 4b), which is greater than the rate of  $0.3 \pm 0.1$  mm/yr generated for the NHF unit generated by the same method (RAHN, 1994), or the rate of 0.3-0.5 mm/yr for the eastern Aar massif from MICHALSKI and SOOM (1990), although this was calculated using an age-altitude relationship. Combined with the eastward younging in the apatite F-T ages, this suggests that there was later but faster exhumation of material towards the east. No displacement of the apatite ages across the Glarus Overthrust was detected, as in the case of the NHF unit (RAHN, 1994).

Despite the caution needed in interpreting apparent *exhumation rates*, the rate calculated for this study can be accepted as a reasonable value for the Glarus Alps for the following reasons. During the Alpine metamorphic phase, the Helvetic nappes and Infrahelvetic Complex were buried beneath a Penninic and Austroalpine nappe pile that was 12 km thick (GROSHONG et al., 1984). The Glarus Overthrust, which marks the boundary between these tectonic units is itself now exposed at elevations of 1000-3000 m (SCHMID, 1975), attesting to the exhumation of once-buried rock. As a horizontal feature that was independent of late Miocene-Pliocene surface topography, using the Glarus Overthrust as a reference plane for apatite fission-track dates, tests whether the 110 °C paleo-isotherm was significantly perturbed by topography on the scale of the spacing between sample sites and therefore likely to give spurious exhumation rates. The fact that a better correspondence between apa-

ite ages and perpendicular distance from the Glarus Overthrust was observed both in this study (Fig. 4b) and that of RAHN (1994) may mean that the Glarus Overthrust was parallel to the critical paleo-isotherm, and may still have been a planar feature, only attaining its current arched geometry since 3.5 Ma. In addition, our own structural investigations of the north-dipping limb of the Glarus Overthrust near Pizol (Fig. 1) have revealed that normal displacements of up to a few metres on northward-dipping, steeply-inclined cleavage-parallel fault planes commonly affect the main thrust boundary. These minor normal faults may have accommodated arching of the Glarus Overthrust and would therefore support our hypothesis that this arching post-dates the main, Calanda phase thrust displacement. This interpretation challenges PFIFFNER's (1985) conclusion that the Glarus Overthrust was arched by Calanda phase shortening of the Infrahelvetic Complex in its footwall.

Valley relief in Calfeisental reaches 2000 m near to its mouth at Vättis, and its separation from the adjacent, parallel valley of Weisstannental to the north is 6–10 km (Fig. 1), meaning that the present day topography can be modelled with an amplitude of 2000 m and a wavelength of 12–20 km. For a topographic wavelength of 12–20 km where perturbations of the critical paleo-isotherm existed, differences in its depth would have been important for samples separated by lateral distances of a quarter wavelength, i.e., 3–5 km; this is the spacing between sites in Calfeisental. Assuming that the current topography is a reasonable analogue for the topography of the Glarus Alps in the late Miocene-Pliocene, or at least that the area was no more rugged than at present, figure 6 of STÜWE et al. (1994) can be used to estimate the perturbation of the critical paleo-isotherm. Assuming that steady-state had been reached, a denudation rate of 0.8 mm/yr would have caused perturbations in the isotherm ranging between 200 and 500 m for a topographic wavelength of 12–20 km. From equation (4) of STÜWE et al. (1994), this range of perturbations would result in a 10–25% ( $\pm 0.2$  mm/yr) overestimation in the actual exhumation rate. We do not feel that this represents a significant error in the apparent exhumation rate given the uncertainties in the fission-track analytical procedure and in view of the fact that this is the same as the error involved in the graphical calculation of the exhumation rate (Fig. 4b). We also feel that the geothermal gradients assumed in STÜWE et al.'s (1994) model may not be applicable to the Alps in the Miocene, and coupled with the gross ap-

proximations required in applying this model to the Glarus Alps, the calculation is unjustified.

Indirect evidence that uplift of the area since the early Miocene has indeed been driven by erosion of material can be provided using a simple calculation that was developed by BROWN (1991) and has been used in other mountain belts (FOSTER et al., 1994). Adopting a paleo-geothermal gradient of 35 °C/km (ERDELBROCK, 1994), and assuming a mean paleo-surface temperature of 10 °C, the depth to the pre-late Miocene 110  $\pm$  10 °C isotherm was  $\sim$  3000 m. Hence, the amount of material removed since the late Miocene is at least 3000 m. Added to this, localities which passed through this isotherm in the late Miocene are now elevated to  $\sim$  2500 m (samples FTA 1–3 and KAW 2608), i.e., 500 m above the present mean surface elevation in the Glarus Alps, so an additional 500 m of material has been removed assuming that the present mean surface elevation is similar to that in the late Miocene. Therefore, at least 3500 m of regional denudation can be accounted for since the late Miocene.

Tectonic uplift of the Aar massif and its cover nappes was initially accommodated by basement shortening on folds and/or faults (BURKHARD, 1990) associated with the last stages of continent-continent collision. BURKHARD (1990) concluded that its root zone was already backfolded and steepened before 15 Ma (i.e., mid-Miocene) because there is no discontinuity in the apatite ages across the present day structure. However, the faster, younger, late Miocene to Recent exhumation of the eastern Aar massif and overlying nappes could reflect lateral variations in the isostatic response to continued tectonic thickening of the European crust. Alternatively, erosion of material following tectonism and mountain building may have induced passive isostatic rebound of the crust. The maximum present local (geodetic) surface uplift rate in the region, of 1.4 mm/yr, is centred around Chur (JEAN-RICHARD, 1975; SCHAER et al., 1975). The increase in the Miocene-Pliocene exhumation rates in this direction, documented above, may indicate that a Pliocene local surface uplift maximum centred around Chur continues to the present day. This hypothesis supports the conclusion of FLISCH (1986, in MICHALSKI and SOOM, 1990) that the beginning of the Chur uplift was around 3 Ma.

## 7. Conclusions

Fission-track age determinations on zircons and apatites from a series of samples from the Infrahelvetic Sardona unit of eastern Switzerland pro-

vide new information about the exhumation pattern of the eastern Aar massif and the overlying Glarus nappes:

(a) Zircon ages are mixed Variscan and Alpine ages, exhibiting a modal age of 160 Ma. Partial annealing of the tracks, despite peak paleotemperatures of around 300 °C, may be explorable by the short effective heating time for the peak metamorphism of only 1 Myr.

(b) Apatite cooling ages from the Sardona unit of  $3.5 \pm 0.1$  Ma and  $4.5 \pm 0.4$  Ma (i.e., late Miocene-Pliocene) suggest that there was an eastward younging in the onset of exhumation of material between the eastern Aar massif and Calfeisental.

(c) Apatite ages show better correspondence between age and perpendicular distance from the Glarus Overthrust than between age and elevation above mean sea level, suggesting that arching of the Glarus Overthrust may be a very young feature.

(d) The calculated exhumation rate for the region Calfeisental-Pizol was  $0.8 \pm 0.2$  mm/yr and may indicate that there was a Pliocene surface uplift maximum centred around Chur that continues to the present day.

(e) At least 3500 m of denudation can be accounted for since the early Miocene, suggesting that erosional surface processes are the principal cause for the Miocene-Recent regional exhumation of material.

### Acknowledgements

This work was undertaken by J.L. as part of D.Phil. research project at the University of Oxford, supervised by Philip Allen and John Platt, and funded by a student sponsorship provided by Shell International Petroleum Company Limited. The F-T project was conceived in collaboration with Meinert Rahn, who has continued to contribute towards the end product through fruitful discussions. Rod Brown and Kit Johnson suggested significant improvements to the original version of the manuscript. Diane Seaward and Adrian Pfiffner are thanked for their thorough, challenging and speedy reviews of the manuscript.

### References

BOSTICK, N.H., CASHMAN, S.M., MCCULLOGH, T.H. and WADELL, T.H. (1978): Gradients of vitrinite reflectance and present temperature in the Los Angeles and Ventura basins, California. In: OLTZ, D.F. (ed.): Low Temperature Metamorphism of Kerogen and Clay Minerals. Pacific Section, Soc. Econ. Paleont. Mineral., Los Angeles, 65-89.

BROWN, R.W. (1991): Backstripping apatite fission-track "stratigraphy": A method for resolving the erosional and isostatic rebound components of tectonic uplift histories. *Geology* 19/1, 74-77.

BURKHARD, M. (1990): Aspects of the large-scale Miocene deformation in the most external part of the Swiss Alps (Subalpine Molasse to Jura fold belt). *Eclogae geol. Helv.* 83/3, 559-583.

ERDELBROCK, K. (1994): Diagenese und schwache Metamorphose im Helvetikum der Ostschweiz (Inkohlung und Illit- "Kristallinität"). Unpublished Ph. D. Thesis, RWTH, University of Aachen.

FOSTER, D.A., GLEADOW, A.J.W. and MORTIMER, G. (1994): Rapid Pliocene exhumation in the Karakoram (Pakistan) revealed by fission-track thermochronology of the K2 gneiss. *Geology* 22, 19-22.

FREY, M. (1986): Very low-grade metamorphism of the Alps - an introduction. *Schweiz. Mineral. Petrogr. Mitt.* 66/1/2, 13-27.

FREY, M. (1988): Discontinuous inverse metamorphic zonation, Glarus Alps, Switzerland: evidence from illite 'crystallinity' data. *Schweiz. Mineral. Petrogr. Mitt.* 68/2, 171-183.

FREY, M., HUNZIKER, J.C., ROGGWILLER, P. and SCHINDLER, C. (1973): Progressive niedriggradige Metamorphose glaukonitführender Horizonte in den helvetischen Alpen der Ostschweiz. *Contrib. Mineral. Petrol.* 39, 185-218.

FREY, M., TEICHMÜLLER, M., TEICHMÜLLER, R., MULLIS, J., KÜNZI, B., BREITSCHMID, A., GRUNER, U. and SCHWIZER, B. (1980): Very low-grade metamorphism in the external parts of the Central Alps: illite crystallinity, coal rank and fluid inclusion data. *Eclogae geol. Helv.* 73, 173-203.

FLISCH, M. (1986): Die Hebungs geschichte der ostalpinen Silvretta-Decke seit der mittleren Kreide. *Bull. Ver. schweiz. Petroleum-Geol. u. Ing.* 53, 23-49.

GALBRAITH, R. (1992): Statistical models for mixed ages. Abstracts with programs, 7th International Workshop on Fission Track Thermochronology, University of Pennsylvania, 7.

GROSHONG, R.H. JR., PFIFFNER, O.A. and PRINGLE, L.R. (1984): Strain partitioning in the Helvetic thrust belt of Eastern Switzerland from the leading edge to the external zone. *Journal of Structural Geology* 6(1/2), 5-18.

HOMEWOOD, P., ALLEN, P.A. and WILLIAMS, G.D. (1986): Dynamics of the Molasse basin of western Switzerland. In: ALLEN, P.A. and HOMEWOOD, P. (eds): Foreland Basins. Int. Assoc. Sediment. Spec. Publ. 8, 199-217.

HUNZIKER, J.C., FREY, M., CLAUER, N., DALLMEYER, R.D., FRIEDRICHSEN, H., FLEHMIG, W., HOCHSTRASSER, K., ROGGWILER, P. and SCHWANDER, H. (1986): The evolution of illite to muscovite: mineralogical and isotopic data from the Glarus Alps, Switzerland. *Contrib. Mineral. Petrol.* 92, 157-180.

HUNZIKER, J.C., DESMONS, J. and HURFORD, A.J. (1992): Thirty-two years of geochronological work in the Central and Western Alps: a review on seven maps. *Mém. Géol. (Lausanne)*, No. 13, pp. 59.

HURFORD, A.J. (1990): Standardization of fission track dating calibration: recommendation by the Fission Track Working Group of the IUGS Subcommission on Geochronology. *Chem. Geol. (Isotope Geoscience section)* 80, 171-178.

HURFORD, A.J. (1991): Uplift and cooling pathways derived from fission track analysis and mica dating: a review. *Geol. Rdsch.* 80/2, 349-368.

HURFORD, A.J., FLISCH, M. and JÄGER, E. (1989): Unravelling the thermo-tectonic evolution of the Alps: a contribution from fission track analysis and mica

dating. In: COWARD, M.P., DIETRICH, D. and PARK R.G. (eds): Alpine Tectonics. Geol. Soc. London Spec. Publ. 45, 369–398.

JEANRICHARD, F. (1975): Summary of geodetic studies of recent crustal movements in Switzerland. Tectonophysics 29, 289–292.

LIHOU, J. (1993): The earliest evolution of the North Helvetic foreland basin as recorded by the Sardona unit of eastern Switzerland. Abstract supplement No. 1, TERRA nova 5, 67.

MICHALSKI, I. and SOOM, M. (1990): The Alpine thermo-tectonic evolution of the Aar and Gotthard massifs, Central Switzerland: Fission Track ages on zircon and apatite and K–Ar mica ages. Schweiz. Mineral. Petrogr. Mitt. 70/3, 373–388.

MILNES, A.G. and PFIFFNER, O.A. (1977): Structural development of the Infrahelvetic Complex, eastern Switzerland. Eclogae geol. Helv. 70/1, 83–95.

MILNES, A.G. and PFIFFNER, O.A. (1980): Tectonic evolution of the Central Alps in the cross section St. Gallen–Como. Eclogae geol. Helv. 73/2, 619–633.

PARRISH, R. (1983): Cenozoic thermal evolution and tectonics of the Coast Mountains of British Columbia 1: Fission track dating, apparent uplift rates and patterns of uplift. Tectonics 2, 601–631.

PFIFFNER, O.A. (1985): Displacements along thrust faults. Eclogae geol. Helv. 78/2, 313–333.

PFIFFNER, O.A. (1986): Evolution of the north Alpine foreland basin in the Central Alps. In: ALLEN, P.A. and HOMEWOOD, P. (eds): Foreland Basins. Int. Assoc. Sediment. Spec. Publ. 8, 219–228.

RAHN, M. (1994): Incipient Metamorphism of the Glarus Alps: Petrology of the Taveyanne Greywacke and Fission Track Dating. Unpublished Ph. D. Thesis, University of Basel. pp. 42.

RAHN, M., MULLIS, J., ERDELBROCK, K., FREY, M. (1994): Very low-grade metamorphism of the Taveyanne greywacke, Glarus Alps, Switzerland. J. met. Geol. 12/5, 625–641.

RAHN, M., MULLIS, J., ERDELBROCK, K. and FREY, M. (1995): Alpine Metamorphism in the North Helvetic Flysch of the Glarus Alps, Switzerland. Eclogae geol. Helv. 88/1, 157–178.

SCHAER, J.P., REIMER, G.M. and WAGNER, G.A. (1975): Actual and ancient uplift rate in the Gotthard region, Swiss Alps: a comparison between precise levelling and fission-track apatite age. Tectonophysics 29, 293–300.

SCHMID, S.M. (1975): The Glarus Overthrust: field evidence and mechanical model. Eclogae geol. Helv. 68/2, 247–280.

SPICHER, A. (1980): Tektonische Karte der Schweiz, 1 : 500 000 (2nd edition). Schweiz. Geol. Kommission. Kümmerly and Frey, Bern.

STÜWE, K., WHITE, L. and BROWN, R. (1994): The influence of eroding topography on steady-state isotherms. Application to fission track analysis. Earth Planet. Sci. Letters 124, 63–74.

TAGAMI, T., CARTER, A. and HURFORD, A.J. (1995): Natural long-term annealing of the zircon fission-track system in a Vienna Basin deep borehole: constraints upon the closure temperature and partial annealing zone. Chem. Geol. (Isotope Geoscience section), in press.

TRÜMPY, R. (1967): Geologischer Führer der Schweiz, Heft 7. Wepf & Co., Basel, 649–696.

TRÜMPY, R. (1980): Geology of Switzerland, a Guide-Book, Part A: An outline of the geology of Switzerland. Wepf & Co., Basel, pp. 104.

TRÜMPY, R. and TROMMSDORFF, V. (1980): Excursion No. IV: Alps of Eastern Switzerland. In TRÜMPY, R. (ed.), Geology of Switzerland, a Guide-Book. Wepf & Co., Basel, 211–260.

WANG, H., FREY, M. and STERN, W.B. (1995): Diagenesis and incipient metamorphism of the Helvetic Alps, eastern Switzerland: clay mineral data. Schweiz. Mineral. Petrogr. Mitt. 75, 187–199.

WINKLER, W. (1983): Stratigraphie, Sedimentologie und Sedimentpetrographie des Schlieren-Flysches (Zentralschweiz). Beiträge zur Geologischen Karte der Schweiz, Neue Folge 158. Schweiz. Geol. Komm., pp. 105.

WINKLER, W., HURFORD, A.J., VON SALIS PERCH-NIELSON, K. and ODIN, G.S. (1990): Fission-track and nannofossil ages from a Paleocene bentonite in the Schlieren Flysch (Central Alps, Switzerland). Schweiz. Mineral. Petrogr. Mitt. 70/3, 389–396.

YAMADA, R., TAGAMI, T., NISHIMURA, S. and ITO, H. (1995): Annealing kinetics of fission tracks in zircon: an experimental study. Chem. Geol. (Isotope Geoscience section), in press.

Manuscript received December 13, 1994; revision accepted March 7, 1995.