**Zeitschrift:** Schweizerische mineralogische und petrographische Mitteilungen =

Bulletin suisse de minéralogie et pétrographie

**Band:** 77 (1997)

Heft: 3

**Artikel:** Pb isotope and trace element signatures of polymetamorphic rocks from

the Silvretta nappe, a comparison

Autor: Köppel, Viktor / Hansmann, Werner / Maggetti, Marino

**DOI:** https://doi.org/10.5169/seals-58488

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 25.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Pb isotope and trace element signatures of polymetamorphic rocks from the Silvretta nappe, a comparison

by Viktor Köppel<sup>1</sup>, Werner Hansmann<sup>1</sup> and Marino Maggetti<sup>2</sup>

#### Abstract

The polymetamorphic crystalline basement of the Silvretta nappe is composed of paragneisses, amphibolites, eclogites and igneous rocks which are broadly subdivided into an older, Late Precambrian to Cambrian group comprising gabbros, diorites, tonalites and granites and a younger, Ordovician group of mainly granites. Lead isotope signatures of major mineral phases from the magmatic rocks complement the whole rock trace element data and support in most cases the petrogenetic conclusions obtained from trace element signatures. Similar proportions of mantle and crustal lead were observed in meta-volcanic and basic to intermediate meta-igneous rocks with geochemical signatures indicative of a MORB, IAB and I-type VAG origin of their protoliths. Higher proportions of crustal lead were observed in an IAB-type gabbro. A MORB type eclogite and an S-type augengneiss contained crustal lead. The lead of a younger S-type COLG granitic gneiss is dominated by crustal lead, and metasediments were found to contain crustal lead with the highest  $\mu$  value observed in the region.

The data pattern suggests that the magmas which originated from a depleted mantle source were contaminated by a crustal Pb-component from the metasediments. The lead isotope ratios of meta-igneous rocks indicate an increasing contribution of crustal lead as cratonisation progressed and thereby reflect the evolution of the crystalline core of the Silvretta nappe from a primitive island arc to a mature crust.

Keywords: Pb isotopes, trace element data, depleted mantle, crustal lead, cratonisation, Austroalpine.

## Introduction

While geochemical characteristics of magmatic rocks commonly serve to distinguish different tectonic settings, lead isotope signatures reflect the principal reservoirs of the lithosphere in which their magmas were generated. These major reservoirs are the upper crust where high  $\mu\text{-} (= ^{238}\text{U}/^{204}\text{Pb})$  and high W-values (=  $^{232}\text{Th}/^{204}\text{Pb})$  prevail, the lower crust with low  $\mu\text{-}$  and high W-values, and the normal, i.e. more or less depleted mantle with low  $\mu\text{-}$  and W-values. In contrast to most trace elements, lead concentrations of mantle derived rocks are low and their Pb-isotope ratios are therefore sensitive to contamination and therefore offer the opportunity to detect minor contaminations.

The formation of new continental crust commonly begins with the formation of primitive island arcs along subduction zones. At this stage mantle material and sediments will be involved, possibly also fragments of old continental crust. Active island arcs will progressively cratonise and accrete to continental margins. The lead isotope pattern of the accompanying magmatism will reflect this trend. In the early formed rocks the mantle signature of lead will remain discernible while at later stages, when sediments and possibly older crustal material are increasingly involved in orogenic processes, the rocks will obtain a transient lead isotope signal of the average continental crust. The lead isotopes will then evolve depending on the  $\mu$ - and W-values of the newly formed rocks (e.g. Doe and Zartman, 1979).

The composition of the lead isotope signal at the time of crystallization or metamorphic recrystallization is best represented in minerals with very low  $\mu$ - and W-values such as feldspars from igneous rocks or galenas from associated sulfide deposits. Such lead-isotopic ratios reflect the time

<sup>&</sup>lt;sup>1</sup> Departement für Erdwissenschaften, ETH-Zentrum, CH-8092 Zürich, Switzerland. E-mail: koeppel@erdw.ethz.ch.

<sup>&</sup>lt;sup>2</sup> Institut de Minéralogie et Pétrographie, Université de Fribourg, Pérolles, CH-1700 Fribourg, Switzerland.

integrated  $\mu$ - and W-values of a relatively large volume of the crust and thus they are reasonably representative for the continental crust. The evolution of average continental crustal lead has been modelled, for example, by STACEY and KRAMERS (1975).

This study aimed at testing whether or not lead isotopes of polymetamorphic igneous rocks yield information that is compatible with their geochemical signatures. Previous lead isotope studies of metamorphic rocks demonstrated that the mobility of lead across lithological boundaries is generally restricted to a few tens of meters or even less (Cumming et al., 1987; Peretti and Köppel, 1986). Curti (1987) demonstrated that the lead in mica-poor metapsammites from the Monte Rosa area did not homogenise under Alpidic lower am-

phibolite facies conditions. Mobility is, however, greatly enhanced by fluid flow in shear zones and in rocks with abundant water bearing phases such as micas.

Ten samples representative of the major lithologic units of the Silvretta basement rocks were selected for lead isotope analyses (Tab. 3). They comprise a MORB-type eclogite and an amphibolite, two gabbros with an island-arc signature, a metatonalite and metadiorite with a volcanic arc signature, an older, S-type augengneiss with a volcanic arc and a younger S-type augengneiss with a collisional granite signature. Two paragneiss samples were included. Their major and trace element compositions are listed in table 1 and 2. The characterization of the older and younger orthogneisses is based on the SiO<sub>2</sub>–Zr/TiO<sub>2</sub> (FLOYD and WINCHES-




Fig. 1 Generalised geologic map of the Silvretta thrust sheet (after MAGGETTI and FLISCH, 1993).

Sample numbers: 1 = RB 64, 2 = RB 92, 3 = Ma 640, 4 = Ma 124a, 5 = ST 221, 6 = Ma 1061, 7 = Ma 252, 8 Ma 193, 9 = Ma 943, 10 = Si 586.

Tab. 1 XRF Analyses.

Type Sample weight%	A Ma 640	E Ma 124a	O St 221	O Ma 1061	O Ma 252	O Ma 193	O Ma 943	Y Si 586
SiO <sub>2</sub>	49.19	48.50	49.83	54.72	57.84	67.73	72.99	75.86
TiO <sub>2</sub>	2.26	1.68	0.42	0.60	0.53	0.29	0.39	0.11
$Al_2O_3$	14.82	16.10	12.12	19.02	16.33	16.10	13.49	12.67
$Fe_2O_3$	6.29	3.46	1.52	2.57	2.27	1.21	0.44	0.22
FeO	7.25	9.01	8.91	4.42	5.71	2.26	2.07	0.96
MnO	0.34	0.21	0.21	0.14	0.14	0.09	0.04	0.03
MgO	5.94	7.21	16.21	4.93	4.10	1.59	1.04	0.30
CaO	6.98	11.11	5.55	8.64	8.08	4.21	1.23	0.42
Na <sub>2</sub> O	4.68	3.72	1.87	3.27	2.71	4.18	2.64	3.03
K <sub>2</sub> O	0.58	0.14	0.42	0.22	0.80	1.41	3.68	0.96
$P_2O_5$	0.31	0.14	0.02	0.16	0.11	0.12	0.10	0.13
L.O.I.	1.08	0.04	2.75	1.76	0.63	1.06	1.75	1.08
Total	100.08	100.36	99.83	100.45	99.35	100.25	99.86	99.77
$Fe_{tot}(Fe_2O_3)$	14.35	13.65	11.42	7.48	8.71	3.63	2.74	1.28
ppm								
Nb	19	6	1	2	_	_	7	6
Zr	109	98	18	54	55	105	142	67
Y	41	66	2	12	13	9	21	17
Sr	147	47	249	618	238	334	148	41
Rb	2	3	4	2	30	38	107	203
Th	_	1	1	-	<u> -</u>	7	14	14
Pb	_	10	1	3	_	_	25	17
Ga	21	16	9	15	17	18	15	13
Zn	113	101	80	73	112	65	44	13
Cu	46	6	1	1	22	18	19	14
Ni	36	63	61	45	20	15	22	2
v	342	321	169	213	238	62	39	2 8
Cr	77	199	747	115	52	25	26	9
Ba	127	31	83	159	318	726	769	100

L.O.I. = loss on ignition, P = paragneiss, E = eclogite, A = amphibolite, O = older orthogneiss, Y = younger orthogneiss

TER, 1978) and the Zr/TiO<sub>2</sub>-Nb/Y (WINCHESTER and FLOYD, 1977) discrimination diagrams.

## Geologic setting

The Silvretta thrust sheet (Fig. 1) is part of the pre-Permian basement of the Alps. MAGGETTI et al. (1990) distinguish four major groups of crystalline rocks: (1) paragneisses, (2) eclogites and amphibolites, (3) older orthogneisses and (4) younger orthogneisses (= "Flüelagranitic association"). The complex geologic evolution of this Gondwana polymetamorphic basement was summarised by MAGGETTI and FLISCH (1993). Detailed geochemical studies were undertaken during the last years by the Silvretta working group of the University of Fribourg. In the following, we present a short synthesis of the results.

The paragneisses comprise dominantly biotite-plagioclase gneisses and micaschists. BOLLIN et al. (1993 a, b) interpreted these rocks as cyclic greywacky and pelitic turbidites which were deposited onto a shallow continental shelf. BIINO et al. (1994) reported Sm-Nd model ages of approx. 1.7 Ga for biotite-plagioclase gneisses and micaschists which represents an "average" source age of the detritus. Detrital zircons yielded upper incept ages of 1.5 to 1.9 Ga, which also reflect an "average" Proterozoic source age of the sediments (Grauer and Arnold, 1968).

The metasedimentary rocks are closely associated with eclogites and amphibolites that were derived from N- and T-type tholeitic protoliths (MAGGETTI et al., 1987; MAGGETTI and GALETTI, 1988). According to MAGGETTI et al. (1987) the intercalations of these metabasic rocks with terrigeneous paragneisses represent an original composite layering of igneous basic rocks and sediments. The geotectonic setting could correspond to a divergent oceanic plate boundary that was located close to a continent which provided detritus or, less likely, to a rifting stage of a continental plat-

	Ma 640	Ma 124a	ST 221	Ma 1061	Ma 252	Ma 193	Ma 943	Si 586
La	18.4	5.73	1.742	8.12	7.4	12.03	30.41	6.17
Ce	41.0	15.3	3.929	19.8	16.7	22.34	60.74	15.08
Pr	12 <del></del> 16	1.75	0.544	2.81	_	2.38	6.99	1.72
Nd	24.8	7.93	2.26	11.99	7.8	8.24	27.18	6.42
Sm	6.95	2.34	0.681	2.82	1.90	1.66	5.2	1.84
Eu	2.27	0.72	0.517	0.96	0.71	0.62	1.36	0.15
Gd	7.5	3.39	0.615	2.41	_	1.427	4.66	1.80
Tb	1.29	0.8	0.121	0.259	0.305	0.193	0.66	0.41
Dy	×-	5.89	0.757	2.112	-	1.19	3.57	2.79
Ho	-	1.46	0.186	0.434		0.252	0.79	0.62
Er	-	3.75	0.499	1.125		0.72	2.05	1.77
Tm	_	0.57	0.094	0.156	a <del></del> a	0.107	0.29	0.33
Yb	4.1	3.85	0.617	1.039	1.29	0.789	2.07	2.23
Lu	0.65	0.62	0.096	0.151	0.221	0.139	0.34	0.34

Tab. 2 REE analyses, ppm, Ma 640 and Ma 252 by INA (MAGGETTI et al., 1987), all others by ICP-MS (SARM CRPG, Vandœuvre, France).

form with formation of new oceanic crust. Similar gneiss-amphibolite associations throughout the Alpine domain were interpreted as indicative of either incipient oceanic rifting or back arc environments (e.g. BOUCHARDON et al., 1989; Pa-QUETTE et al., 1989; MESSIGA et al., 1992; MILLER and THOENI, 1995). Another possibility is that the protoliths of the amphibolites and the paragneisses were originally generated in different geotectonic environments. The metabasic rocks may have accreted together with continental-derived sediments (= paragneisses) in a wedge along an active plate margin, and both were subsequently subducted. However, this model can hardly explain the intimate interlayering of both amphibolites and paragneisses.

Major and trace element chemistry of the amphibolite (Ma 640) and the eclogite (Ma 124a) point to MORB compositions (Fig. 2, Tab. 1, 2). The precursor rocks correspond to basic liquids (PEARCE, 1983). Both are, however, enriched in LIL-Elements (Fig. 3). MAGGETTI and GALETTI




Fig. 2 V-Ti/1000 plot (SHERVAIS, 1982; HODDER, 1985). MORB = mid ocean ridge basalt, IAB = island arc basalt.

(1988) explained this by the mobile behaviour of the alkalis during post-magmatic alterations. Lithophile elements can also be enriched in magmas generated from a metasomatised subcontinental lithosphere, or by intracrustal contamination during ascent of basic magmas derived from a "normal" source. PAQUETTE et al. (1989) explained similar LILE enrichments of eclogites from the Alpine External Massifs by 4–5% crustal contamination of tholeitic magmas which were emplaced into thinned continental crust during the initial stages of oceanic rifting.

Already in pre-Ordovician time the metasediments and basic rocks were multiply metamorphosed, first under amphibolite and eclogite and then again under amphibolite facies conditions (MAGGETTI and GALETTI, 1988). After decompression, they were intruded by the protoliths of the older orthogneisses that include ultramafic, gabbroic, dioritic to tonalitic and granitic rocks.




Fig. 3 N(ormal)-MORB normalised plot of metabasites.

Zircon U-Pb ages of these rocks indicate crystallization ages of 610 Ma for a diorite and 530-520 Ma for gabbros, tonalites (SCHALTEGGER et al., this volume) and granitic gneisses (MÜLLER et al., 1995, 1996). The tholeitic basic and the intermediate members of this suite exhibit I-type characteristics (MAGGETTI et al., 1990) whereas the granitic members are of S-type origin (POLLER, 1994 a, b) (Fig. 4). This magmatism has an overall island (volcanic) arc affinity (Fig. 5) (MAGGETTI et




Fig. 4 ASI-Fe<sub>tot</sub> plot to distinguish I- and S-type granitoids after NORMAN et al. (1992). ASI = aluminum saturation index.




Fig. 5 Rb-(Y+Nb) discrimination plot for different tectonic regimes (PEARCE et al., 1984). COLG = syn-collisional granites, ORG = ocean ridge granites, VAG = volcanic arc granites, WPG = within plate granites.

al., 1990; POLLER, 1994 a, b). A detailed discussion of the trace element geochemistry is given by SCHALTEGGER et al. (this volume).

During a second period of magmatism (475–440 Ma) the protoliths of the so-called younger orthogneisses comprising gabbros and granitoids intruded (FLISCH, 1987; LIEBETRAU, 1995). The latter show S-type signatures (Fig. 4) (LIEBETRAU and NÄGLER, 1994; LIEBETRAU, 1995) and are related to collisional regimes (Fig. 5), documenting in the view of LIEBETRAU (1995) differing metasedimentary sources rather than a transition from an oceanic-continent to continent-continent collisional regime.

Nd model ages of 900 to 1000 Ma were reported of basic to intermediate igneous rocks of the older suite (SCHALTEGGER et al., this volume) and of ca. 1700 Ma for older as well a younger granitic gneisses (POLLER, 1994b; POLLER et al., 1994; LIEBETRAU, 1995). These results probably reflect variable mixtures of 500–600 Ma depleted mantle material with crustal components with a mean age of > 1700 Ma. During late Carboniferous the whole Silvretta basement underwent amphibolite facies metamorphism and pervasive deformation (FREI et al., 1995; PROSPERT and BIINO, 1996).

In summary, the metasediments of the Silvretta nappe contain mid-Proterozoic and older components which were deposited in a late Proterozoic oceanic trough. Subduction of this oceanic crust gave rise to two major magmatic cycles in Cambrian and Ordovician times. Cratonization ended with an intensive Variscan amphibolite facies metamorphism. Trace element and REE compositions show the influence of both mantle and crust.

## Analytical procedure

The hand-picked mineral separates (plag, bi, musc, kf, hbl) were treated with hot (120 °C) 2N HCl for 60–80 minutes. After removal of the acid, they were rinsed 2× with destilled water and dried. By this treatment only biotite was markedly corroded. Samples weighing between 25–120 mg were dissolved in 4 ml of a mixture of 6N HCl and 2.3N HF. After complete dissolution the clear solutions were evaporated to dryness and redissolved in 2 ml of 6N HCl. A <sup>205</sup>Pb tracer was added to the solution which then was again evaporated to dryness. The evaporate was dissolved in a 12:1 mixture of 1.5N HBr and 2N of HCl.

Pb was separated using ion-exchange columns filled with 0.5 ml of DOWEX AG1x8 resin (100–200 mesh) and applying an HBr procedure. The purified resin was preconditioned with 2 resin

Tab. 3 Fractionation corrected lead-isotope ratios and lead concentrations of mineral separates.

Sample	$^{206}Pb/^{204}Pb \pm$	$^{207}\text{Pb}/^{204}\text{Pb} \pm$	$^{208}\text{Pb}/^{204}\text{Pb} \pm$	ppm Pb
RB 64 paragneiss (Kno	tengneis), N Scalettapa	ss 789 690/174 510		
plag (V)	18.297 .011	15.661 .012	38.580 .050	8.19
bi (V)	19.032 .013	15.695 .015	39.657 .048	5.20
RB 92 paragneiss (Kno	tengneis), Dischmatal,	Scalettariser 790 000/175	5 400	
bi (V)	18.639 .012	15.673 .015	38.991 .047	4.60
plag (V)	18.266 .009	15.654 .007	38.398 .046	20.80
Ma 640 amphibolite (M	IORB), Sertig, Rhynerh	orn 782 250/177 900		
hbl (V)	20.474 .025	15.706 .029	40.278 .070	0.39
repeat new sample	20.486 .012	15.712 .014	40.291 .048	0.42
plag (V)	18.937 .013	15.597 .014	38.874 .054	1.59
Ma 124a eclogite (MOI	RB), Inntal, Punt Nova	800 900/172 700		
fsp (pV)	18.178 .013	15.610 .016	38.338 .049	4.04
px (pV)	18.298 .013	15.600 .016	38.412 .050	2.95
ST 221 older orthognei	ss, gabbro, Val Sarsura	798 250/176 050		
plag (pV?)	18.157 .013	15.585 .015	37.804 .045	1.98
Ma 1061 older orthogn	eiss, gabbro, Val Tuoi 80	5 480/188 720		
hbl. (V)	18.602 .011	15.576 .014	38.127 .055	0.37
repeat, new sample	18.584 .014	15.567 .016	38.101 .048	0.42
plag(V)	18.005 .011	15.539 .013	37.668 .045	6.75
Ma 252 older orthogne	iss, metadiorite, Val Sag	liains 799 400/186 000		
plag (V)	19.805 .011	15.664 .013	38.007 .045	5.21
hbl (V)	21.385 .015	15.751 .015	38.795 .049	12.03
		lite, Inntal, God Pradits	ch 802 350/176 500	
bi (V)	19.321 .018	15.612 .015	37.835 .037	0.53
repeat	19.314 .034	15.606 .027	37.822 .070	0.70
plag (V)	19.215 .013	15.601 .015	37.742 .048	5.34
Ma 943 older orthogne	iss, augengneiss type M	önchalp, E Tällikopf, Mö	inchalp 789 800/189 500	
bi (V)	137.663 .039	21.939 .006	42.638 .050	7.50
plag (V)	18.881 .026	15.646 .022	42.208 .058	0.34
musc (V)	18.135 .011	15.598 .012	38.642 .045	3.85
Si 586 younger orthogn	eiss, augengneiss type I	Flüela, Radönt 793 220/1	78 760	
musc (V)	19.636 .013	15.700 .016	38.650 .047	2.23
kf (V)	18.356 .011	15.638 .013	38.099 .045	36.89

V = Variscan, pV = pre Variscan

The quoted errors are  $2\sigma$  standard deviations and include the error introduced by the fractionation correction.

volumes of 1N HBr. After the sample was loaded onto the column, the resin was flushed with 3 volumes of 1N HBr and 1 volume of 2N HCl. Pb was then extracted by adding 3 volumes of 7.5N HCl. Pb was purified by repeating the same procedure using 0.25 ml of resin volume and accordingly reduced acid volumes. The analytical Pb blank contribution of 0.2 ng was in all cases < 0.5% of the total Pb and thus was negligible.

Pb was loaded on Re filaments and analyzed with a Finnigan MAT 261 thermal ion mass spec-

trometer. Fractionation effects on the isotope ratios were corrected by a factor

c = 1 + fxDamu

(f =  $1.3 \pm 03$  permil [2 $\sigma$ ]; Damu = difference in atomic mass units between numerator and denominator of the isotope ratio), determined by replicate analysis of the NBS standard reference material 981.

Experience has shown that occasionally lead isotopes from samples may fractionate during mass spectrometric analysis far in excess of the

fractionation observed when analysing the NBS standard lead (KNILL, 1996). This excessive fractionation is not recognisable by a trend of increasing isotopic ratios during analyses. We therefore analysed two minerals of most of the rock samples. Extremely flat or steep slopes in the <sup>207</sup>Pb/<sup>204</sup>Pb–<sup>206</sup>Pb/<sup>204</sup>Pb diagrams would then either indicate that one of the samples fractionated excessively or else that the mineral pair was not in isotopic equilibrium.

#### Results and discussion

Lead isotope analyses are listed in table 3 and the data points are plotted in figure 6. Because of their polymetamorphic nature we did not attempt to correct the isotopic ratios for in situ decayed U and Th and therefore did not determine their U and Th concentrations. The discussion of the present-day lead isotope ratios is therefore focused on the coupled uranogenic <sup>207</sup>Pb/<sup>204</sup>Pb <sup>206</sup>Pb/<sup>204</sup>Pb ratios. In most cases the feldspar lead is comparatively unradiogenic which agrees with the common observation that the U/Pb ratio in feldspars is lower than in other silicates and thus approximates the initial lead composition at the time of feldspar (re-)crystallization. An exception is the feldspar-lead of sample Ma 943 which is more radiogenic than that of the muscovite. According to microscopic evidence most of the minerals (re-)crystallized during the Variscan metamorphism, exceptions are the eclogite (Ma 124a) and the plagioclase of the gabbro (ST 221). The




Fig. 6 (A) <sup>206</sup>Pb/<sup>204</sup>Pb-<sup>207</sup>Pb/<sup>204</sup>Pb-diagram with the growth curve for average Pb of the continental crust according to STACY and KRAMERS (1975) with time-marks at intervals of 100 Ma.

plagioclase of the eclogite forms a symplectitic intergrowth with hornblende which is interpreted as a retrograde reaction product during uplift.

The tie lines (Fig. 6) between coexisting mineral pairs represent isochrons if the minerals contained the same initial common lead at the time of crystallization and if their U-Pb system remained closed in the past. Because of the relatively small spread of the data the error of the slopes of the tielines is high and thus prevents the calculation of meaningful ages. For example, the slope of sample Si 586 (s =  $0.0456 \pm .016$ ) would still be compatible with a Variscan isochron age of 330 Ma (s = 0.0530). The scatter of the slopes in figure 6 can be entirely attributed to the analytical uncertainties stated in table 3. Only the biotite of sample Ma 943 yielded a sufficiently high <sup>206</sup>Pb/<sup>204</sup>Pb ratio of 137.7 to allow in combination with the coexisting plagioclase and muscovite the calculation of a  $^{207}$ Pb/ $^{204}$ Pb age of 330 ± 5 Ma. The age reflects the Variscan metamorphism in the Silvretta nappe (FREI et al., 1995). The highly radiogenic lead is probably caused by the presence of a uranium rich phase in the analyzed sample.

The isotope signatures of the data points cover a spectrum ranging from low <sup>207</sup>Pb/<sup>204</sup>Pb (low μ) ratios relative to <sup>206</sup>Pb/<sup>204</sup>Pb to relatively high <sup>207</sup>Pb/<sup>204</sup>Pb ratios (high μ) (Fig. 6). The lead of the paragneisses yielded the highest μ values (9.9 to 10) which indicates a long residence time of the lead in a relatively uranium rich environment. Such a signature is typical for paragneisses of the East Alpine nappes and of the Southern Alps (KÖPPEL et al., 1993).

Most of the basic to intermediate members of the older orthogneiss suite contain a mantle lead component, i.e. the metagabbro (Ma 1061), the metatonalite (Ma 193) and within the analytical uncertainty also the metadiorite (Ma 252). The volcanic-arc gabbro (ST 221) and the S-type augengneiss (Ma 943) contain lead that is isotopically similar to average crustal lead. Of the two samples from the amphibolite-eclogite group, the lead of the amphibolite (Ma 640) is only little contaminated by crustal lead. The Pb-concentrations of its the major constituents (0.4 to 1.6 ppm, Tab. 3) suggest a somewhat higher whole rock lead concentration relative to N-type MORB (< 1 ppm). The amphibolite is significantly enriched in LILelements (Fig. 3). The eclogite (Ma 124a) consisted prior to retrograde reequilibration of ca. 35% garnet and 55% omphacite plus minor amounts of zoisite and rutile. In contrast to the amphibolite, the eclogite displays an average crustal lead signature. The lead concentration of the pyroxene and retrograde plagioclase (3 to 4 ppm, Tab. 3) also point to higher whole rock lead concentrations than one would expect in N-type MORB. The eclogite is also enriched in LIL-elements (Fig. 3), but less so than the amphibolite.

The data points do not reflect an origin of the lead solely from a depleted mantle. We have outlined in figure 7 the data field of the present-day MORB and the growth-curve for mantle lead according to ZARTMAN and HAINES (1988). The growth curve with  $\mu = 40$  illustrates the lead-evolution of a mantle derived rock with an initial  $^{206}\text{Pb}/^{204}\text{Pb} = 17.43$  and  $^{207}\text{Pb}/^{204}\text{Pb} = 15.37$  and an age of 600 Ma which corresponds to the oldest reported age of igneous rocks of the Silvretta nappe (SCHALTEGGER et al., this volume). Evidently none of the samples contains a lead that was solely derived from a depleted mantle source. This would also hold if we had chosen a more radiogenic initial lead-composition, comparable to the relatively radiogenic lead of some of the present-day MORB. Note that in spite of the high  $\mu$ value the growth curve plots below the mantle curve of ZARTMAN and HAINES (1988). This is a result of their model which accounts for recycling of crustal material with high 207Pb/204Pb into the mantle. To produce the <sup>207</sup>Pb/<sup>204</sup>Pb ratios observed in




Fig. 7  $^{206}\text{Pb}/^{204}\text{Pb}-^{207}\text{Pb}/^{204}\text{Pb}$ -diagram with the growth curve (SK) for average Pb of the continental crust according to STACY and KRAMERS (1975) and time-marks at 200 Ma intervals. MORB designates the field the present-day Pb-isotopic composition of mid-ocean ridge basalts. The mantle growth curve by ZARTMAN and HAINES (1988) indicates the direction along which the MORB field has moved in the past. A growth curve with time-marks at 100 Ma intervals and with  $\mu=40$  is shown which originates at the mantle growth curve 600 Ma ago. EM I, EM II and HIMU denote components of isotopically anomalous mantle lead. The arrows indicate the position of these fields ca. 600 Ma ago. Symbols as in figure 6.

the samples would require that mantle derived magmas with  $\mu$  values between 20 and 25 were generated 2000 Ma ago. There is no evidence, however, of such high protolith ages of the I-type meta-granitoids or of the amphibolites. Moreover, the  $^{206}Pb/^{204}Pb$  ratios would in this case vary between about 21.5 and 23.3.

Excluding a crustal lead contribution during magma genesis, ascent and cooling, the lead isotope pattern of the basic to intermediate members of the older orthogneisses thus suggests an origin from an anomalous mantle. An anomalously enriched mantle may result from metasomatism by dewatering reactions of subducted sediments (EM II, ZINDLER and HART, 1986) such as the paragneisses (Fig. 7). A further possibility would be a derivation from an EM I type mantle which was contaminated by metasomatism originating, for example, from subducted old lower crust which normally has low U/Pb and high Th/Pb ratios. Although such an origin cannot be ruled out, it appears unlikely because the uranogenic lead is not obviously retarded and the thorogenic ratios are not abnormally high. A contribution from a HIMU mantle source is unlikely in view of the relatively low <sup>206</sup>Pb/<sup>204</sup>Pb but high <sup>207</sup>Pb/<sup>204</sup>Pb ratios of most of the data (Fig. 7).

Alternatively, if the source region of the magma had approximately the isotopic charcteristics seen in present-day MORB, a contribution of crustal lead from metasediments during magma genesis and emplacement would also explain the data. Assuming that the lead contaminant derived from sediments (RB 64 and RB 92), then their Pbcontribution to samples Ma 640, Ma 1061 and Ma 252 may be estimated from figure 6 to amount to 25% to 50% depending on the Pb-isotopic composition of the mantle lead. Assuming the same mixing components, the meta-tonalite (Ma 193) would then be somewhat less, and the meta-gabbro (ST 221) and the eclogite (Ma 124a) more contaminated (50-80%). Due to the contrasting Pb concentration of mantle derived magmas (approx. 1 ppm) and rocks of the continental crust (> 10 ppm) contamination by assimilation of sediments would have amounted to approximately 2% to 5% which is in reasonable agreement with the estimate of a maximum contribution of 10% on the basis of Sm/Nd isotopic systematics (SCHALTEGGER et al., this volume). A further possibility is contamination by submarine hydrothermal alteration in the presence of sediments. It certainly could have affected the basalts which were the protoliths of the amphibolites. Also the crustal lead isotope signature of the eclogite (Ma 124a) may be the result of a hydrothermal sea-floor alteration. The eclogites occur as alternating, 5 to

Sample	V-Ti/1000	Zr/Y–Zr	Ti–Zr	Cr–Y	Rb-(Y+Nb)	Fe(tot)- ASI	Pb-Isotope signature
Eclogite Ma 124a	MORB	MORB(?)	MORB	MORB	-	_	average crust
Amphibolite Ma 640	MORB	MORB	MORB(?)	MORB(?)	-	_	mantle-crust
Gabbro Ma 1061	IAB	-?	IAB	IAB	_	I-type	mantle-crust
Gabbro ST 221	IAB	-?	IAB	-?	_	I-type	crust-mantle
Biotite bearing							
metatonalite Ma 193	_	_	_	_	VAG	I-type	mantle-crust
Metadiorite Ma 252*	_	_	_	_	VAG	I-Type	mantle-crust
Augengneiss (type						• •	
Mönchalp) Ma 943	<del>-</del>	_	_	_	VAG	S-type	average crust
Augengneiss (type						. 1	
Flüela) Si 586	-	_	_	_	COLG	S-Type	crust

Tab. 4 Summary of geochemical characteristics.

100 cm thick layers of differing eclogitic modal composition within a 500 m thick complex of metabasites which is surrounded by metasediments. Thus, the protoliths of the eclogites were most likely oceanic basalts.

Both augengneisses, the older (Ma 943) and the younger orthogneiss (Si 586) are, based on the SiO<sub>2</sub>-ASI discrimination diagram, S-type granites. Accordingly, the uranogenic lead of the older orthogneiss rocks is of average crustal composition and is thus typical of lead of mature arcs, and the lead of the younger orthogneiss exhibits a slightly more pronounced crustal signature.

### **Conclusions**

The lead isotope data support in most cases the interpretation of the geochemical signatures (Tab. 4). The isotopic signature of the amphibolite (Ma 640) indicates the presence of a mantle component in agreement with its geochemical characteristics that suggest a MORB origin. The geochemical signatures of the four I-type intrusives indicate an island arc basalt and volcanic arc setting which agrees with the presence of a mantle derived lead component in these samples. Based on the aluminum saturation index the VAG-type metagranite (Ma 943) and a syncollisional metagranite (Si 586) are S-type granites. Their isotope signature indicates a crustal origin of the lead and thus supports an S-type origin.

In spite of the polymetamorphic nature of the meta-igneous rocks, their geochemical and lead isotope signatures are in the majority of cases compatible. The only exception to this is the eclogite (Ma 124a) which contains an average crustal lead but the geochemical signatures are indicative

of a MORB origin. However, from its intimate association with metasediments one may plausibly conclude that hydrothermal sea floor alteration was responsible for the crustal signature of the lead.

The lead isotope data support the idea that new crust was generated first by mantle differentiation in an island arc situation and subsequently with increasing participation of crustally derived metasediments which contained early Proterozoic detritus. The lead isotope data do not indicate the presence of remnants of an old, for example early Proterozoic or even Archean crust.

### Acknowledgements

We thank I. Ivanov-Bucher and W. Wittwer for preparing the mineral separates and R. Frei for reviewing the manuscript; his critical comments were of great help in preparing the final version of this paper. We are grateful to C. Prospert who contributed figure 1.

#### References

BIINO, G., MEISEL, T.C., NÄGLER, T.F. and KRAMERS, J.-D. (1994): Whole rock chemistry and isotope chemistry of metasediments in the Silvretta nappe and the early crustal history of the alpine basement. Abstracts 174. Annual meeting Swiss Academy of Science, 4–8 October 1994, 38.

Bollin, R. (1993a): Geochemie und seltene Erden (SE) der Paragneise (Biotit- und Feldspatknotengneise) der Silvrettadecke (Kanton Graubünden/Schweiz). Schweiz. Mineral. Petrogr. Mitt. 73/1, 130.

Bollin, R. (1993b): Metamorphe Entwicklung und Ursprung der Paragneise (Biotitplagioklasgneise, Plagioklasknotengneise) der Silvretta. Unpublished Ph. D. Thesis, University of Freiburg, Switzerland.

BOUCHARDON, J.L., SANTALLIER, D., BRIAND, B., MÉNOT,

<sup>\*</sup> The discrimination diagrams are not strictly applicable to this rock as its low quartz content originates probably from metamorphic reactions.

R.P. and PIBAULE, M. (1989): Eclogites in the French Palaeozoic orogen: Geodynamic significance. Tectonophysics 169, 317–332.

CUMMING, G.L., KÖPPEL, V. and FERRARIO, A. (1987): A lead isotope study of the northeastern Ivrea zone and the adjoining Ceneri zone (N-Italy): evidence for a contaminated subcontinental mantle. Contrib. Mineral. Petrol. 97, 19–30.

CURTI, E. (1987): Lead and oxygen isotope evidence for the origin of the Monte Rosa gold lode deposits (Western Alps, Italy): A comparison with Archean

lode deposits. Econ. Geol. 82, 2115–2140.

DOE, B.R. and ZARTMAN, R.E. (1979): Plumbotectonics, the Phanerozoic. In: BARNES, H.L. (ed.): Geochemistry of Hydrothermal Ore Deposits, 2nd ed. J. Wiley

& Sons, New York, 22–70.

FLISCH, M. (1987): Teil 1: Geologische, petrographische und isotopengeologische Untersuchungen an und isotopengeologische Gesteinen des Silvrettakristallins. Teil 2: Die Hebungsgeschichte der oberostalpinen Silvretta-Decke seit der mittleren Kreide. Unpublished Ph. D Thesis, University of Bern, Switzerland.

FLOYD, P.A. and WINCHESTER, J.A. (1978): Identification and discrimination of altered and metamorphosed volcanic rocks using immobile elements. Chem.

Geol. 21, 291-306.

Frei, R., Biino, G.G. and Prospert, C. (1995): Dating a Variscan pressure-temperature loop with staurolite.

Geology 23, 1095-1098.

GRAUERT, B. and ARNOLD, A. (1968): Deutung diskordanter Zirkonalter der Silvrettadecke und des Gotthardmassivs (Schweizer Alpen). Contrib. Mineral. Petrogr. 20, 34–56.

HODDER. A.P.W. (1985): Depth of origin of basalts inferred from Ti/V ratios and a comparison with the K<sub>2</sub>O-depth relationship for island arc volcanics.

Chem. Geol. 48, 3-16.

KNILL, M. (1996): The Pb-Zn-As-Tl-Ba-deposit at Lengenbach, Binn Valley, Switzerland. Petrogenesis based on combined geochemical and isotopical (U, Pb, Rb, Sr, S, O, C) investigations. Beiträge zur Geologie der Schweiz, Geotechnische Serie, Lief. 90, pp. 74.

KÖPPEL, V., NEUBAUER, F. and SCHROLL, E. (1993): Prealpidic ore deposits in the Central, Eastern and Southern Alps. In: VON RAUMER and NEUBAUER (eds): The pre-Mesozoic Geology of the Alps, Springer Verlag, Heidelberg, 145–162.

LIEBETRAU, V. (1995): Petrographie, Geochemie und Datierung der Orthogneise der sog. "Flüelagranitischen Assoziation" (Silvrettakristallin, Graubünden). Unpublished Ph. D. Thesis, University of

Freiburg, Switzerland. LIEBETRAU, V. and NÄGLER, T. (1994): Geochronologische und geochemische Diskussion der sogenannten "flüelagranitischen Assoziation" des Silvrettakristallins (Graubünden/Schweiz). Schweiz. Mineral.

Petrogr. Mitt. 74, 265–268. MAGGETTI, M. and GALETTI, G. (1988): Evolution of the Silvretta eclogites: magmatic and metamorphic events. Schweiz. Mineral. Petrogr. Mitt. 68, 467–484.

- MAGGETTI, M., GALETTI, G. and STOSCH H.-G. (1987): Eclogites from the Silvretta nappe (Switzerland): geochemical constraints on the nature and geotectonic setting of their protoliths. Chem. Geol. 64, 319-334.
- MAGGETTI, M., GALETTI, G. and STOSCH, H.-G. (1990): Geochemische Argumente zur Genese der "Alteren Orthogneise" der Silvretta. Schweiz. Mineral. Petrogr. Mitt. 70, 103–107.
- MAGGETTI, M. and FLISCH, M. (1993): Evolution of the

Silvretta Nappe. In: von RAUMER, J.F. and NEUBAUER, F. (eds): Pre-Mesozoic Geology in the Alps. Springer, Berlin, 469-484.

MESSIGA, B., TRIBUZIO, R. and SCAMBELLURI, M. (1992): Mafic eclogites from the Valosio crystalline massif (Ligurian Alps, Italy). Schweiz. Mineral. Petrogr. Mitt. 72, 365–377.

MILLER, C. and THOENI, M. (1995): Origin of eclogites from the Austroalpine Ötztal basement (Tirol, Austria): Geochemistry and Sm-Nd vs Rb-Sr isotope systematics. Chem. Geol. 122, 199-225.

MÜLLER, B., KLÖTZLI, U. and FLISCH, M. (1995): U-Pb and Pb-Pb zircon dating of the older orthogneiss suite in the Silvretta nappe, eastern Alps: Cadomian magmatism in the upper Austro-Alpine realm. Geol. Rundschau 84, 457–465.

MÜLLER, B., KLÖTZLI, U., SCHALTEGGER, U. and FLISCH, M. (1996): Early Cambrian oceanic plagiogranite in the Silvretta nappe, eastern Alps: Geochemical, zircon U-Pb and Rb-Sr data from garnet-hornblendeplagioclase gneisses. Geol. Rundschau 25, 822-831.

NORMAN, M.D., LEEMAN, W.P. and MERTZMAN, S.A. (1992): Granites and rhyolites from the northwestern USA: Temporal variation in magmatic processes and relations to tectonic setting. Transactions of the Royal Society of Edinburgh, Earth Sciences, 83, 71-81.

PAQUETTE, J.L., MÉNOT, R.P. and PEUCOT, J.J. (1989): REE, Sm-Nd and U-Pb zircon study of eclogites from Alpine External Massifs (Western Alps): Evidence for crustal contamination. Earth Planet. Sci. Lett. 96, 181-198.

PEARCE, J.A. (1983): A "users" guide to basalt discrimination diagrams. Unpubl. report, The Open Univer-

sity, Milton Keynes, 37 pp.

PERETTI, A and KÖPPEL, V. (1986): Geochemical and lead isotope evidence for a mid-oceanic ridge type mineralization within a polymetamorphic ophiolite complex (Monte del Forno, North Italy/Switzerland). Earth and Planet. Sc. Lett. 80, 252–264.

POLLER, U. (1994a): Der Mönchalpgneis der Silvrettadecke (Graubünden): Geochemie und Sm-Nd-Modellalter. Schweiz. Mineral. Petrogr. Mitt. 74,

269-272

POLLER, U. (1994b): Petrographie, Geochemie und Datierung der Augengneise Typ Mönchalp (Ältere Orthogneise) des Silvrettakristallins, Graubünden-Schweiz. Unpublished Ph. D. Thesis, University of

Freiburg, Switzerland.

POLLER, U., LIEBETRAU, V., NÄGLER, T. and MAGGETTI, M. (1994): A study of crustal evolution and Tectonic setting applying Rb/Sr, Sm/Nd and trace element analyses (Silvretta, Switzerland). Abstracts 8th Int. Conference on Geochronology, Cosmochronology and Isotope Geology, Berkeley, US GS Circular

PROSPERT, C. and BIINO, G.G. (1996): The Al<sub>2</sub>SiO<sub>5</sub> veins and segregations of the Silvretta thrust sheet: Structural and metamorphic implications for a Carboniferous history. Mitt. Österr. Min. Ges. 141, 180–181.

SCHALTEGGER, U., NÄGLER, TH.F., CORFU, F., MAGGETTI, M., GALETTI, G. and STOSCH, H.-G. (1997): A Cambrian island arc system in the Silvretta nappe: constraints from geochemistry and geochronology. Schweiz. Mineral. Petrogr. Mitt. 77, 337–350. Shervais, J.W. (1982): Ti–V plots and the petrogenesis of

modern and ophiolitic lavas. Earth Planet. Sci. Lett.

59, 101-118.

STACEY, J.S. and KRAMERS, J.D. (1975): Approximation of terrestrial lead isotope evolution by a two-stage model. Earth and Planet. Sci. Lett. 26, 207-221.

- Taylor, S.R. and McLennan, S.M. (1985): The continental crust: its composition and evolution. Blackwell, Oxford.
- WINCHESTER, J.A. and FLOYD, P.A. (1977): Geochemical discrimination of different magma series and their differentiation products using immobile elements. Chem. Geol. 20, 325–343.
- ZARTMAN, R.E. and HAINES, S.M. (1988): The plumbotectonic model for lead isotopic systematics among
- major terrestrial reservoirs. A case for bi-directional transport. Geochim Cosmochim Acta 52, 1327–1340. ZINDLER, A. and HART, S. (1986): Chemical geodynamics. Ann. Rev. Earth Planet. Sci 14, 493–571.

Manuscript received February 5, 1997; revised manuscript accepted August 25, 1997.