

**Zeitschrift:** Schweizerische mineralogische und petrographische Mitteilungen =  
Bulletin suisse de minéralogie et pétrographie

**Band:** 79 (1999)

**Heft:** 3

**Artikel:** The Gemicic S-type granites in southeastern Slovakia : Late  
Palaeozoic or Alpine intrusions? Evidence from electron-microprobe  
dating of monazite

**Autor:** Finger, Fritz / Broska, Igor

**DOI:** <https://doi.org/10.5169/seals-60218>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 13.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

SHORT COMMUNICATION

## The Gemicic S-type granites in southeastern Slovakia: Late Palaeozoic or Alpine intrusions? Evidence from electron-microprobe dating of monazite

by *Fritz Finger<sup>1</sup>* and *Igor Broska<sup>2</sup>*

### Abstract

The method of total-Pb monazite dating by means of the electron microprobe has been applied to broadly constrain the age of granite samples from the Gemicic tectonic unit in southeastern Slovakia. The aim was to determine whether these granites intruded in the late Palaeozoic or during the Alpine orogeny in the Cretaceous. Both views are currently held in the geological literature. Monazites from three different localities consistently yielded Permian Th-U-Pb model ages, with weighted averages for single samples at 272±11 Ma, 276±13 Ma and 273±13 Ma. The results thus suggest that the S-type granites of the Gemicicum formed essentially in mid-Permian times.

**Keywords:** Carpathians, Slovakia, Gemicicum, granites, monazite, geochronology.

### Introduction

The Gemicicum represents the southernmost basement-bearing tectonic unit of the Inner (BIELY et al., 1996), or Central (MAHEL', 1986; PLAŠIENKA et al., 1997) Western Carpathians. During the Alpine orogeny, it was thrust onto the Veporicum (Fig.1). The Gemicicum itself includes several internal thrust sheets. It is uncertain whether these formed during Alpine or pre-Alpine deformational events. According to GRECULA (1982), a phase of intra-Gemicic nappe stacking may have occurred in the Permian. Lithologically, the Gemicicum consists mainly of poly-metamorphic Lower Palaeozoic strata, with basic to intermediate volcano-sedimentary formations in the north (Klatov and Rakovec groups) and volcanogenic flysch-type metasediments in the south (Gelnica group). Variscan regional metamorphism reached amphibolite facies conditions in the Klatov group and greenschist facies grade in the Rakovec and Gelnica groups (KRIST et al., 1992; FARYAD, 1997). Alpine metamorphism was generally at lower greenschist facies grade.

Tin-bearing S-type granites intruded into the folded Variscan basement. Although only small

surface exposures are present (e.g. PETRIK and KOHÚT, 1997), geophysical data and boreholes suggest that large masses of these granites are hidden below the metavolcanics and metasediments (VOZÁR et al., 1996). According to field evidence, at least some of the granites were included in the internal Gemicic nappe system, and their intrusion ages thus constitute important time limits for this thrust tectonics (GRECULA, 1982, 1995).

In the past, a Permian age has been assumed for the Gemicic granites, relying on Rb-Sr whole-rock isochrons (KOVACH et al., 1986, CAMBEL et al., 1989). However, a recent deep seismic profile has revealed structures which are interpreted in favour of an important Alpine (Cretaceous?) plutonic event in the Gemicic unit (VOZÁR et al., 1996). Transparent and (due to their small amplitude) indistinct seismic zones in the upper crust were regarded as (syn-collisional) granitoid bodies intruded into the Alpine structures. This has raised some doubts concerning the reliability of the Permian Rb-Sr whole-rock ages obtained from the surface exposures of Gemicic granites (VOZÁROVA and VOZÁR, 1996), all the more so as Rb-Sr and K-Ar muscovite dating provided mostly Alpine values for these rocks (KANTOR and

<sup>1</sup> Institut für Mineralogie der Universität Salzburg, Hellbrunnerstrasse 34, A-5020 Salzburg, Austria.  
<Friedrich.Finger@sbg.ac.at>

<sup>2</sup> Geological Institute of the Slovak Academy of Sciences, Dubravská Cesta 9, 842 26 Bratislava, Slovakia.

RYBÁR, 1979). The Permian ages still await confirmation through dating of high-T minerals like zircon or monazite.

In this paper, we present chemical Th–U–Pb model ages of monazites from three samples of Gemicic granites. Two samples were obtained from the northwestern part of the Gemicicum (Hnilec body), whilst one comes from the southwestern corner of the unit (Betliar body). This latter granite body has been considered as the most potential candidate for an Alpine-age intrusion from geological reasoning, considering the seismic data (VOZÁR *et al.*, 1996; VOZÁROVA and VOZÁR, 1996).

### The dated samples

Granite samples GZ-1 and GZ-3 are medium-grained, whilst GZ-15 is coarser grained with phryic K-feldspar (for sample localities, see Fig.1). Apart from large amounts of perthitic K-feldspar, oligoclase and quartz, the rocks contain some muscovite and rare biotite. Accessory phases are

zircon, monazite, apatite, xenotime, rutile, garnet and tourmaline. All three samples show a severe mylonitic low-T overprint and are in the strict sense metagranites, a feature which holds true for the Gemicic granites in general (PETRIK and KOHÚT, 1997). Geochemically, all three samples can be described as S-type granites. As is typical for the Gemicic granite suite (e.g. BROSKA and UHER, 1998), they are high in  $\text{SiO}_2$  (~70%), strongly peraluminous ( $\text{A/CNK} \sim 1.3$ ) and rich in phosphorus.

### Dating results

Several recent studies (e.g. COCHERIE *et al.*, 1998 and further references therein) have shown that chemical Th–U–Pb dating of monazite by means of the electron microprobe is a useful geochronological tool, as it provides moderately precise monazite crystallization ages. The theoretical basis of the method, its advantages and major error risks can be taken from the work of MONTEL *et al.* (1996). The analytical procedure used in the pre-




Fig. 1 Geological sketch map of the Western Carpathians showing the major basement massifs and the sample localities of this study (GZ-1: Hnilec, cliff 780 m above sea level, 800 m NE from elevation point Peklisko; GZ-3: Hnilec, outcrop 1205 m above sea level, 220 m NE from elevation point Surovec; GZ-15: Betliar, cliff 3250 m SW from elevation point Volovec II). Inset shows the position of the map area (white star) in the Alpine-Carpathian chain.

Tab. 1 Th, U, Pb contents (wt. % elements), Th\* values and model ages of the analysed monazites from samples GZ-1, GZ-3 and GZ-15. Model ages and  $2\sigma$  errors were calculated after the method of MONTEL et al. (1996), Th\* values have been recast with these model ages using the equation given in SUZUKI et al. (1991). For each sample, a weighted average age (95 % c.l.) is given.

Grain	Th	U	Pb	Th*	Age
<b>SAMPLE GZ-1</b>					
m1	10.618	0.762	0.149	13.082	<b>256 ± 24</b>
m2	11.489	1.213	0.192	15.419	<b>279 ± 20</b>
m3	13.340	1.371	0.219	17.779	<b>276 ± 18</b> <b>272 ± 11</b>
<b>SAMPLE GZ-3</b>					
m1	2.714	0.057	0.029	2.898	<b>228 ± 108</b>
m2	12.550	1.416	0.207	17.134	<b>271 ± 18</b>
m3	10.732	0.980	0.179	13.910	<b>289 ± 23</b>
m4	2.583	0.048	0.038	2.740	<b>307 ± 114</b>
m5	3.243	0.061	0.042	3.441	<b>272 ± 91</b>
m6	3.298	0.066	0.045	3.510	<b>285 ± 90</b>
m7	3.556	0.036	0.038	3.672	<b>234 ± 86</b> <b>276 ± 13</b>
<b>SAMPLE GZ-15</b>					
m1	6.377	0.123	0.075	6.775	<b>249 ± 46</b>
m2	9.166	0.664	0.145	11.317	<b>287 ± 28</b>
m3	9.203	0.683	0.133	11.414	<b>262 ± 28</b>
m4	10.145	1.299	0.170	14.349	<b>266 ± 22</b>
m5	17.049	1.462	0.273	21.786	<b>281 ± 14</b>
m6	9.613	0.421	0.124	10.974	<b>254 ± 29</b> <b>273 ± 13</b>

sent study follows the working routine in the Salzburg laboratory, described in FINGER and HELMY (1998).

Monazites were located in normal polished thin sections by means of BSE imaging. All three samples contained several monazite grains per thin section, the largest of which were chosen for analysis. The analyses were carried out with a defocused beam (5  $\mu\text{m}$ ), which was generally placed into the grain centres. This was done to avoid potential lead loss effects which may occur at the rim zones of monazite (SUZUKI et al., 1994). The risk of picking up inherited lead in the core zones was considered minor, due to the fractionated nature of the granites and the relatively high proposed magma temperatures of around 750–800 °C (PETRIK and KOHÚT, 1997). Additionally, BSE imaging provided no evidence of inherited cores. The Th, U, Pb concentrations and calculated model ages with errors ( $2\sigma$ ) are given in Tab. 1. Using an accelerating voltage of 15 kV, a probe current of 250 nA and counting times of 200 s for Pb (M $\alpha$ ), 50 s for U (M $\alpha$ ) and 30 s for Th (M $\alpha$ ), the 2-sigma

errors and detection limits resulting from the counting statistics of the microprobe were typically 0.011 wt. % for Pb, 0.03 wt. % for U and 0.04 wt. % for Th.

In figure 2 the results are graphically presented in a Th\* vs Pb diagram, together with data for a monazite age standard. This standard was used to independently control the analytical accuracy of the dating procedure (FINGER and HELMY, 1998). In the Th\* vs Pb diagram, monazites of one age should plot along a straight line through zero, given that common lead contents and lead loss were unimportant. The slope of this line is a direct function of the age. The Th\* parameter has been calculated according to SUZUKI et al. (1991). It includes the measured Th plus a certain amount of theoretical Th, that would have produced the same amount of lead as the measured U (at the model age).

The thin section of sample GZ-1 contained only three monazites of sufficient size. These had high brabantite components ( $\text{CaTh}[\text{PO}_4]_2$ ), and thus gave well defined model ages between  $256 \pm 24$  Ma and  $279 \pm 20$  Ma (weighted average age:  $272 \pm 11$  Ma). In sample GZ-3, seven grains were analysed. Most had rather low Th contents and, consequently, low contents of radiogenic lead and large errors in the age calculation (Tab. 1). Two high-Th grains provided ages of  $271 \pm 18$  Ma and  $289 \pm 23$  Ma, respectively. The weighted average




Fig. 2 Total Pb vs. Th\* isochron diagram after SUZUKI et al. (1991) using the values given in Tab. 1. Also shown are data from laboratory standard F-5 obtained during the same analytical session. The F-5 monazites were taken from a fraction, which was dated by U-Pb isotope dilution – mass spectrometry techniques at a concordant age of  $341 \pm 2$  Ma (FRIEDL, 1997). The recommended age could be perfectly reproduced (weighted average:  $342 \pm 7$  Ma; MSWD 0.54). The time scale shown is based on the position of zero-intercept isochrons. The two drawn isochrons refer to the calculated weighted average of all sample and standard data points, respectively (95 % c.l.). Size of symbols corresponds approximately to the  $\pm 1\sigma$  error of the Pb analysis.

age of all seven analyses is  $276 \pm 13$  Ma. Six grains were analysed in *sample GZ-15*. These had a Th\* variation from ca. 7 to 22 wt% and gave consistent Permian model ages, clustering around a weighted mean of  $273 \pm 13$  Ma. The linear Th\*-Pb covariation of all data points and their trend towards zero (Fig. 2) argues against substantial lead loss or inheritance effects, and also against the presence of large amounts of common lead. Therefore, we assume that the model ages correspond more or less to the magmatic crystallization ages of the monazites.

### Conclusions

The obtained monazite model ages lend support to the classic concept that the tin-bearing Gemicic S-type granite suite has a Permian formation age (see e.g. CAMBEL et al., 1989). Also, the Betliar granite in the south is obviously a Permian intrusion. Our data suggest that these Gemicic S-type granites formed at ca. 270 Ma, which means at a time when large parts of the Palaeozoic basement in the Carpathians and Alps underwent a phase of rifting (VOZÁR, 1997; NEUBAUER et al., 1999). High heat flow at this time apparently caused melting of lower-crustal metasedimentary sources. Later, the granites were incorporated into the nappe system of the Gemicic unit. However, the new age data for the granites provide no information as to whether this happened in the mid/late Permian or during the Alpine orogeny. Furthermore, it cannot be ruled out, of course, that younger granite bodies of Cretaceous age are present in the upper crust as well and are not currently exposed at the surface.

### Acknowledgements

François Bussy and Urs Schaltegger are thanked for their helpful reviews. Gudrun Riegler has drawn the figures, Robin Cooke revised the English. Many thanks to both. The work was funded by the Austrian National Bank, grant 7163.

### References

BIELY, A., BEZÁK, V., ELEČKO, M., GROSS, P., KALIČIAK, M., KONEČNY, V., LEXA, J., MELLO, J., NEMČOK, J., POTFAJ, M., RAKÚS, M., VASS, D., VOZÁR, J. and VOZÁROVÁ, A. (1996): Explanation to geological map of Slovakia 1 : 500 000. Dionyz Štúr Publishers, Bratislava, 76 p.

BROSKA, I. and UHER, P. (1998): Brown and blue schorl from the Spiš-Gemer granite, Slovakia: composition and genetic relations. *J. Czech Geol. Soc.* 43, 9–16.

CAMBEL, B., BAGDASARJAN, G.P., VESELSKY, J. and GUKASJAN, R.CH. (1989): Rb-Sr geochronology of the leucocratic granitoid rocks from the Spišsko-gemerské rudoohorie Mts. and Veporicum. *Geol. Zbor. Geol. carpath.*, 40, 323–332.

COCHERIE, A., LEGENDRE, O., PEUCAT, J.J. and KOUAME-LAN, A.N. (1998): Geochronology of polygenetic monazites constrained by in situ electron microprobe Th-U-total lead determination: Implications for lead behaviour in monazite. *Geochim. Cosmochim. Acta.* 62, 2475–2497.

FARYAD, S.W. (1997): Metamorphic petrology of the Early Paleozoic low-grade rocks in the Gemicic. In: GRECULA, P., HOVORKA D., PUTIŠ, M. (eds): Geological evolution of the Western Carpathians. *Mineralia Slovaca-Monography*, 235–252.

FINGER, F. and HELMY, H.M. (1998): Composition and total-Pb model ages of monazite from high-grade paragneisses in the Abu Swayel area, southern Eastern Desert, Egypt. *Mineral. Petrol.*, 62, 269–289.

FRIEDL, G. (1997): U/Pb-Datierungen an Zirkonen und Monaziten aus Gesteinen vom österreichischen Anteil der Böhmischem Masse. Ph.D Thesis, Univ Salzburg, 242p.

GRECULA, P. (1982): Gemicic – Segment of the Paleotethyan riftogenous basin. Alfa, Bratislava, 263 p.

GRECULA, P. (Ed.) (1995): Mineral deposits of the Slovak Ore Mountains. Vol. 1. Geocomplex Press, Košice, 831 p.

KANTOR, J. and RYBÁR, M. (1979): Radiometric ages and polyphasic character of Gemicic granites. *Geol. Zbor. Geol. carpath.*, 30/4, 433–447.

KOVÁČ, A., SVINGOR, E. and GRECULA, P. (1986): Rb-Sr isotopic ages of granitoids from the Spišsko-Gemerské rudoohorie Mts. Western. *Mineralia slov.*, 18, 1–14.

KRIST, E., KORIKOVSKIJ, S. P., PUTIŠ, M., JANÁK, M. and FARYAD, S.W. (1992): Geology and petrology of metamorphic rocks of the Western Carpathian crystalline complexes. Comenius Univ. Press, Bratislava, 324 p.

MAHEL, M. (1986): Geological structure of the Czechoslovak Carpathian. Paleoalpine units. Veda publishers Bratislava, 503 p.

MONTEL, J.M., FORET, S., VESCHAMBRE, M., NICOLLET, CH. and PROVOST A. (1996): A fast, reliable, inexpensive in-situ dating technique: Electron microprobe ages on monazite. *Chem. Geol.*, 131, 37–53.

NEUBAUER, F., HOINKES, G., SASSI, F.S., HANDLER, R., HÖCK, V., KOLLER, F., FRANK, W. (1999): Pre-Alpine metamorphism in the Eastern Alps. *Schweiz. Mineral. Petrogr. Mitt.*, 79, 41–62.

PETRÍK, I. and KOHÚT, M. (1997): The evolution of granitoid magmatism during the Hercynian orogen in the Western Carpathians. In: GRECULA, P., HOVORKA D., PUTIŠ, M. (eds): Geological evolution of the Western Carpathians. *Mineralia Slovaca – Monography*, 235–252.

PLAŠIENKA, D., GRECULA, P., PUTIŠ, M., HOVORKA, D. and KOVÁČ, M. (1997): Evolution and structure of the Western Carpathians: an overview. In: GRECULA, P., HOVORKA, D., PUTIŠ, M. (eds.): Geological evolution of the Western Carpathians. *Mineralia slovaca – Monography*, Bratislava, p. 1–24.

SUZUKI, K., ADACHI, M. and TANAKA, T. (1991): Middle Precambrian provenance of Jurassic sandstone in the Mino Terrane, central Japan: Th-U-total Pb evidence from an electron microprobe monazite study. *Sedimentary Geol.* 75, 141–147.

SUZUKI, K., ADACHI, M. and KAJIZUKA, I. (1994): Electron microprobe observations of Pb diffusion in

metamorphosed detrital monazites. *Earth Planet. Sci. Lett.*, 128, 391–405.

VOZÁROVÁ, A. and VOZÁR, J. (1996): Terranes of the West Carpathians – North Pannonian domain. *Slovak Geol. Magazine* 1, 65–85.

VOZÁR, J., TOMEK, Č., VOZÁROVÁ, A., MELLO, J. and IVANIČKA, J. (1996): Seismic section G-1. *Geol. Práce*, Spr. 101, 32–34.

VOZÁR, J. (1997): Rift-related volcanics in the Permian of the Western Carpathians. In: GRECULA, P., HOVORKA, D., PUTIŠ, M. (eds): *Geological evolution of the Western Carpathians. Mineralia Slovaca – Monography*, 225–234.

Manuscript received March 12, 1999; revision accepted August 30, 1999.