Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	80 (2000)
Heft:	2
Artikel:	Chemical compositon of piemontites and reaction relations of piemontite and spessartine in piemontite-quartz schists of central Shikoku, Sanbagawa metomorphic belt, Japan
Autor:	Izadyar, Javad
DOI:	https://doi.org/10.5169/seals-60961

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Chemical composition of piemontites and reaction relations of piemontite and spessartine in piemontite-quartz schists of central Shikoku, Sanbagawa metamorphic belt, Japan

by Javad Izadyar^{1,2}

Abstract

Chemical composition of piemontite and reaction relations between spessartine and piemontite from the Sanbagawa piemontite-quartz schists in central Shikoku have been studied. The Sanbagawa metamorphic belt was formed during a regional Cretaceous intermediate high-pressure type of metamorphism present throughout Southwest Japan. This study clarifies that the Sanbagawa piemontite-quartz schists can be divided into talc-bearing and talc-free types. Piemontite in the talc-bearing type (Asemi-gawa area) exhibits two zones, whereby average chemical compositions of the core and rim based on the piemontite (Pi), pistacite (Ps) and clinozoisite (Czo) end-members are: $X_{Pi} = 0.26$, $X_{Ps} = 0.09$, $X_{Czo} = 0.65$ and $X_{Pi} = 0.21$, $X_{Ps} = 0.19$, $X_{Czo} = 0.60$, respectively. In the Besshi area, average chemical compositions of the core and rim are: $X_{Pi} = 0.17$, $X_{Ps} = 0.13$, $X_{Czo} = 0.70$ and $X_{Pi} = 0.18$, $X_{Ps} = 0.16$, $X_{Czo} = 0.66$, respectively. However, sometimes Fe³⁺ and Mn³⁺ decrease but Al increases from core to rim. Piemontite in the talc-free type exhibits complex zoning patterns. In the Asemi-gawa area, three zones can be seen: a core: ($X_{Pi} = 0.18$, $X_{Ps} = 0.15$, $X_{Czo} = 0.67$), a mantle: ($X_{Pi} = 0.04$, $X_{Ps} = 0.21$, $X_{Czo} = 0.74$), and a rim: ($X_{Pi} = 0.19$, $X_{Czo} = 0.71$). In the Besshi area, two types of zoning can be identified; in one type a decrease of Mn and Fe and an increase of Al, and in another type Mn increases whilst Al and Fe decrease (from core to rim).

The Sanbagawa belt is one of the most extensively studied metamorphic complexes, and thus, gives a good opportunity for finding a relation between chemical variation of piemontite and changing P,T, fO_2 and bulk composition. In general, the piemontite from the Asemi-gawa region contains higher piemontite and lower pistacite and clinozoisite end-members than that from the Besshi area. By considering the difference of metamorphic grade, similarity of bulk composition and constant oxygen fugacity between two areas, piemontite with higher epidote end-member appears to be stable at the higher grade. Based on textural and chemical evidences two steps in the relation between piemontite and spessartine can be distinguished. Spessartine formation is due to the reaction between piemontite and braunite because of the increasing temperature.

Keywords: high-pressure metamorphism, piemontite, spessartine, Japan, Sanbagawa.

Introduction

Piemontite-quartz schists are highly oxidized manganese-rich metamorphic rocks which have been frequently reported in high-pressure metamorphic belts, for example in Europe (REINECKE, 1986), New Zealand (KAWACHI et al., 1983) and Japan (MINAKAWA, 1992). Studying such rocks is important from different points of view. As source of unusual suites of Mn-rich minerals some of which are rare or unique species (e.g. Mn-rich pyroxene, Mn-rich garnet), it provides the best opportunity to obtain valuable information on the crystal chemistry of unusual minerals, and the chemical variations of high-Mn minerals reveal important information on the metamorphic P-T-t history of the host rocks. Piemontite (Pi), pistacite (Ps) and clinozoisite (Czo) are three end-members which can explain the chemical composition of piemontite within its basic formula $Ca_2(Al, Mn^{3+}, Fe^{3+})_3Si_3O_{12}(OH)$. Piemontite is the Mn end-member of the solid solution and is characterized by a unique pleochroism. Some amounts of Mn²⁺, Sr and Pb can substitute on the

¹ Department of Geology and Mineralogy, Graduate School of Science, Kyoto University, Kyoto, Japan.

² Present address: Department of Geology, Faculty of Science, Zanjan University, Zanjan, Iran. <jav7d@mavara.com>

J. IZADYAR



Ca site (DEER et al., 1992; KESKINEN and LIOU, 1979; KAWACHI et al., 1983). The chemical composition of natural piemontite varies from 33 mol% $Mn^{3+} + Fe^{3+} [Mn^{3+} mol\% = Mn^{3+} \times 100 / (Mn^{3+} + Al)]$ + Fe^{3+}] to 52 mol% which is wider than that of epidote (REINECKE, 1986). The highest Mn³⁺ content has been found for piemontite coexisting with braunite (CHOPIN, 1978) or ardennite (REI-NECKE, 1986). Experimental investigations on Fefree piemontite by ANASTASIOU and LANGER (1976, 1977) revealed that piemontite with Al/Mn ratios ranging from 5/1 to 5/7 can be synthesized at the Mn₂O₃-MnO₂ buffer and 7-15 kbar fluid pressure. On the other hand, KESKINEN and LIOU (1979) synthesized piemontite at 2 kbar and fO₂ buffered at Cu₂O/Cu and CuO/Cu₂O, yielding an Al/Mn ratio of 2:1.

In central Shikoku, Sanbagawa belt, Japan, piemontite with variable chemical compositions and complex zoning patterns is abundant in highly oxidized quartz schists. Thus the object of this contribution is to present: chemical composition of piemontites and other Mn-rich minerals from various assemblages; the effect of increasing metamorphic grade on chemical composition of piemontite; the relation between piemontite zonation and Sanbagawa metamorphic conditions; and spessartine-forming reaction derived from textural and chemical evidence.

Geological setting, sample locality and analytical procedure

The Sanbagawa metamorphic belt is an area in Southwest Japan that was subjected to a Cretaceous regional metamorphism of intermediate high pressure (Fig. 1). A large proportion of the Sanbagawa belt consists of metapelites interbedded with varying amounts of oceanic crustal material and is accompanied by ultramafic and mafic tectonic blocks in the highest grade part of central Shikoku (e.g. WALLIS et al., 1992). Quartz schists, e.g., manganiferous and ferrugenous metacherts, are common in the schists (BANNO and SAKAI, 1989). In central Shikoku, the Sanbagawa metamorphic belt is widest and can be divided into four mineral zones; chlorite, garnet, albite-biotite and oligoclase-biotite zones, based on the appearance of index minerals in pelitic schists (e.g. HI-GASHINO, 1975; ENAMI, 1983; BANNO and SAKAI, 1989). Metamorphic conditions in this area were estimated at 250-300 °C and 5-6 kbar for the lower chlorite zone, and 610 °C and 10-12 kbar for the oligoclase-biotite zone (BANNO and SAKAI, 1989). Thus, the overall slope of the P-T trajectory from the chlorite zone to the albite-biotite zone is positive (prograde path of metamorphism), but chemical zonation of some minerals such as amphibole and garnet suggest also a retrograde path (BANNO and SAKAI, 1989). For further details on the geology and petrology of this area, the reader is referred to BANNO and SAKAI (1989), HIGASHI-NO (1990), and WALLIS and BANNO (1990).

The studied samples were obtained from the garnet zone of the Asemi-gawa area and the albite-biotite zone of the Besshi area (Fig. 1).

Chemical analyses were obtained by a scanning electron microscope (Hitachi S550) with a Kevex energy-dispersive X-ray analytical system using the correction subroutine of Magic V program of Kevex 7000 μ system at Kyoto University. Standards were: pure metals for Mn, Cr and Ni; albite, wollastonite and orthoclase for Na, Ca and K; and oxides for Al, Fe, Mg, Ti.

Details of the analytical procedure follow MORI and KANEHIRA (1984) and HIRAJIMA and BANNO (1991). Back-scattered electron (BSE) images were obtained using a GW-BSE detector system of the scanning electron microscope (Hitachi S530 and S550) at 20 kV and a beam current of about 1000 pA.

Petrography

Piemontite-quartz schists from the Asemi-gawa area commonly show compositional banding with alternating piemontite-rich and quartz-rich bands. The width of the piemontite-rich bands ranges from 0.5 to 1.5 mm. They are mainly composed of piemontite, garnet, talc and hematite with subordinate amounts of quartz, phengite, albite and braunite. The quartz-rich bands (1-2 mm thick) mostly contain quartz, phengite, albite, talc and chlorite with minor amounts of piemontite, garnet, hematite and braunite. In comparison to the Asemi-gawa area, samples from the Besshi area do not show clear compositional banding. The Sanbagawa piemontite-quartz schists can be divided into talc-bearing and talc-free types and their mineral assemblages in the relevant areas are as follows (abbreviations following KRETZ, 1983):

Grt zone (Asemi-gawa area): Qtz + Ab + Phn + Chl + Grt + Piemontite + Braunite + Ap + Hem ± Tlc ± Crossite.

Bt zone (Besshi area): Qtz + Ab + Phn + Chl + Grt + Piemontite + Braunite + Ap + Hem ± Tlc ± Barroisite ± Dol.

For comparison, the mineral parageneses of piemontite-quartz schists, pelitic schists and hematite-bearing quartz schists are listed in figure 2.

		Chlorite zone	Garnet zone	Albite-biotite zone	Oligoclase- biotite zone
Pelitic schist	Chlorite Garnet Biotite Hornblende Lawsonite Albite				– Oligoclase
Hematite- Qtz schist	Winchite Crossite Hornblende Chlorite Sodic- pyroxene Garnet Albite				
Piemontite- Qtz schist	Chlorite Garnet Piemontite Braunite Talc Albite Crossite Barroisite Hematite				

Fig. 2 The stability of silicate minerals in pelitic schists, hematite-quartz schists and piemontite-quartz schists from the Sanbagawa belt. Solid lines: major constituents; dashed lines: minor constituents. Quartz, phengite and epidote are always present. Phlogopite occurs as a secondary mineral in the piemontite-quartz schist. Mineral assemblages of the pelitic schist and the hematite-quartz schist are adapted from ENAMI (1983) and ENAMI et al. (1994).

<i>Tab.</i> 1	Representative ana	vses of p	hengite,	phlogopite.	talc.	chlorite and	d amphibol	le
				a set of the set of th			a company of	

	Asemi-gawa						Besshi			
Point No.	Phengite 18	Chlorite 145	Crossite 739	Phlogopite 1415	Talc 156	Phengite 34	Chlorite 22	Barroisite 17	Talc 44	
SiO ₂	48.93 (0.04)	30.22 (0.04)	59.17 (0.04)	44.61 (0.04)	63.15 (0.05)	48.18	29.83	54,53 (0.05)	62.73	
TiO ₂	0.48 (0.01)	-	0.03 (0.00)	-	_	1.07	-		-	
Al_2O_3	26.28 (0.04)	19.16 (0.03)	7.19 (0.02)	11.15 (0.03)		27.71	18.88	6.26 (0.02)	_	
Fe ₂ O ₃	3.99 (0.02)	0.78 (0.01)	7.19 (0.03)	0.81 (0.01)	0.07 (0.00)	4.38	0.74	7.39 (0.03)	0.15	
Mn ₂ O ₃	0.45 (0.01)		0.31 (0.01)	.	_	0.20		1.56 (0.01)	_	
MnO	-	2.32 (0.02)	-	1.12 (0.01)	0.33 (0.01)	_	1.37 (0.01)	_	0.34	
MgO	3.17 (0.02)	31.36 (0.05)	14.69 (0.04)	24.80 (0.04)	30.96 (0.05)	3.17	31.31	17.63 (0.04)	30.93	
CaO	_	0.07 (0.00)	0.59 (0.01)		_	-	-	5.32 (0.02)	an <u>-</u> a 18	
Na ₂ O	0.60 (0.01)	—	7.00 (0.04)	-		1.22 (0.02)	and epote	5.77 (0.04)	_	
K ₂ O	9.53 (0.02)	-	0.05 (0.00)	10.12 (0.03)	-	9.78 (0.03)	-	-	_	
Total	93.43	83.91	96.22	92.61	94.51	95.71	82.13	98.46	94.15	
	O = 22	O = 28	O = 23	O = 22	O = 22	O = 22	O = 28	O = 23	O = 22	
Si	6.67	5.85	8.05	6.36	8.00	6.45	5.90	7.47	8.00	
Al ^{IV}	1.33	2.15	0.00	1.64	-	1.55	2.10	0.53		
Al ^{VI}	2.89	2.22	1.15	0.23	_	2.82	2.30	0.48	11.01	
Ti	0.05	_	0.003	_	-	0.10	1. N. <u>2</u> 8. N	54527	3100 <u> </u>	
Fe ³⁺	0.40	0.11	0.74	0.09	0.007	0.44	0.10	0.76	0.01	
Mn ³⁺	0.05	-	0.03	-	_	0.02		0.16		
Mn ²⁺		0.34		0.13	0.03	-	0.23		0.03	
Mg	0.64	9.04	2.98	5.27	5.87	0.63	9.22	3.60	5.89	
Ca		0.01	0.08		_		_	0.78		
Na	0.16	1 1 1 1 1 1	1.85	-	_	0.32	510 - C	1.53	191-19	
K	1.66	—	0.01	1.84	_	1.67	-			
Total	13.85	19.72	14.89	15.56	13.90	14.00	19.85	15.31	13.93	

Numbers in parentheses indicate precision of each element. Notice that in the Besshi samples only precision is mentioned that differs with the relevant mineral in the Asemi-gawa samples.

Chemical composition of Mn-poor minerals

Talc can be distinguished from phengite by its lower interference colour under the microscope. Talc occurs as a tabular aggregate or as an intercalation with phengite or chlorite. The composition of talc is very close to the ideal formula (Tab. 1).

Phengite, a major phyllosilicate in the studied rocks, occurs both in the matrix and as an inclusion in albite porphyroblasts. Phengite in the Asemi-gawa area is higher in Si than in the Besshi area reflecting the difference of maximum attained temperature during metamorphism (Tab. 1).

Chlorite is in contact with talc and phengite and is commonly associated with piemontite, garnet, braunite and hematite. The composition of the analyzed chlorite is close to the ideal endmember of clinochlore (Tab. 1).

Phlogopite can be found only replacing phengite at the margin in the Asemi-gawa area but is neither in contact with chlorite nor talc. The phlogopite composition deviates from the ideal end-member by excess Si on the tetrahedral site which could be accompanied by excess Al on the octahedral position and a K deficiency (Tab. 1).

The colourless amphiboles in the Asemi-gawa samples occur only as inclusions in albite and garnet porphyroblasts, and their compositions correspond to crossite (Tab. 1). In the Besshi samples they are matrix phases and are magnesiokataphorite or barroisite.

Porphyroblasts of albite commonly contain abundant inclusions of piemontite, hematite, phengite, talc, colourless amphibole, quartz and rarely garnet.

Hematite occurs in the matrix and as inclusions in albite and garnet, and its average composition is $(Fe_{1,90}Mn_{0.08}Ti_{0.01})_{1.99}O_3$. Careful analyses did not reveal any compositional differences between the two areas.

For further details on the chemical composition and the petrographical significance of the talc-phengite-albite assemblage see IZADYAR et al. (2000).

Chemical composition of Mn-rich minerals

PIEMONTITE

Piemontites occur as subhedral and euhedral crystals and show strong pleochroism with X = yellow and Z = pinkish red. Piemontites are present in the matrix as well as enclosed by garnet and albite in piemontite-rich and quartz-rich bands. The structural formula was computed following SMITH and ALBEE (1967) and, for most analyzed piemortites, shows Si = 3.00, Al + Fe³⁺ +

 Mn^{3+} = 3.00 and Ca = 2.00 per 12.5 oxygens. Therefore, substitution of Al for Si is not indicated by the data. Usually, Mn is present as Mn^{3+} in the octahedral sites but in some piemontites, a certain amount of Mn^{2+} may substitute for Ca.

In piemontites from the talc-bearing schists two zones can be distinguished. The inner zone (core) often forms large crystals; the outer zone is





Fig. 4 Chemical composition of the piemontite from the talc-bearing schists.

Point No.	5112	5212	1919	2019	332	122
Core/Rim	Core	Rim	Core	Rim	Core	Rim
SiO ₂ wt%	37.74 (0.04)	38.55	38.12	39.00	37.61	38.43
TiO ₂	0.11(0.00)	0.07	0.06	0.07	0.04	0.06
Al_2O_3	20.55 (0.03)	19.07	20.59	19.51	20.30	18.81
Fe ₂ O ₃	2.90 (0.02)	9.78	3.82	9.34	6.11 (0.03)	9 10
Mn_2O_3	14.58 (0.04)	11.75	13.28	12.70	11.35	12 34
CaO	22.99 (0.03)	19.81	22.48	18.98	23.04	19.64
Total	98.87	99.03	98.35	99.60	98.45	98.38
No. of cations base	ed on O = 12.5					10
Si	3.01	3.07	3.04	3.08	3.01	3.05
Ti	0.01	0.01	0.01	0.01	0.00	0.01
Al	1.94	1.80	1.94	1.81	1.92	1.78
Fe ³⁺	0.18	0.58	0.24	0.56	0.37	0.55
Mn^{3+}	0.88	0.62	0.81	0.61	0.69	0.67
Total	3.01	3.01	3.00	2.99	2.98	3.01
Ca	1.96	1.69	1.93	1.60	1.98	1.69
Mn^{2+}	- 19 a	0.01	_	0.14	_	0.08
Total	1.96	1.70	1.93	1.74	1.98	1.77
End members					en e	(z-
Pi	29.34	20.66	27.00	20.33	23.00	22.33
Ps	6.00	19.33	8.00	18.66	12.33	18.33
Czo	64.66	60.01	65.00	61.01	64.67	59.34

Tab. 2 Representative analyses of the Asemi-gawa piemontites from the talc-bearing schists.

Tab. 2 (cont.) Representative analyses of the Besshi piemontites from the talc-bearing schists.

DINI						
Point No.	311	312	39	310	45	46
Core/Rim	Core	Rim	Core	Rim	Core	Rim
SiO ₂ wt%	37.50 (0.04)	37.82	37.89	37.89	37.58	37.47
TiO ₂	-	-	_	<u> </u>		
Al_2O_3	20.87 (0.03)	21.59	21.25	21.19	21.86	20.15
Fe_2O_3	11.12 (0.04)	10.61	8.89 (0.03)	9.31	4.60(0.03)	717
Mn_2O_3	6.69 (0.03)	6.32	7.03	7.42	10.42	10.93
CaO	20.29 (0.03)	20.63	21.30	20.85	21.98	20.95
Total	96.47	96.97	96.36	96.67	96.44	96.67
No. of cations based	l on O = 12.5					
Si	2.92	2.92	2.95	2.94	2.90	2.93
Ti	-					-
Al	1.92	1.97	1.95	1.94	2.00	1.85
Fe ³⁺	0.65	0.62	0.52	0.54	0.26	0.42
Mn^{3+}	0.40	0.37	0.42	0.44	0.60	0.65
Total	2.97	2.96	2.89	2.92	2.86	2.92
Ca	1.70	1.70	1.77	1.73	1.83	1.75
Mn^{2+}	-				_	_
Total	1.70	1.70	1.77	1.73	1.83	1.75
End members						
Pi	13.33	12.30	14.00	14.66	20.00	21.66
Ps	21.60	20.60	17.33	18.00	8.66	14.00
Czo	65.07	67.10	68.67	67.34	71.34	64.34

Numbers in parentheses indicate precision of each element. Other columns have similar precision as column one, otherwise it is mentioned.

All of the Mn and Fe are considered as Mn_2O_3 and Fe_2O_3 . Mn²⁺ and Mn³⁺ are computed based on structural formula. a narrow rim surrounding the core (Fig. 3). In the Asemi samples, the rim contains more Fe³⁺ but lower Mn^{3+} and Al than the core (average core composition: $X_{Pi} = 0.26$, $X_{Ps} = 0.09$, $X_{Czo} = 0.65$; average rim composition: $X_{Pi} = 0.21$, $X_{Ps} = 0.19$, $X_{Czo} = 0.60$; $X_{Pi} = Mn^{3+}/(Al + Mn^{3+} + Fe^{3+})$; cf. Fig. 4 and Tab. 2). Inspection of table 2 reveals that the rim is lower in Ca than the core and, in this case, some divalent Mn may substitute for Ca. In the Besshi samples, piemontite does not exhibit clear zoning but sometimes slight zonation could be seen, in

which case the rim is richer in Fe³⁺ and Mn³⁺ than the core (average core composition: $X_{Pi} = 0.17$, $X_{Ps} = 0.13$, $X_{Czo} = 0.70$; average rim composition: $X_{Pi} = 0.18$, $X_{Ps} = 0.16$, $X_{Czo} = 0.66$; cf. Fig. 4 and Tab. 2). Sometimes another type of zoning is visible where Fe³⁺ and Mn³⁺ decrease but Al increases from core to rim (cf. points 311–312 in Tab. 2).

Piemontites in the talc-free type show complex chemical zoning patterns. The cores are highly enriched in Mn³⁺ and are surrounded by a mantle which is rich in Al and Fe³⁺. Zoning is more

Tab. 3 Representative analyses of the Asemi-gawa piemontites from the talc-free schists.

Point No.	136	137	138	131	132	133	11	12	13	
Core/Rim	Core	Mantle	Rim	Core	Mantle	Rim	Core	Mantle	Rim	
SiO2 wt%	38.21 (0.04)	37.89	39.96	38.09	38.28	38.65	38.38	38.48	39.24	
TiO ₂	0.00(0.00)	0.08	0.00	-	_	0.05	0.04	-	0.03	
Al ₂ Õ ₃	20.04 (0.03)	22.61	22.11	21.32	22.86	22.00	20.97	22.64	22.46	
Fe ₂ O ₃	7.59 (0.03)	10.88 (0.04)	8.49	6.52	10.34	9.63	7.52	10.77	10.57	
Mn ₂ O ₃	9.44 (0.03)	2.04(0.02)	5.38 (0.02)	8.90	2.52 (0.02)	4.68 (0.02)	8.58	2.23	3.87 (0.02)	
CaÕ	21.27 (0.03)	23.05	22.25	22.47	23.01	22.83	22.18	23.14	22.89	
Total	96.55	96.55	98.19	97.30	97.01	97.84	97.67	97.26	99.06	
No. of cation	No. of cations based on $O = 12.5$									
Si	3.10	3.10	3.10	3.05	3.05	3.05	3.05	3.05	3.05	
Ti	0.00	0.01	0.00	—	_	0.01	0.01	-	0.00	
Al	1.91	2.14	2.05	2.02	2.15	2.06	1.98	2.14	2.07	
Fe ³⁺	0.46	0.66	0.50	0.40	0.62	0.57	0.45	0.64	0.62	
Mn ³⁺	0.58	0.12	0.32	0.55	0.15	0.28	0.52	0.13	0.23	
Total	2.95	2.93	2.87	2.97	2.92	2.92	2.96	2.91	2.92	
Са	1.85	1.98	1.88	1.94	1.97	1.94	1.90	1.98	1.93	
Mn ²⁺		_		_	-	-	-	-	-	
Total	1.85	1.98	1.88	1.94	1.97	1.94	1.90	1.98	1.93	
End membe	ers									
Pi	19.33	4.00	10.66	18.33	5.00	9.33	17.33	4.33	7.66	
Ps	15.33	22.00	16.66	13.33	20.66	19.00	15.00	21.33	20.66	
Czo	65.34	74.00	72.68	68.34	74.34	71.67	67.67	74.34	71.68	

Tab. 3 (cont.) Representative analyses of the Besshi piemontites from the talc-free schists.

Point No. Core/Rim	16 Core	17 Rim	18 Core	19 Rim	11 Core	12 Rim	28 Core	29 Rim		
SiO ₂ wt%	38.35	38.20	37.79	38.33	39.04	40.18	38.58	38.57		
TiO	_	_		_						
AlaOa	22.22	23.02	21.20	22.97	22.23	21.32	22.67	23.30		
Fe ₂ O ₂	11.72	12.41	10.77	11.76	12.31	10.63	10.54	12.07		
Mn ₂ O ₃	3.24	2.16	5.01	1.32	4.41	6.45	4.07	1.40		
CaÔ	21.89	21.69	21.47	22.19	20.97	20.57	21.85	22.34		
Total	97.42	97.48	96.24	96.57	98.96	99.15	97.71	97.68		
No. of cations based on $Q = 12.5$										
Si	3.00	2.98	3.00	3.10	2.95	3.00	3.00	3.10		
Ti	-			-	_	-	-			
Al	2.03	2.17	1.97	2.13	1.98	1.90	2.10	2.10		
Fe ³⁺	0.68	0.76	0.70	0.68	0.70	0.60	0.58	0.68		
Mn ³⁺	0.19	0.09	0.29	0.08	0.25	0.40	0.30	0.08		
Total	2.90	3.02	2.96	2.89	2.93	2.90	2.98	2.86		
Ca	1.94	1.89	1.97	1.94	1.70	1.67	1.92	1.94		
Mn ²⁺	_			_	_	_				
Total	7.84	7.89	7.93	7.93	7.58	7.54	7.90	7.90		
End members										
Pi	6.33	3.00	9.66	2.66	8.33	13.33	10.00	2.66		
Ps	22.66	25.33	23.33	22.66	23.33	20.00	19.33	22.66		
Czo	71.01	71.67	67.01	74.68	68.34	66.67	70.67	74.68		

Numbers in p arentheses indicate precision of each element. Other columns have similar precision as column one, otherwise it is mentioned. All of the Mn and Fe are considered as Mn_2O_3 and Fe_2O_3 . Mn^{2+} and Mn^{3+} are computed based on structural formula.

	In piemontite	rich band	In quartz-rich band	Enclosed in	albite C	hanged rim of gar	net at contact to n	natrix
Point No.	27	43c	130	150	1519	1619	4619	4191
SiO2 wt%	37.39 (0.04)	37.20	37.46	37.92	37.61	37.33	38.98	39.34
TiO ₂	0.23 (0.01)	0.28	0.25	_	0.13 (0.00)	0.14	0.15	0.16
Al_2O_3	19.58 (0.03)	19.01	19.62	19.71	19.21	19.67	19.24 (0.04)	20.67
Fe ₂ O ₃	1.74 (0.02)	1.96	1.73	1.93	4.02 (0.03)	2.65 (0.03)	3.05	3.75
MnO	36.92 (0.06)	36.28	35.71	36.24	25.40 (0.05)	31.03	32.31	28.29
MgO	0.71(0.01)	0.42	0.57	0.63	0.27	0.17	0.50	0.27
CaO	4.08 (0.02)	3.97	4.47	4.36	12.05 (0.03)	8.95	7.04	9.64
Total	100.65	99.12	99.81	100.79	98.69	99.94	101.27	102.12
No. of catio	ons based on 12	oxygens						- 65 - 5
Si	3.04	3.05	3.04	3.04	3.03	3.00	3.07	3.05
Ti	0.01	0.01	0.01		0.008	0.003	0.008	0.01
Al	1.86	1.84	1.87	1.87	1.83	1.87	1.89	1.89
Fe ²⁺	—	-	0.07	0.02	0.02	0.11		
Fe ³⁺	0.09	0.12	0.10	0.12	0.17	0.13	0.11	0.11
Mn	2.55	2.52	2.46	2.47	1.74	2.13	2.18	1.86
Mg	0.08	0.05	0.07	0.08	0.03	0.02	0.05	0.03
Ca	0.34	0.34	0.38	0.37	1.04	0.80	0.60	0.80
Total	7.97	7.93	7.93	7.95	7.92	7.97	7.93	7.86
End memb	bers							
Sps	85.86	86.60	84.50	84.50	60.40	71.70	76.70	66.42
Pyr	2.70	1.70	2.40	2.74	1.04	0.70	1.60	1.07
Grs	6.84	5.60	8.10	6.76	27.66	20.40	15.50	23.11
And	4.60	6.10	5.00	6.00	8.50	6.50	5.50	5.50
Alm	0.00	0.00	0.00	0.00	2.40	0.70	0.70	3.90

Tab. 4 Representative analyses of garnet from the talc-bearing schist in the Asemi-gawa area.

Numbers in parentheses indicate precision of each element. Other columns have similar precision as column one, otherwise it is mentioned.

All of the Fe is considered as Fe_2O_3 .

Fe²⁺ and Fe³⁺ are computed based on structural formula.

Tab. 5 Representative analyses of braunite.

Point No.	1832	2032	2132
	Asemi-gawa Be	sshi-Kokury	o Besshi-Seki
SiO ₂ wt%	10.44 (0.02)	10.34	10.73
TiO ₂	0.11(0.00)	0.15	0.14
Al_2O_3	0.40(0.02)	0.30	0.35
Fe_2O_3 (total)	12.66 (0.04)	12.35	12.24
Mn_2O_3 (total)	76.38 (0.08)	76.25	77.23
MgO	0.09(0.00)	0.00	0.08
CaO	0.28 (0.01)	0.26	0.10
Total	100.36	99.65	100.87
No. of cations b	based on $O = 12$		
Si	1.02	1.01	1.04
Ti	0.01	0.01	0.01
Al	0.05	0.03	0.03
Fe ³⁺	0.93	0.91	0.89
Mn^{3+}	5.07	5.09	5.37
Mn^{2+}	0.60	0.61	0.53
Mg	0.01	0.00	0.01
Ca	0.03	0.03	0.01
Total	7.72	7.69	7.89

Numbers in parentheses indicate precision of each element. Other columns have similar precision as column one. Mn²⁺ and Mn³⁺ are computed based on structural formula.



Fig. 5 Back scattered electron image of piemontite from the talc-free piemontite-quartz schist showing three zones: 1 = core with enrichment of Mn^{3+} -content, 2 = mantle with depletion in Mn^{3+} -content, and 3 = rim with enrichment in Mn^{3+} -content.

complicated by the presence of an outer rim which is further enriched in Mn^{3+} (Fig. 5). In the Asemi-gawa area, three generations of piemontites are visible even within one grain. In this case, the core with an average composition of $X_{Pi} =$ 0.18, $X_{Ps} = 0.15$, $X_{Czo} = 0.67$ is surrounded by a mantle with an average composition of $X_{Pi} = 0.04$, $X_{Ps} = 0.21$, $X_{Czo} = 0.74$. The mantle, in turn, is surrounded by a rim characterized by $X_{Pi} = 0.10$, $X_{Ps} = 0.19$, $X_{Czo} = 0.71$ (cf. Fig. 6 and Tab. 3). In the Besshi piemontites, zonation is not very distinct, but two types may be observed: from core to rim, Mn and Fe decrease and Al increases (cf. points 18–19 and 28–29 in Tab. 3); and Mn increases and Al and Fe decrease (cf. points 11–12 in Tab. 3).

GARNET

Porphyroblasts of garnet occur in both piemontite- and quartz-rich bands. Garnet in piemontiterich bands is euhedral and its size ranges from 1 to 3 mm. This type of garnet is poikiloblastic, containing many inclusions of piemontite, quartz, hematite, braunite, talc and amphibole. Chemically, these garnets are homogeneous Ca–Fe-bearing



Fig. 6 Chemical composition and compositional variation of piemontite from the talc-free schists.

spessartines ($X_{Sps} = 0.85$, $X_{Grs} = 0.06$ and $X_{And} = 0.05$). The garnet in quartz-rich bands includes only small amounts of hematite, piemontite and braunite, and garnet is also a Ca–Fe-bearing spessartine ($X_{Sps} = 0.84$, $X_{Grs} = 0.07$ and $X_{And} = 0.06$). In one sample, garnet occurs within an albite porphyroblast in the quartz-rich band and it is chemically the same as that outside albite (Tab. 4).

BRAUNITE

Braunite usually occurs both in the matrix and as inclusions in garnet in the piemontite-rich bands and rarely as inclusions in albite and garnet in the quartz-rich bands. Commonly, it is euhedral and chemically homogeneous. There are no differences between chemical compositions of braunite in the Asemi-gawa and Besshi areas (Tab. 5).

Variation of chemical composition of piemontite with changing metamorphic grade

Experimental investigations on piemontites were performed on Fe³⁺-free piemontites by ANASTA-SIOU and LANGER (1976, 1977) and KESKINEN and LIOU (1979), but most of the natural piemontites contain some Fe³⁺. In order to correlate the experimental results to Fe³⁺-bearing piemontite, they compared their data to epidote stability studies of HOLDWAY (1972). ANASTASIOU and LANGER concluded that the Mn³⁺-bearing phase shows higher temperature stability than Fe³⁺-bearing epidote whilst KESKINEN and LIOU (1979) considered that introduction of Fe in piemontite will extend its stability towards higher temperature. Because the Sanbagawa belt is one of the most extensively studied metamorphic complexes (TAKASU et al., 1994), it provides the best opportunity to study the relation between chemical variation of piemontite and changing P,T, fO₂ and bulk-rock composition. Thus, the samples were collected from the garnet zone in the Asemi-gawa area and albite-biotite zone of the Besshi area. Comparing the average core compositions of piemontites in the talc-bearing type between both areas reveals that the Asemi piemontite is richer in piemontite but poorer in pistacite and clinozoisite components (Asemi-gawa: $X_{Pi} = 0.26$, $X_{Ps} = 0.09$, $X_{Czo} = 0.65$; Besshi: $X_{Pi} = 0.15$, $X_{Ps} = 0.17$, $X_{Czo} = 0.68$; cf. Fig. 4 and Tab. 2). The same results were obtained by comparing average core compositions of piemontites in the talc-free samples (Asemi-gawa: $X_{Pi} = 0.18, X_{Ps} = 0.15, X_{Czo} = 0.67$; Besshi: $X_{Pi} = 0.09$, $X_{Ps} = 0.22, X_{Czo} = 0.69$; cf. Tab. 3). The chemical composition of piemontite is mainly controlled by

		2	1		0				
Point No. Core/Rim	95 Core	96 Rim	97 Changed Rim	3419 Core	3319 Rim	3519 Changed Rim	2191 Core	1191 Rim	3191 Changed Rim
SiO ₂ wt%	38.11 (0.04)	38.49	37.49	37.94	36.34	37.69	38.28	37.85	39.11
TiO ₂	0.04 (0.00)	0.06	0.04	0.04	0.07	0.06	C-RUIS	0.04	0.05
Al_2O_3	20.22 (0.03)	19.23	19.84	20.15	18.19	19.24	19.35	20.02	20.19
Fe ₂ O ₃	5.40 (0.03)	8.50	5.49	5.88	9.38	9.08	9.32	5.98	7.02
Mn ₂ O ₃	12.16 (0.03)	11.63	12.87	12.00	12.88	13.58	11.73	11.52	13.07
CaO	22.23 (0.03)	18.91	21.89	23.11	19.45	19.29	19.92	22.86	20.25
Total	98.16	96.82	97.62	99.12	96.31	98.94	98.60	98.27	99.69
No. of catio	ons based on O	= 25					28.217.1	1-1089F	WARD STREET
Si	3.05	3.10	3.00	3.02	3.00	3.02	3.05	3.02	3.08
Ti	0.002	0.004	0.002	0.002	0.004	0.003	r s <u>e</u> der s	0.002	0.003
Al	1.91	1.83	1.89	1.90	1.77	1.82	1.84	1.90	1.87
Fe ³⁺	0.32	0.51	0.33	0.35	0.58	0.54	0.56	0.36	0.41
Mn ³⁺	0.74	0.64	0.77	0.73	0.64	0.63	0.71	0.70	0.71
Total	2.972	2.984	2.99	2.982	2.994	2.99	3.11	2.962	2.993
Ca	1.91	1.64	1.90	1.98	1.73	1.65	1.70	1.97	1.71
Mn^{2+}		0.07	0.02	-	0.17	0.19	-	- 1	0.07
Total	1.91	1.71	1.92	1.98	1.90	1.84	1.70	1.97	1.78
End memb	ers						L P I	0.08	
Pi	24.66	21.33	25.66	24.33	21.41	21.00	23.66	23.33	23.66
Ps	10.66	17.00	11.00	11.66	19.40	18.00	18.66	12.00	13.66
Czo	64.68	61.67	63.34	64.01	59.19	61.00	57.68	64.67	62.68

Tab. 6 Representative analyses of piemontites at contact to garnet.

Numbers in parentheses indicate precision of each element. Other columns have similar precision as column one. All of the Mn and Fe are considered as Mn_2O_3 and Fe_2O_3 . Mn^{2+} and Mn^{3+} are computed based on structural formula.

oxygen fugacity, bulk composition, pressure and temperature (KESKINEN and LIOU, 1979). To investigate the fO₂ effect, mineralogical and geochemical evidences were used. Braunite and hematite are two fO₂ indicator minerals in the studied samples, and they were analyzed in detail. The results showed that braunite and hematite are homogenous and that there is no difference on chemical compositions of braunite and hematite between the Asemi-gawa and the Besshi areas. In fact, both areas are located within oxidized assemblage according to the CHOPIN (1978) definition because Mn³⁺ is present in the minerals such as piemontite and braunite, while Mn²⁺ is mostly incorporated in spessartine. Thus, the oxygen fugacity is inferred to be similar in the Asemi and Besshi areas. This conclusion was also supported by determining oxidation ratio (Fe₂O₃ \times 100/ $(Fe_2O_3 + FeO))$ in the whole rock. In both areas, these ratios are around 100% (Tab. 7). Geochemical investigations on the Sanbagawa piemontitequartz schists show that bulk compositions of the two areas are almost identical (Tab. 7). Therefore, the difference in composition of piemontite between the Asemi-gawa and Besshi areas must result from the difference in metamorphic grade. Thus, it appears that the epidote end-member component in piemontite increases from the garnet zone to the albite-biotite zone. Zoned crystals are good indicators of changing metamorphic conditions and are well-studied in the Sanbagawa belt (BANNO and SAKAI, 1989; OTSUKI and BAN-NO, 1990). Even though such observations in other

minerals have been widely used to clarify metamorphic condition of the Sanbagawa belt, little attention has been given to chemical zonation of piemontite. This study shows that piemontite composition is sensitive to changing metamorphic grade. In the Sanbagawa belt, thus, the piemontite compositions show together with experimental results (KESKINEN and LIOU, 1979), that the chemical variation from core to rim in the talc-bearing and from core to mantle in the talc-free types may be related to prograde metamorphism. Additionally, the chemical trends from mantle to rim in the talc-free type may be attributed to the retrograde path of the Sanbagawa metamorphism.

Reaction relations between piemontite and spessartine

Based on textural and chemical criteria two different steps in the relation between piemontite and spessartine can be distinguished. The first step is the spessartine formation. Texturally, spessartine containing many inclusions of piemontite and braunite does not occur as a fine-grained matrix mineral but only as inclusion in albite porphyroblasts. On the other hand, textural relations indicate that braunite occurs as inclusion in spessartine or as a matrix mineral close to the core of piemontite from the talc-bearing type. Therefore, it may coexist with the piemontite core. The partitioning of Mn³⁺, Fe³⁺ and Al between piemontite (Pi) and braunite (Br) confirms the assumption of their coexistence. Distribution coefficients have been calculated by the equations:

$$\begin{split} [K_{D(Al-Mn)} &= (X_{Al}/X_{Mn})_{Pi}/(X_{Al}/X_{Mn})_{Br} \\ X_{Al} &= Al \ / \ Al + Mn] \\ [K_{D(Fe-Mn)} &= (X_{Fe}/X_{Mn})_{Pi}/(X_{Fe}/X_{Mn})_{Br} \\ X_{Mn} &= Mn \ / \ Fe + Mn] \end{split}$$

showing that relative to braunite, piemontite is enriched in Al and Fe³⁺ and depleted in Mn³⁺ (Fig. 7). From the textural observations it is concluded that spessartine may be formed through a reaction involving piemontite and braunite. Experimental work by KESKINEN and LIOU (1979) shows that spessartine is a product of piemontite breakdown due to increasing temperature and/or decreasing oxygen fugacity. Thus, spessartine formation may be formed during the prograde path of the Sanbagawa metamorphism.

The second step is the changing chemical composition of piemontite at contact to spessartine.

Tab. 7 Representative bulk composition of piemontite-quartz schists in the Sanbagawa belt.

715.9	Bessh	ii area	Asemi area
Sample no.	1-1aQ	B4C	1213
SiO ₂ wt%	75.67	78.05	78.26
TiO_2	0.37	0.45	0.4
Al_2O_3	5.77	8.01	7.33
Fe ₂ O ₃	3.49	3.7	3.46
FeO	0.17	0.15	0.1
MnO	0.19	0.18	0.22
MgO	2.78	2.15	2.94
CaO	1.53	1.61	1.24
Na ₂ O	1.19	1.36	1.29
K ₂ Õ	0.61	1.03	0.72
P_2O_5	0.18	0.15	0.17
L.O.I.	6.95	2.6	2.97
Total	98.9	99.44	99.1
Oxidation Rat	io*95.36	96.1	97.19

 $*Fe_2O_3 \times 100/(FeO + Fe_2O_3)$



Fig. 7 Distribution of Mn^{3+} between coexisting piemontite and braunite.

Although, the preservation of compositional zoning of Al, Mn and Fe in zoned piemontite is a primary feature produced during growth, there may be some evidences of secondary cation exchange: in the talc-bearing types, for example, the chemical composition of piemontite changes whenever it occurs as inclusion in spessartine or in contact to it in matrix. This change was observed in the rim in a very limited area, about 33 µm away from the piemontite-spessartine contact. At the contact, piemontite tends to be, on average, richer in Mn ($X_{Pi} = 0.24$) and poorer in Fe and Ca ($X_{Czo} =$ 0.62 and $X_{Ps} = 0.13$) as compared to rim ($X_{Pi} =$ 0.22, $X_{Czo} = 0.62$, $X_{Ps} = 0.16$; cf. Tab. 6 and Fig. 8). Chemical change can also be seen at the piemontite-spessartine contact of the garnet, where it becomes richer in Ca and Fe ($X_{Grs} = 0.18$, $X_{And} =$ 0.09) and poorer in Mn ($X_{sps} = 0.73$) compared to the inner part ($X_{sps} = 0.85$, $X_{Grs} = 0.10$, $X_{And} = 0.09$; cf. Tab. 4 and Fig. 8).

Although charge balance calculations in garnets are equivocal, the distribution of cations in the structure of studied garnet did not show any incorporation of trivalent Mn in the octahedral sites (Tab. 4). Indeed, for the mentioned fo₂ conditions, in which Mn can be present both as Mn²⁺ and Mn³⁺, all of the available Mn³⁺ was preferably incorporated in the braunite or piemontite structure and there is no excess of Mn³⁺ to contribute



Fig. 8 Compositional variation of piemontite at the contact to spessartine.

to the garnet structure. Also, this conclusion is well supported by the fact that Mn³⁺-bearing garnets have been synthesized only at pressure higher than 25 kbar (NISHIZAWA and KOIZOMI, 1975) which is very different from the Sanbagawa metamorphic condition. Furthermore, optical absorption spectra and theoretical considerations suggest that Mn is almost totally divalent in garnet structure (FRENTRUP and LANGER, 1982).

Concluding remarks

The following results were obtained from the petrographic and mineralogical study on piemontitequartz schists from the Sanbagwa metamorphic belt:

(1) Piemontites of variable composition and complex chemical zoning are the most abundant Mn-silicates in highly oxidized Mn-rich quartz schists.

(2) The Asemi-gawa piemontites are richer in piemontite component than those from the Besshi area. The difference of metamorphic grade indicates that piemontite with higher epidote contents is stable at higher grade because bulk composition and oxygen fugacity are similar in both areas.

(3) Preservation of Mn, Al, Ca and Fe zoning in piemontite provides the best opportunity to unravel the metamorphic history of the region. The chemical evolution of piemontite can be compared with minerals like garnet. This study revealed prograde and retrograde paths of the Sanbagawa metamorphism.

(4) Besides piemontite, other Mn–Al–Ca silicates are also sensitive to changing metamorphic conditions. In the Sanbagawa belt, spessartine formation and cation exchange between spessartine and piemontite are examples of such sensitivity.

Acknowledgements

The results presented here are part of the author's doctoral thesis. The author wishes to express appreciation to Emeritus Prof. S. Banno and Doctors T. Hirajima and K. Tomita for their valuable helps and critical discussions during the time spent at Kyoto university. The author would like to express his sincere thanks to Prof. R. Gieré for his critical comments on several versions of this manuscript. This study was generously supported by the Ministry of Education of Japan.

References

- ANASTASIOU, P. and LANGER, K. (1976): Synthese und Stabilität von Piemontit, Ca₂Al_{3-p}Mn³⁺_p(Si₂O₇/SiO₄/ O/OH). Fortschr. Mineral., 54, Bh. 1, 3–4.
- ANASTASIOU, P. and LANGER, K. (1977): Synthesis and physical properties of piemontite Ca_2Al_{3-p}

 $Mn_{\rm p}^{\rm 3p}(Si_2O_7/SiO_4/O/OH).$ Contrib. Mineral. Petrol., 60, 225–245.

- BANNO, S. and SAKAI, C. (1989): Geology and metamorphic evolution of the Sanbagawa metamorphic belt, Japan. In: DALY, J.S., CLIFF, R.A. and YARDLEY, B.W.D. (eds): Evolution of metamorphic belts. Geol. Soc. Spec. Pub., 43, 519–532.
- CHOPIN, Ć. (1978): Les paragenèses réduites ou oxydées de concentrations manganésifères des "schistes lustrés" de Haute-Maurienne (Alpes françaises). Bull. Minéral., 101, 514–531.
- DEER, W.A., HOWIE, R.A. and ZUSSMAN, J. (1992): An introduction to rock forming minerals. Longman Scientific & Technical, 2nd edition, 549 pp.
- ENAMI, M. (1983): Petrology of pelitic schists in the oligoclase-biotite zone of the Sanbagawa metamorphic terrain, Japan: phase equilibria in the highest grade zone of a high-pressure intermediate type of metamorphic belt. J. Metamorphic Geol., 1, 141–161.
- ENAMI, M., WALLIS, S.R. and BANNO, Y. (1994): Paragenesis of sodic pyroxene-bearing quartz schist: implication for the P-T history of the Sanbagawa belt. Contrib. Mineral. Petrol., 116, 182–198.
- FRENTRUP, K.R. and LANGER, K. (1982): Microscope absorption spectrometry of silicate microcrystals in the range 4000–5000 cm⁻¹ and its application to garnet end-members synthesized at high pressures. In: SCHREYER, W. (ed.): High pressure research in geoscience. Schweizerbart, Stuttgart, 247–258.
- HIGASHINO, T. (1975): Biotite zone of Sanbagawa metamorphic terrain in the Siragayama area, central Shikoku, Japan. J. Geol. Soc. Japan, 81, 653–670.
- HIGASHINO, T. (1990): The higher grade metamorphic zonation of the Sanbagawa metamorphic belt in central Shikoku, Japan. J. Metamorphic Geol., 8, 413–423.
- HIRAJIMA, T. and BANNO, S. (1991): Electron-microprobe analysis of rock forming minerals with Kevexdelta IV (Quantum detector). Hitachi Scientific Instrument News, 34, 2–7.
- strument News, 34, 2–7. HOLDWAY, M.J. (1972): Thermal stability of Al–Fe epidotes as a function of fO_2 and Fe content. Contrib. Mineral. Petrol., 37, 307–340.
- IZADYAR, J., HIRAJIMA, T. and NAKAMURA, D. (2000): Talc-phengite-albite assemblage in piemontitequartz schists of the Sanbagawa metamorphic belt, central Shikoku, Japan. Island Arc, 9, 145–158.
- KAWACHI, Y., GRAPES, R.H., COOMBS, D.S. and DOWSE, M. (1983): Mineralogy and petrology of a piemontite-bearing schist, western Otago, New Zealand. J. Metamorphic Geol., 1, 353–372.
- KESKINEN, M. and LIOU, J.G. (1979): Synthesis and stability relations of Mn–Al piemontite. Amer. Mineral., 64, 317–328.
- KRETZ, R. (1983): Symbols for rock-forming minerals. Amer. Mineral., 63, 1023–1052.
- MINAKAWA, T. (1992): Study on characteristic mineral assemblages and formation process of metamorphosed manganese ore deposit in the Sanbagawa belt. Memorial of Faculty of Science, Ehime University, 1, 1–74.
- MORI, K. and KANEHIRA, K. (1984): X-ray energy spectrometry for electron probe analysis. J. Geol. Soc. Japan, 90, 271–285.
- NISHIZAWA, H. and KOIZUMI, M. (1975): Synthesis and infrared spectra of Ca₃Mn₂Si₃O₁₂ and Cd₃B₂Si₃O₁₂ (B: Al, Ga, Cr, V, Fe, Mn) garnets. Amer. Mineral., 60, 84–87.
- OTSUKI, M. and BANNO, S. (1990): Prograde and retrograde metamorphism of hematite-bearing basic schists in the Sanbagawa belt in central Shikoku. J. Metamorphic Geol., 8, 425–439.
- REINECKE, T. (1986): Crystal chemistry and reaction relation of piemontites and thulites from highly oxidized

loww grade metamorphic rocks at Vitali, Andros Islannd, Greece. Contrib. Mineral. Petrol., 93, 56-76.

- SMITHH, D. and ALBEE A. (1967): Petrology of a piemont-itee-bearing gneiss, San Gorgonio Pass, California. Contrib. Mineral. Petrol., 16, 189–203.
 TAKASSU, A., WALLIS, S.R., BANNO, S. and DALLMEYER, R.J.D. (1994): Evolution of the Sanbagawa metamor-which halve Longon Lithog 22, 110, 122.
- phhic belt, Japan. Lithos, 33, 119-133.
- WALLLIS, S.R. and BANNO, S. (1990): The Sanbagawa belt treends in research. J. Metamorphic Geol., 8, 393–399.
- WALLIS, S.R., BANNO, S. and RADVANCE, M. (1992): Kinematics, structure and relationship to metamor-phism of east-west flow in the Sanbagawa belt southwest, Japan. Island Arc, 1, 176-185.

Manuscript received March 28, 1999; revision accepted February 28, 2000.