Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =

Bulletin suisse de minéralogie et pétrographie

Band: 81 (2001)

Heft: 2

Artikel: Evidence for Lower Paleazoic magmatism in the Eastern Southalpine

basement : zircon geochronology from Comelico porphyroids

Autor: Meli, Sandro / Klötzli, Urs S.

DOI: https://doi.org/10.5169/seals-61684

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 01.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Evidence for Lower Paleozoic magmatism in the Eastern Southalpine basement: zircon geochronology from Comelico porphyroids

by Sandro Meli¹ and Urs S. Klötzli²

Abstract

Multigrain conventional U/Pb and single grain ²⁰⁷Pb*/²⁰⁶Pb* evaporation data on zircons are presented for two samples of acidic metavolcanic rocks from the Eastern Southalpine basement (Comelico, North-Eastern Italy). Three zircon populations have been distinguished in each sample, reflecting different contributions in the metavolcanics. Clear, colourless, elongated crystals are thought to be single stage magmatic products, whilst zircons belonging to other populations contain inherited cores mantled by magmatic overgrowths. Elongated crystals yielded concordant U/Pb ages (479 ± 8 and 485 ± 8 Ma), whereas other zircon populations were discordant; the inherited components have poorly constrained Archean apparent ages. Evaporation measurements performed on a core-bearing crystal yielded ages ranging from 646 to 715 Ma, which are interpreted as discordant ages of the core; Th/U zoning also points to a magmatic signature of the inherited cores. The conventional U/Pb data represent the first radiometric age determinations of the pre-Variscan acidic volcanism in the Southalpine domain, and the concordant ages constrain the acidic volcanic activity in the Eastern Southalpine basement within the Arenig. The new isotopic ages are consistent with the biostratigraphy and lithostratigraphy of a nearby basement outcrop (Agordo), which has been correlated to Comelico. Moreover, it is also possible to correlate the Southalpine Ordovician volcanism with a volcanic event in the Austroalpine domain (Eisenerz), for which a comparable age has been inferred by means of biostratigraphy.

Keywords: porphyroids, acidic metavolcanics, U/Pb geochronology, Southalpine basement, Ordovician.

1. Introduction

In the Eastern Alps (Fig. 1), the occurrence of metavolcanic layers ("porphyroids") both in the Austroalpine (HEINISCH, 1981) and in the Southalpine (Bellieni and Sassi, 1981) domains document the existence of a widespread pre-Variscan acidic magmatic activity, possibly connected to an orogenic event. An Upper Ordovician age was assigned to the protoliths of the Southalpine porphyroids on a lithostratigraphic basis, at the Caradocian-Ashgillian boundary (SASSI et al., 1979); the assessment was based upon a long distance correlation between the rock sequence in which the Comelico porphyroids occur and that described by FLAJS and SCHÖNLAUB (1976) in the Northern Greywacke Zone (Austria). The hypothesis was strengthened by some palynologic data

from the Agordo phyllites (Col di Foglia Fm.), which point to an Upper Cambrian sedimentation age for the pelitic protoliths underlying the nearby porphyroids (KALVACHEVA et al., 1986): the sedimentation age represents a maximum age for Agordo porphyroids. However, radiometric dating on the metavolcanics was lacking, and the eruption age was loosely constrained. There has been much debate about the geodynamic setting (e.g., HEINISCH and SCHMIDT, 1976; BÖGEL et al., 1979; SASSI and SCHMIDT, 1982; SASSI et al., 1987; HEINISCH, 1988; LOESCHKE, 1989; MELI, 1998) and the major orogenic cycles to which the volcanics can be genetically linked (Caledonian vs. Cadomian). The existence of a "Caledonian" event was maintained by Borsi et al. (1975), Sassi and SCHMIDT (1982), EBNER et al. (1987), on the basis of: (i) the occurrence of an Upper Cambrian-

¹ Dipartimento di Mineralogia e Petrologia, C.so Garibaldi 37, I-35122 Padova, Italy. <sandrom@unipr.it> Present address: Dipartimento di Scienze della Terra, Parco Area delle Scienze 157/A, I-43100 Parma, Italy.

² Laboratorium für Geochronologie, Geozentrum, Universität Wien, Althanstraße 14, A-1090 Wien, Österreich.

Lower Ordovician metamorphic event in some Austroalpine and Southalpine complexes (Borsi et al., 1973, DEL MORO et al., 1984); (ii) the occurrence of Late Ordovician peraluminous granitoids (Borsi et al., 1973); (iii) the possible occurrence of basal conglomerates overlying the Ordovician crystalline rocks (Sassi and Zanferrari, 1972); (iv) the occurrence of intra-Ordovician volcanic activity; this latter was not dated by means of radiometric geochronology. The existence of a "Cadomian" event was maintained, among others, by VAI (1975), and HANDLER et al., (1997), on the basis of (i) geochronological discrepancies of the above mentioned metamorphic and magmatic events with Caledonides s.s.; (ii) 40Ar/39Ar ages on detrital white micas; (iii) chronostratigraphic arguments about the Ordovician molasse in the Carnic Alps. The tectonic significance of both events is still under debate.

Several attempts have been made to date the emplacement of the porphyroid protoliths, but the Variscan overprint prevented any primary age estimate, either with Rb/Sr or K/Ar methods. In this contribution new U/Pb and ²⁰⁷Pb*/²⁰⁶Pb* (* = radiogenic) data are presented, to solve the uncertainty on the age of this widespread magmatic activity.

2. Geologic outline

The Eastern Southalpine Basement outcrops in three different areas (Fig. 1): a belt oriented WNW-ESE, close to the Periadriatic Line, a narrow belt between Trento and Agordo, and a small area close to Recoaro. The metamorphic basement is separated from the overlying Upper Paleozoic clastic sediments ("Verrucano" conglomerates and Val Gardena Sandstones) by an unconformity (Sassi and Spiess, 1993). Acidic metavolcanics and metavolcanoclastics occur as thick layers interbedded with phyllites (Sassi and Zirpoli, 1989); the metavolcanic layers are widespread throughout the whole basement outcrops, with the only exception of Recoaro area (Sassi and Zirpoli, 1968).

SASSI and ZIRPOLI (1989) divided the Southalpine metamorphic basement into three units by means of lithostratigraphic criteria: (1) a Lower Pelitic-Psammitic Complex (LPC); (2) a Volcano-Sedimentary Complex (VSC); (3) an Upper Pelitic-Psammitic Complex (UPC). LPC and UPC are very similar, consisting both of quartzphyllites; their distinction is possible only on the basis of the relative stratigraphic position with respect to the VSC, which has been considered as a guide level (SASSI et al.,1974). This latter consists of a metapelitic-psammitic sequence, interlayered by (i) thick layers of acidic metavolcanics (porphyroids); (ii) a thin, discontinuous level of mafic metavolcanics; (iii) carbonate-bearing phyllites and carbonate schists (SASSI and ZIRPOLI, 1989). At present, no detailed mapping of the three complexes is available.

The Southalpine basement escaped Alpine metamorphism, and only gentle folding and brittle deformations of Alpine age occur (SASSI and Spiess, 1993); the features of the Variscan metamorphism, which developed in two stages at about 345 and 320 Ma (DEL MORO et al., 1980, 1984; MELI, 1994), are therefore preserved. The P-T conditions recorded in the outcropping rocks range within the greenschist facies (see NEUBAU-ER et al., 1999, for a review), and a slight increase of the metamorphic grade is recognizable northwards and westwards, from the chlorite-zone (Comelico) to the almandine-zone (Val Sarentino) (CARDIN et al., 1985; MAZZOLI and SASSI, 1988). Microstructural and petrological studies demonstrated that the whole Southalpine crystalline basement shared the same metamorphic history during Paleozoic (SASSI and ZIRPOLI, 1989).

Petrographic and geochemical features of the acidic metavolcanics are consistent with an anatectic origin of the magmas, generated by partial melting of metapelites; the discriminant diagrams for acidic rocks indicate a late- to post-orogenic geodynamic setting at the time of emplacement (MELI, 1998). BELLIENI and SASSI (1981) included the pre-Variscan eruptives in the "Upper Ordovician Granite-Rhyolite Association", which also comprises coeval granitoids occurring in the Austroalpine domain. MAZZOLI and SASSI (1992) pointed out the geochemical similarities between porphyroids and granitoids, and referred the Ordovician plutonism and volcanism to a unique cycle of magma generation. The sporadic mafic metavolcanics occurring in the upper part of the Volcano-Sedimentary Complex (SASSI and ZIR-POLI, 1989) probably have a within-plate alkaline affinity (VISONÀ and ZANFERRARI, 1985), but their primary age is loosely constrained.

In this work, we focus our attention on the Comelico area, where the degree of recrystallization of the protoliths is lower than that observed in other sectors of the basement (SASSI et al., 1979); moreover, the stratigraphy is better defined and comparable with that of other sites, for which biostratigraphic data are available (ZANFERRARI, 1985; KALVACHEVA et al., 1986). In Comelico, the porphyroids outcrop in two localities: M. Elmo (ME), and M. Cavallino (MC) (Fig. 2); field work demonstrated that ME and MC por-

phyroids are distinct lithologic units (MELI, 1994), being separated by a thick metasedimentary body. Samples ME37 and MC63 were collected from ME and MC, respectively.

3. Petrography

Comelico porphyroids are characterized by a marked grain size bimodality: large crystals, interpreted as relics of magmatic phenocrysts, occur in a very fine grained matrix, which represents a recrystallyzed volcanic microcrystalline and/or glassy groundmass. The pre-metamorphic mineralogical association consists of quartz, K-feldspar, plagioclase, ilmenite, biotite, apatite and zircon. The present feldspar compositions do not reflect their pristine chemistry, as plagioclase has been transformed into albite + epidote ± sericite or albite + calcite + sericite, and alkali mobilization caused the growth of both chess-board albite upon pre-existing K-feldspar (Callegari and De Pieri, 1966) and microcline overgrowth on plagioclase. Quartz porphyroclasts preserve a typ-




Fig. 1 Geological sketch map of the Eastern Alps (simplified from: Heinisch, 1981; Sassi et al., 1985; Barth et al., 1994; Handler et al., 1997; Frey et al., 1999). The letters refer to porphyroid outcrops: (A) Kitzbühel; (B) Zeller Furche-Dienten; (C) Radmer, Blasseneck and Ratschengraben; (D) Eisenerz; (E) Hohe Veitsch; (F) Bressanone-Val Sarentino; (G) Val di Funes; (H) Comelico; (I) Thurntaler; (J) Val Sugana; (K) Cima d'Arzon-Cima d'Asta; (L) Agordo.




Fig. 2 Geological sketch map of the Comelico area (simplified from: BIANCHI et al., 1930; CASTIGLIONI et al., 1940; CANTELLI et al., 1971; MELI, 1994; HEINISCH et al., 2000). Sampling areas represented by stars: Monte Elmo (ME), N 46°43'08", E 12°24'27"; Monte Cavallino (MC), N 46°40'06", E 12°31'12". P. L. = Periadriatic Line.

ically magmatic embayment. Chlorite + Fe-Ti oxide pseudomorphs point to the occurrence of biotite phenocrysts in the volcanic assemblage. Garnet relics, which are thought to be restitic (MELI, 1998), are often present in MC, while their occurrence has not been detected in ME.

The metamorphic assemblage of Comelico porphyroids is consistent with the Variscan zoneography described by MAZZOLI and SASSI (1988): only chlorite is present as index mineral; other metamorphic minerals are quartz, albite, K-feldspar, muscovite and ilmenite.

4. Zircon typology

Zircon concentrates were obtained using standard techniques; the typology characterization was done following the classification scheme of Pupin (1980). Accordingly, three populations have been recognized in both samples, as shown in figure 3.

Population 37A: clear and colourless crystals, long prismatic, containing few inclusions; no visible cores. The typology distribution clusters around subtype S1.

Population 37B: clear and colourless crystals, short prismatic, with few inclusions; visible cores. The typology distribution clusters around subtype S7.

Population 37C: clear to slightly turbid, pale pink crystals, short prismatic, with few inclusions; visible cores. The typology distribution clusters around subtype S12.

Population 63A: clear and colourless crystals, long prismatic, containing few inclusions; no visible cores. The typology distribution clusters around subtype S1.

Population 63B: clear to turbid, pale pink crystals, short prismatic, with few inclusions; visible cores. The typology distribution clusters around subtype S12.

Population 63C: rounded, slightly elongated crystals, turbid; strongly metamictic. They cannot be classified typologically.

Internal features like cores and inclusions refer to observations from optical microscopy. A cathodoluminescence image performed on a crystal from population 63B confirmed the occurrence of cores in the less elongated crystals (Fig. 4).

Populations 37A and 63A are thought to represent true magmatic products, owing to the elongation, the clearness, the absence of appreciable inclusions and the S1 clustering (PUPIN, 1980). Core-bearing populations 37B, 37C and 63B represent mixtures of inherited crystals and magmatic overgrowths. Hubich and Loeschke (1993) in an extensive cathodoluminescence study of zircons coming from the same area, showed that cores are always present in crystals of the same shape. Population 63C is rounded and strongly metamictic: zircons belonging to this group are thought to be detrital crystals which have been incorporated either into the source or into the melt from wallrocks. Due to their presumably high lattice damages, they have not been analyzed.

5. ²⁰⁶Pb/²³⁸U, ²⁰⁷Pb/²³⁵U, and ²⁰⁷Pb*/²⁰⁶Pb* data

5.1. ANALYTICAL TECHNIQUES

Single grain evaporation Pb–Pb analyses and multigrain conventional U–Pb zircon dating were performed on a Finnigan MAT 262 multicollector mass spectrometer; in both conventional and evaporation runs we employed a secondary electron multiplier (SEM)-ion counter detection system. Pb fractionation for the SEM-ion counter is in the range of 0.200 \pm 0.087% per mass unit.




Fig. 3 SEM photomicrographs of representative zircon crystals form the populations found in samples ME37 and MC63: A = 37A; B = 37B; C = 37C; D = 63A; E = 63B; F = 63C. A, B, C, E, F are SE images; D is a BSE image.

The U-Pb multigrain analyses followed the procedures described by Krogh (1973) and Parrish et al. (1987). All zircon populations were mechanically abraded employing an air abrasion technique (Krogh, 1982). Elemental ratios were determined using a mixed ²⁰⁵Pb–²³³U–²³⁵U spike; U and Pb concentrations were not determined by ID as sample weighing was not possible; a rough

estimate of U concentrations is given in table 1. Pb fractionation was corrected using NBS 981 and NBS 982 standard measurements (0.1070 \pm 0.046% per mass unit). The total procedural blank for all fractions was 12 pg for Pb and 0.1 pg for U.

Single zircon evaporation dating followed modified procedures originally described by KOBER (1987); full details about the applied tech-




Fig. 4 CL image of a core-bearing zircon crystal belonging to population 63B; field of view: $180 \times 120 \mu m$.

nique are summarized by KLÖTZLI (1997). ²⁰⁷Pb/ ²⁰⁶Pb and ²⁰⁸Pb/²⁰⁶Pb ratios were corrected by factors derived from NBS SRM 982 standard measurements. Only high temperature steps (> 1350 °C) with ²⁰⁶Pb/²⁰⁴Pb > 5000 were used for age calculations, and no common lead correction has been applied.

Age calculation and error statistics were made using a modified version of "ROCKAGE 3" from the Geological Survey of Canada and "ISO-PLOT" of LUDWIG (1992); decay constants are those recommended by STEIGER and JÄGER (1977).

5.2. RESULTS

The conventional U/Pb method has been applied to few crystals belonging to populations 37A, 37B, 37C, 63A, 63B: the results are reported in table 1 and in figure 5. The error ellipses of 37A and 63A overlap the concordia line; the weighted average of ²⁰⁷Pb/²⁰⁶Pb, ²⁰⁷Pb/²³⁵U, and ²⁰⁶Pb/²³⁸U ages is 479 ± 8 and 485 ± 8 Ma for ME63 and ME37, respectively. Other populations are discordant; 37B and 37C are undistinguishable within the error limits, and 63B shows a significant departure from the rest of the data points.

Evaporation technique was not suitable for lead measurements on elongated zircons 37A and 63A. As the crystals were of the highest quality, a missing silica gel effect is postulated, due to a shift of silica evaporation towards higher temperatures (KLÖTZLI-CHOWANETZ et al., 1997). Therefore, this technique has been applied only to one corebearing crystal from population 37B; data are summarized in table 2 and reported in the frequency histogram of figure 6. The high T steps yield measurements which cluster around an age of 681 ± 7 Ma; no age zoning can be detected within the crystal. Fig. 7 shows spatial variations of ²⁰⁸Pb*/²⁰⁶Pb* towards the crystal interior, not accompanied by a similar variation of ²⁰⁷Pb*/ ²⁰⁶Pb*. As ²⁰⁸Pb* solely descends from the decay of ²³²Th, such features are thought to reveal changing Th/U ratios in different zircon domains, probed at different evaporation temperatures.

Tab. 1 U-Pb data on zircon populations from Comelico porphyroids.

| | Isotopic ratios | | | | | | | |
|------|-----------------|--------------------|---------------|--------------------------------------|--|----------------------|--|--|
| Pop. | 206Pb a 204Pb | 207Pb b | | 206Pb b | ²⁰⁷ Pb ²⁰⁶ Pb | | | |
| 37A | 889 | 0.61308 = | 1.42% | 0.07799 ± 1.40 | 0.05701 ± | .60% 0.3116 | | |
| 37B | 896 | 0.68779 = | 1.28% | 0.08072 ± 1.40 | $0.06179 \pm$ | .58% 0.2884 | | |
| 37C | 903 | 0.69057 | 1.58% | 0.08079 ± 1.64 | $0.06199 \pm$ | .64% 0.2773 | | |
| 63A | 831 | 0.60517 = | 0.90% | 0.07704 ± 0.52 | 2% 0.05696 ± | .35% 0.3998 | | |
| 63B | 722 | 1.51320 = | ± 0.76% | 0.09950 ± 0.84 | % 0.11026 ± | .26% 0.4771 | | |
| 7.3% | N° of Th/U | | Apparent ages | | | | | |
| Pop. | crystals | U ppm ^c | today | $\frac{207 \text{Pb}}{235 \text{U}}$ | $\frac{206 \text{Pb}}{238 \text{U}}$ | $\frac{207}{206}$ Pb | | |
| 37A | 5 | 1500 | 1.00 | 485.5 ± 5.4 | 484.1 ± 6.6 | 491.9 ± 13.1 | | |
| 37B | 7 | 5000 | 0.93 | 531.5 ± 5.4 | 500.5 ± 6.8 | 666.9 ± 12.3 | | |
| 37C | 7 | 5000 | 0.89 | 533.1 ± 6.6 | 500.9 ± 7.8 | 673.7 ± 13.7 | | |
| 63A | 5 | 2000 | 1.28 | 480.5 ± 3.4 | 478.5 ± 2.4 | 490.3 ± 7.8 | | |
| 63B | 4 | 8000 | 1.64 | 935.7 ± 4.6 | 611.6 ± 4.8 | 1803.8 ± 8.4 | | |

Errors on isotopic ratios are 2σ of the mean in %; apparent age errors are 2σ of the mean in Ma.

^b ratios corrected for the common lead, assuming that its isotopic composition is predicted by the two-stage model of STACEY and KRAMERS (1975).

c approximate values.

6. Discussion

6.1. INTERPRETATION OF U-Pb AND Pb-Pb AGES

Zircon crystals which are thought to bear inherited cores are discordant. Hubich and Loeschke (1993) demonstrated the existence of older cores in the interior of the S7 (S12) clustering populations, without any evidence of multiple recycling of the grains. Therefore, a discordance produced by mixing of two components could be a realistic hypothesis; however, applying the integrating conventional U/Pb method, it is not possible to ascertain whether the core component is concordant by itself.

Elongated crystals are concordant; nevertheless, the age sequence ²⁰⁷Pb/²⁰⁶Pb > ²⁰⁷Pb/²³⁵U > ²⁰⁶Pb/²³⁸U is found. This could be due to an underestimation of the ²⁰⁷Pb/²⁰⁴Pb in the common Pb correction, performed employing the Pb evolutionary curve proposed by STACEY and KRAMERS (1975). As the contribution of older crust is dem-

Tab. 2 Evaporation data of a zircon crystal belonging to the population 37B.

| T (°C) | $^{207}Pb/^{206}Pb$ | age | $^{208}Pb/^{206}Pb$ | Th/U |
|--------|---------------------|------------------|---------------------|------|
| 1377 | 0.06223 ± 45 | 681.4 ± 15.1 | 0.12545 ± 67 | 0.40 |
| 1399 | 0.06321 ± 25 | 715.1 ± 8.6 | 0.13029 ± 55 | 0.42 |
| 1420 | 0.06124 ± 36 | 646.6 ± 12.9 | 0.12066 ± 57 | 0.39 |
| 1441 | 0.06228 ± 42 | 682.5 ± 14.4 | 0.13033 ± 57 | 0.42 |
| Total | 0.06224 ± 21 | 681 ± 7 | 0.12685 ± 57 | 0.41 |

Reported ages are propagated weighted mean values calculated from at least 20 measured $^{207}\text{Pb}/^{206}\text{Pb}$ ratios. Errors on isotopic ratios and ages are 2σ of the mean.




Fig. 5 Concordia diagram of U–Pb zircon data from the Comelico porphyroids.

onstrated by the core-bearing populations 37B, 37C and 63B, the inheritance of small amounts of lead with anomalously high ²⁰⁷Pb/²⁰⁴Pb seems probable; unfortunately, the Variscan metamorphic reequilibration of suitable phases prevented more precise common Pb correction.

The upper intercepts give ages of 2628 ± 760 and 3280 ± 51 Ma, but their interpretation is not straightforward, as the data do not allow us to establish whether they represent a discordia generated by a mixing of two concordant ages, or the inherited component is discordant by itself, causing a multiple discordance. In the latter case, upper intercepts do not represent any geological event; however, the 207 Pb/ 206 Pb datum (Tab. 1) constrains the minimum age of the oldest basement rocks in the region at about 1800 Ma.




Fig. 6 207 Pb*/ 206 Pb* age distribution from 37B single grain evaporation measurement.




Fig. 7 Plot of block number versus $^{207}\text{Pb*}/^{206}\text{Pb*}$ and $^{208}\text{Pb*}/^{206}\text{Pb*}$. Block number gives progress of evaporation with steps of increasing T. Each block represent the mean of 20 mass scans. Assigned errors of averaged ratios are 2σ of the mean.

In the light of U–Pb data, the ²⁰⁷Pb*/²⁰⁶Pb* age obtained on a single zircon from population 37B does not match with concordant ages of 37A and 63A populations. The lack of ages around 480–490 Ma in the low T steps are probably related again to a missing silica gel effect in the outer part of the zircon, as happened for 37A and 63A. The evaporation spectrum reveals also a Th/U zoning of the inherited core (Fig. 7), which points towards a magmatic signature of this domain. Therefore, a partial reworking of igneous material by the anatectic melts is suggested.

6.2. REGIONAL IMPLICATIONS

As demonstrated by zircon populations, some differences between ME37 and MC63 are evident: in the former, rounded metamictic zircons are absent, and the inherited cores show different isotopic compositions. Moreover, several restitic garnets have been observed only in the latter, pointing to a medium-scale heterogeneity in the crustal source of the Ordovician volcanics.

The concordant ages of both 37A and 63A constrain the Comelico volcanism within the Arenig (HARLAND et al., 1989). Thus, the new radiometric ages are not contradictory with the Caledonian hypothesis held by Borsi et al. (1975) and EBNER et al. (1987), as the metamorphic event they found in the Eastern Alps is slightly older than the emplacement of acidic volcanics. Moreover, a time resemblance with the Caledonides s.s. is testified by the occurrence of igneous products aged 463-475 Ma in the Connemara region, Ireland (FRIEDRICH et al., 1999a; KINNY et al., 1999). In particular, a set of anatectic products, which include migmatites and granitic pegmatites, has been dated at 467-468 Ma (FRIE-DRICH et al., 1999b). However, the attribution of the Ordovician volcanism to the Caledonian event s.s. is still problematic, as other geologic elements are still lacking.

The extensional tectonic setting hypopthesized on geochemical and petrological grounds (Meli, 1998) fits that described for the Upper Cambrian–Lower Ordovician time span in Central-Western Europe (Pin, 1991). The occurrence of an Early–Middle Ordovician extensional tectonic setting was proposed also for the French Central Massif (Ploquin and Stussi, 1994) and the Armorican Massif (Le Corre, 1994), where a metavolcanic body interlayered with syn-sedimentary beds has been dated by means of U/Pb methods at 466 ± 1 Ma (Bonjour et al., 1988). Therefore, the porphyroids can represent episodes of crustal anatexis and widespread acidic,

peraluminous volcanism, which developed in a thinned continental crust characterized by increased basal heat flow.

The long distance correlations between Austroalpine and Southalpine volcanic events (SASSI and ZIRPOLI, 1989) can be reviewed in the light of the new U/Pb data on Comelico porphyroids. In the Eisenerz metamorphic succession ("D" in Fig. 1), two different porphyroid bodies have been identified (NEUBAUER and SASSI, 1993): the age of the older layer is Lower Ordovician, whereas the younger one is located in the Upper Ordovician, probably being a part of the main pre-Variscan Austroalpine acidic volcanic episode, named "Blasseneck event" from the type locality ("C" in Fig. 1). This latter is younger than Lower Caradocian (FLAJS and SCHÖNLAUB 1976); therefore a long distance chronostratigraphic correlation with Comelico porphyroids does not hold, as a time gap of at least 35 Ma exists (PALMER, 1983; ODIN, 1986; HARLAND et al., 1989; GRADSTEIN and OGG, 1996). Comelico metavolcanics are better correlated with the lower porphyroid level, which underlies the Blasseneck ignimbrites and whose age has been referred to Lower Ordovician. The new radiometric ages suggest that the space-time distribution of the pre-Variscan volcanics is more complex than previously believed, and that pre-Variscan volcanism cannot be considered synchronous throughout the Southalpine and Austroalpine domains. However, within the Southalpine basement, Comelico radiometric ages fit the chronobiostratigraphy of Agordo zone ("L" in Fig. 1), whose metamorphic succession has been correlated with the Comelico units (ZANFERRARI, 1985): the acritarch dated Col di Foglia Fm., which underlies the porphyroids, is older than Arenig (KALVACHEVA et al., 1986). Therefore, at least within the easternmost part of the Southalpine basement, the acidic metavolcanic layers occurring in the Volcano-Sedimentary Complex should have been formed within the same time span. Fig. 8 summarizes the timing of the magmatic and metamorphic events occurred during Paleozoic in the Eastern Southalpine basement, taking into account also these new geochronological data.

7. Conclusions

The U/Pb ages presented in this contribution represent the first geochronologic data of an Ordovician magmatic event in the Southalpine to the east of the Giudicarie Line: other Ordovician (450–480 Ma) magmatic bodies (tonalitic and granitic orthogneisses) are known only in the West-




Fig. 8 Sketch of the timing of Paleozoic magmatic and metamorphic events in the Southalpine domain to the east of the Giudicarie Line (modified after Sassi and Zirpoli, 1989). Sources of geochronological and biostratigraphic data: 1 = Del Moro et al., 1980 (Val di Funes); 2 = Del Moro et al., 1984 (Brunico); 3 = Brondi et al., 1970 (Bolzano); 4 = D'Amico et al., 1980 (Bolzano); 5 = Borsi et al., 1972 (Bressanone); 6 = Borsi et al., 1974 (Cima d'Asta); 7 = Del Moro and Visonà, 1982 (Bressanone); 8 = Barth et al., 1994 (Bolzano and Cima d'Asta); 9 = Kalvacheva et al., 1986 (Agordo); 10 = Meli, 1994 (Val Sarentino and Val Sugana); 11 = this work. Geologic time scale taken from Harland et al. (1989). Abbreviations as in section 2.

ern Southalpine, within the "Scisti dei Laghi" metamorphic unit (SCHALTEGGER and GEBAUER, 1999, and references therein). The age of volcanism in Comelico is constrained within the Lower Ordovician: concordant ages of 479 ± 8 and 485 ± 8 Ma result for MC and ME, respectively. Thus, the "Caledonian" hypothesis is now more substantiated than the alternative "Cadomian" hypothesis. The pre-Variscan volcanism developed earlier than Caradocian—Ashgillian, as previously believed: its correct chronostratigraphic position is within the Arenig. At the present state of know-

ledge, no geological event can be reasonably attributed to the upper intercept ages; however, a minimum age of 1800 Ma is inferred for the oldest component of the source material of the anatectic melts. The differences in isotopic composition between M. Elmo and M. Cavallino zircon populations could be due to a medium-scale heterogeneity within the crustal source of the volcanics, testified also by some petrographic features of ME and MC samples.

Within the easternmost part of the Southalpine basement, the Ordovician volcanism can be considered synchronous, as the new U/Pb ages of Comelico porphyroids agree with the ages inferred for Agordo metavolcanics on biostratigraphic grounds. The Southalpine porphyroids were also correlated with Austroalpine metavolcanics outcropping in the Northern Greywacke Zone; however, the more complex stratigraphy of Austroalpine successions does not allow simple correlations with the Southalpine domain: the main Austroalpine ignimbritic eruption ("Blasseneck event") is younger than Comelico volcanism, whereas an older metavolcanic layer occurring in the Eisenerz area is approximately coeval. Therefore, these new data do not rule out the possibility of considering Southalpine and Austroalpine metavolcanics as being part of a unique cycle of magma generation; however, more detailed geochronology is needed to correctly describe the Ordovician volcanic activity in the Eastern Alps, as sporadic data tend to oversimplify the picture.

Acknowledgements

This research was supported by University of Padova (MURST) and National Council of Research (C.N.R.). We are grateful to Prof. F.P. Sassi, Prof. G. Zirpoli, Prof. G. Serri and Prof. E. Morlotti for their stimulating suggestions. Prof. R.H. Steiger is kindly acknowledged for his comments on a previous version of the manuscript; careful review by Prof. J.D. Kramers and an anonymous reviewer greatly improved the manuscript. Many thanks to T. Ntaflos for the performance of SEM images, to Dr. G. Salviati and N. Armani (MASPEC Institute, Parma) for the performance of CL images and to W. McGregor for the correction of English text.

References

BARTH, S., OBERLI, F. and MEIER, M. (1994): Th–Pb versus U–Pb isotope systematics in allanite from co-genetic rhyolite and granodiorite: implications for geochronology. Earth Planet. Sci. Lett., 124, 149–159.

Bellieni, G. and Sassi, F.P. (1981): New chemical data and a review on the South-Alpine "Pre-Hercynian Rhyolitic Plateau" in the Eastern Alps. IGCP Project n° 5, Newsletter, 3, 22–27.

BIANCHI, A., DAL PIAZ, G. and MERLA, G. (1930): Carta Geologica delle Tre Venezie, F° 4b - Monguelfo,

1:100000. Ufficio Idrografico del Real Magistrato

alle acque, Venezia.

BÖGEL, H., MORTEANI, G., SASSI, F.P., SATIR, M. and SCHMIDT, K. (1979): The Hercynian and pre-Hercynian development of the Eastern Alps. Report on a meeting. N. Jb. Geol. Paläont. Abh., 159, 87–112.

BONJOUR, J.L., PEUCAT, J.J., CHAUVEL, J.J., PARIS, F. and CORNICHET, J. (1988): U-Pb zircon dating of the Early Paleozoic (Arenigian) transgression in Western Brittany (France): a new constraint for the Lower Paleozoic time-scale. Chem. Geol., 72, 329–336. Borsi, S., D'Amico, C. and Del Moro, A. (1974): Studio

radiometrico delle rocce intrusive del massiccio di Cima d'Asta (Trentino). Mem. Soc. Geol. It., 12,

549-571.

Borsi, S., Del Moro, A. and Ferrara, G. (1972): Età radiometriche delle rocce intrusive del massiccio di Bressanone - Ivigna - Monte Croce (Alto Adige). Boll. Soc. Geol. It., 91, 387–406.

Borsi, S., Del Moro, A., Sassi, F.P., Zanferrari, A. and ZIRPOLI, G. (1975): Evento "caledoniano" nelle Alpi? Risposta ad un intervento critico. Boll. Soc.

Geol. It., 94, 275-280.

BORSI, S., DEL MORO, A., SASSI, F.P. and ZIRPOLI, G. (1973): Metamorphic evolution of the Austridic rocks to the south of the Tauern Window (Eastern Alps): radiometric and geopetrologic data. Mem. Soc. Geol. It., 12, 549-571.

Brondi, A., Ghezzo, C., Guasparri, G., Ricci, C.A. and SABATINI, G. (1970): Le vulcaniti paleozoiche nell'area settentrionale del complesso effusivo atesino. Atti Soc. Tosc. Sci. Nat., Mem., s. A., 77, 157–200.

CALLEGARI, E. and DE PIERI, R. (1966): Crystallographical observations on some chess-board albites.

Schweiz. Miner. Petrogr. Mitt., 47, 99-110.

CANTELLI, C., CARLONI, G.C., CASTELLARIN, A., CER-ETTI, E., COLANTONI, P., CREMONINI, G., ELMI, C., Frascari, F., Monesi, A., Pisa, G., Rabbi, E., Toma-DIN, L. and VAI, G.B. (1971): Carta Geologica d'Italia, F° 4c-13 - Monte Cavallino/Ampezzo, 1:100000. Servizio Geologico d'Italia, Roma.

CARDIN, A., PISANI, F., SASSI, F.P., VISONA', D. and ZIRPO-LI, G. (1985): Le metapeliti e le anfiboliti del basamento sudalpino delle Alpi Sarentine (Alpi occidentali). Mem. Sci. Geol. Univ. Padova, 37, 379-406.

CASTIGLIONI, B., LEONARDI, P., MERLA, G., TREVISAN, L. and Zenari, S. (1940): Carta Geologica delle Tre Venezie, F° 12 – Pieve di Cadore, 1:100000. Ufficio Idrografico del Real Magistrato alle acque, Venezia.

D'AMICO, C., DEL MORO, A., FREDDO, A. and PARDINI, G. (1980): Studio radiometrico delle ignimbriti riolitiche atesine, gruppo superiore. Rend. Soc. It. Min-

eral. Petrol., 36, 703-716.

DEL MORO, A. and VISONA, D. (1982): The epiplutonic Hercynian complex of Bressanone (Brixen, Eastern Alps, Italy). Petrologic and radiometric data. N. Jb.

Miner. Abh., 145, 66–85.

DEL MORO, A., SASSI, F.P. and ZIRPOLI, G. (1980): Preliminary results on the radiometric age of the Hercynian metamorphism in the South-Alpine basement of the Eastern Alps. N. Jb. Geol. Paläont. Mh., 707-718.

DEL MORO, A., SASSI, F.P. and ZIRPOLI, G. (1984): Acidic gneisses from Plan De Corones area, and chronological data on South-Alpine basement in Pusteria (Eastern alps). Mem. Sci. Geol. Univ. Padova, 36, 403-412.

EBNER, F., NEUBAUER, F. and STATTEGGER, K. (1987): The Caledonian event in the Eastern Alps: a review. In: FLÜGEL, H.W., SASSI, F.P. and GRECULA, P. (eds): Pre-Variscan and Variscan events in the AlpineMediterranean mountain belts, IGCP Project no 5, Bratislava, 169-182.

FLAJS, G. and SCHÖNLAUB, H.P. (1976): Die biostratigraphische Gliederung des Altpalaozoikums am Polster bei Eisenerz. Verh. Geol. B.-A., 257-303.

FREY, M., DESMONS, J. and NEUBAUER, F. (eds)(1999): The new metamorphic map of the Alps. Schweiz.

Mineral. Petrogr. Mitt., 79(1)

FRIEDRICH, A.M., BOWRING, S.A., MARTIN, M.W. and HODGES, K.V. (1999a): Short-lived continental magmatic arc at Connemara, Western Irish Caledonides; implications for the age of the Grampian orogeny. Geology, 27, 27-30.

FRIEDRICH, A.M., HODGES, K.V., BOWRING, S.A. and MARTIN, M.W. (1999b): Geochronological constraints on the magmatic, metamorphic and thermal evolution of the Connemara Caledonides, Western Ireland. J. Geol. Soc. of London, 156, 1217–1230.

Gradstein, F.M. and Ogg, J.G. (1996): A Phanerozoic Time Scale. Episodes, 19/1–2, 3–5.

HANDLER, R., DALLMEYER, R.D. and NEUBAUER, F. (1997): 40 Ar/39 Ar ages of detrital white mica from Upper Austroalpine units in the Eastern Alps, Austria: evidence for Cadomian and contrasting Varis-

can sources. Geol. Rundsch., 86, 69–80. Harland, W.B., Armstrong, R.L., Cox, A.V., Craig, L.E., SMITH, A.G. and SMITH, D.G. (1989): A geologic time scale. Cambridge University Press, 263 pp.

Heinisch, H. (1981): Zum Ordovizischen "Porphyroid"-Vulkanismus der Ost- und Südalpen, Stratigraphie, Petrographie, Geochemie. Jahrb. Geol. B.-A., 124,

Heinisch, H. (1988): Hinweise auf die Existenz eines passiven Kontinentalrandes im Altpaläozoikum der Nördlichen Grauwackenzone – Ostalpen. Schweiz. Mineral. Petrogr. Mitt., 68, 407–418.

HEINISCH, H. and SCHMIDT, K. (1976): Zur Kaledonischen Orogenese in den Ostalpen. Geol. Rund-

sch., 65, 459-482.

HEINISCH, H., VAN HUSEN, D. and SCHÖNLAUB, H.P. (2000): Geologische Karte der Republik Österreich, 195 – Sillian, 1:50000. Geologischen Bundesanstalt,

HUBICH, V.D. and LOESCHKE, J. (1993): Der Comelico Porphyroid der westlichen Karnischen Alpen (Österreich/Italien): Stratigraphie, Petrographie, Geochemie und Zirkonmorphologie. Jahrb. Geol. B.-A., 136, 351-374.

KALVACHEVA, R., SASSI, F.P. and ZANFERRARI, A. (1986): Acritarch evidence for the Cambrian age of phyllites in the Agordo area (South-Alpine Basement of Eastern Alps, Italy). Rev. Paleobot. Palinol., 48, 311-326.

KINNY, P.D., FRIEND, C.R.L., STRACHAN, R.A., WATT, G.R. and BURNS, I.M. (1999): U-Pb geochronology of regional migmatites in East Sutherland, Scotland; evidence for crustal melting during the Caledonian orogeny. J. Geol. Soc. of London, 156, 1143–1152

KLÖTZLI, U.S. (1997): Single zircon evaporation thermal ionisation mass spectrometry: method and proce-

dures. Analyst, 122, 1239-1248.

KLÖTZLI-CHOWANETZ, E., KLÖTZLI, U.S. and KOLLER, F. (1997). Lower Ordovician migmatisation in the Ötztal crystalline basement (Eastern Alps, Austria): linking U-Pb and Pb-Pb dating with zircon morphology. Schweiz. Mineral. Petrogr. Mitt., 77, 315–324.

KOBER, B. (1987): Single zircon evaporation combined with Pb* emitter-bedding for ²⁰⁷Pb/²⁰⁶Pb age investigations using thermal ion mass spectrometry, and implications to zirconology. Contrib. Mineral. Petrol., 96, 63–71.

Krogh, T.E. (1973): A low contamination method for hydrothermal decomposition of zircon and extraction of U and Pb for isotopic age determinations. Geochim. Cosmochim. Acta, 37, 485-494.

Krogh, T.E. (1982): Improved accuracy of U-Pb zircon ages by the creation of more concordant systems using an air abrasion technique. Geochim. Cosmo-

chim. Acta, 46, 637-649.

LE CORRE, C. (1994): Early tectonic events (Ordovician). In: KEPPIE, J.D. (ed.): Pre-Mesozoic geology in France and related areas, Springer-Verlag, Berlin,

LOESCHKE, J. (1989): Lower Paleozoic volcanism of the Eastern Alps and its geodynamic implications. Geol.

Rundsch., 78, 599–616.

LUDWIG, K.R. (1992): Isoplot: A plotting and regression program for radiogenic-isotope data. USGS, Open-

file report 91–445.

MAZZOLI, C. and SASSI, R. (1988): Caratteri del metamorfismo ercinico nella fillade sudalpina ad ovest di Bressanone. Mem. Sci. Geol. Univ. Padova, 40, 295–314.

MAZZOLI, C. and SASSI, R. (1992): New chemical data on the Upper Ordovician acidic plutonism of the Eastern Alps. In: CARMIGNANI, L. and SASSI, F.P. (eds): Contributions to the geology of Italy, with special regard to the Paleozoic basement, IGCP no 276, Newsletter, 263-277.

MELI, S. (1994): I porfiroidi sudalpini delle Alpi Orientali: studio petrografico e geochimico. Ph.D. Thesis,

University of Padova, Italy, 298 pp.

MELI, S. (1998): Pre-Variscan volcanic activity in the Eastern Alps: the Southalpine porphyroids. Schweiz.

Miner. Petrogr. Mitt. 78, 417–432

- NEUBAUER, F. and Sassi, F.P. (1993): The Austroalpine quartzphyllites and related Paleozoic Formations. In: von RAUMER, J.F. and NEUBAUER, F. (eds): Pre-Mesozoic Geology in the Alps, Springer-Verlag, 423-439
- Neubauer, F., Hoinkes, G., Sassi, F.P., Handler, R., HÖCK, V., KOLLER, F. and FRANK, W. (1999): Pre-Alpine metamorphism of the Eastern Alps. Schweiz. Miner. Petrogr. Mitt. 79, 41–62.

ODIN, G.S. (1986): Recent advances in Phanerozoic Time-Scale calibration. Chem. Geol. (Isot. Geosci.

Sect.), 59, 103-110.

PALMER, A.R. (1983): The decade of North America Geology. 1983 Geologic Time Scale. Geology, 11,

PARRISH, R.R., RODDICK, J.C., LOVERIDGE, W.D. and SULLIVAN, R.W. (1987): Uranium-lead analytical techniques at the geochronology laboratory, Geological Survey of Canada. Geol. Surv. of Canada Paper 87, 3–7.

PIN, C. (1991). Central-Western Europe: major stages of development during Precambrian and Paleozoic times. In: DALLMEYER, R.D. and LECORCHE, J.P. (eds): The West African orogens and Circum-Atlantic correlatives, Springer-Verlag, Berlin, 295–306.

PLOQUIN, A. and STUSSI, J.M. (1994): Felsic plutonism and volcanism in the Massif Central. In: KEPPIE J.D. (ed): Pre-Mesozoic geology in France and related areas. Springer-Verlag, Berlin, 363–378.

Pupin, J.P. (1980): Zircon and granite petrology. Contrib. Mineral. Petrol., 73, 207-220.

SASSI, F.P. and SCHMIDT, L. (1982): The "Caledonian" event in the Eastern Alps: guidelines for its geodynamic interpretation. IGCP Project no 5, Newsletter, 4, 78-81.

SASSI, F.P. and SPIESS, R. (1993): The South-Alpine metamorphic basement in the Eastern Alps. In: von RAUMER, J.F. and NEUBAUER, F. (eds): Pre-Mesozoic geology in the Alps, Springer-Verlag, Berlin, 599-607

SASSI, F.P. and ZANFERRARI, A. (1972): Il significato geologico del complesso del Turntaler (Pusteria), con particolare riguardo alla successione di eventi metamorfici prealpini nel basamento austridico delle Alpi orientali. Boll. Soc. Geol. It., 91, 533-557.

SASSI, F.P. and ZIRPOLI, G. (1968): Il basamento cristallino di Recoaro. Mem. Soc. Geol. It., 7, 227-245.

SASSI, F.P. and ZIRPOLI, G. (1989): The lithostratigraphic sequence in the Southalpine basement of the Eastern Alps. Rend. Soc. Geol. It., 12, 397-402

SASSI, F.P., CAVAZZINI, G. and VISONA, D. (1985): Radiometric geochronology in the Eastern Alps: results and problems. Rend. Soc. It. Mineral. Petrol., 40, 187 - 224.

SASSI, F.P., ZANFERRARI, A. and ZIRPOLI, G. (1974): Some considerations on the South-Alpine basement of the Eastern Alps. N. Jb. Geol. Palaont. Mh., 609-

SASSI, F.P., ZANFERRARI, A. and ZIRPOLI, G. (1979): Nuovi dati sulla stratigrafia e i caratteri chimicopetrografici dei porfiroidi del Comelico (Alpi Ori-

entali). Mem. Soc. Geol. It., 20, 425–434.

SASSI, F.P., ZANFERRARI, A. and ZIRPOLI, G. (1987): The Caledonian event in the Eastern Alps: a review. In: FLUGEL, H.W., SASSI, F.P. and GRECULA, P. (eds): Pre-Variscan and Variscan events in the Alpine-Mediterranean mountain belts, IGCP Project no 5, Bratislava, 431-434.

SCHALTEGGER, U. and GEBAUER, D. (1999): Pre-Alpine geochronology of the Central, Western and Southern Alps. Schweiz. Miner. Petrogr. Mitt. 79, 79–87.

STACEY, J.S. and KRAMERS, J.D. (1975): Approximation of terrestrial lead isotope evolution by a two-stage model. Earth Planet. Sci. Lett., 26, 207–221.

STEIGER, R.H. and JÄGER, E. (1977): Subcommission on geochronology: convention on the use of decay constants in geo- and cosmochronology. Earth Planet. Sci. Lett., 36, 359-362.

VAI, G.B. (1975): Evento "caledoniano" nelle Alpi? Discussione sui rapporti tra radiocronometria e strati-

grafia. Boll. Soc. Geol. It., 94, 269-273.

VISONA, D. and ZANFERRARI, A. (1985): Rifting ercinico nel basamento sudalpino delle Alpi Orientali. In: Cocozza, T. and Ricci, C.A. (eds): Evoluzione stratigrafica, tettonica, metamorfica e magmatica del Paleozoico italiano, Siena, Dip. Sc. Terra, 111.

Zanferrari, A. (1985): Stratigrafia e tettonica del basamento sudalpino delle Alpi Orientali. In: Cocozza, T. and Ricci, C.A. (eds): Evoluzione stratigrafica, tettonica, metamorfica e magmatica del Paleozoico italiano, Siena, Dip. Sc. Terra, 112-113.

Manuscript received July 7, 2000; revision accepted May 19, 2001.