

# Rare earth element concentrations in mafics from the Kizil Da ophiolite (Hatay, Turkey)

Autor(en): **Delaloye, M. / Piskin, Ö. / Voldet, P.**

Objektyp: **Article**

Zeitschrift: **Schweizerische mineralogische und petrographische Mitteilungen  
= Bulletin suisse de minéralogie et pétrographie**

Band (Jahr): **59 (1979)**

Heft 1-2

PDF erstellt am: **24.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-46043>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## **Rare Earth Element Concentrations in Mafics from the Kızıl Dağ Ophiolite (Hatay, Turkey)**

by *Delaloye, M.<sup>1)</sup>, Piskin, Ö.<sup>2)</sup>, Voldet, P.<sup>1)</sup>, Vuagnat, M.<sup>1)</sup> and Wagner, J.-J.<sup>1)</sup>*

### **Abstract**

Abundances of rare earth elements (REE) from the mafic sequence of the Kızıl Dağ – Hatay ophiolite have been measured by neutron activation technique for short lived isotopes. The results for twelve samples from the gabbroic part show a general REE pattern with a depletion in the light REE (LREE) indicating that these rocks were derived from a uniform LREE depleted asthenosphere source (SCHILLING, 1975) at a spreading zone. The REE abundances vary by a factor 6 to 10 between the cumulate gabbros from the ultramafic – transition zone and the high level microgabbros from screens in the dyke complex.

### **Özet**

Kızıl Dağ – Hatay ofiyolitini teşkil eden mafik kayalardaki lantanit elementlerin miktarları nötron aktivasyon metodu ve kısa periyodlu izotopların ölçümüyle tayin edildi. Gabrolara ait 12 numuneden elde edilen neticeler hafif lantanitler için genel bir azalma göstermektedir. Bu da, bu kayaların kaynağının hafif lantanitler için homojen bir azalmaya uğranış genişleme otanındaki bir astenosfer olduğunu gösterir (SCHILLING, 1975). Filon kompleksinde ki ektanlara ait üst seviye mikrogabroları, ultramafik-mafik geçişme sahasındaki gabrolara kıysla 6 ile 10 defa daha fazla miktarda lantanit elementleri ihtiva etmektedirler.

### **1. INTRODUCTION**

Ophiolitic massifs are a major center of interest because they are thought to be pieces of an ancient oceanic crust formed at a spreading axis (GASS, 1968, DEWEY et al., 1973, MOORES et al., 1974). However, it is very difficult to determine if the ophiolites were part of a mid-ocean ridge or part of a back arc basin. Although we cannot actually decide from the geochemistry of the REE to which paleoenvironment the Kızıl Dağ ophiolite belongs we can use the REE

---

<sup>1)</sup> Mineralogy Department, The University, CH-1211 Geneva 4, Switzerland.

<sup>2)</sup> Ege University, Izmir, Turkey.

concentrations to elucidate the processes related to the formation of magma from the upper mantle. This study may help to characterize ophiolites from different settings.

A systematic investigation of the REE of the different members of the Kızıl Dagh ophiolite is in progress and in the present paper we discuss our data on the gabbroic member.

## 2. GEOLOGICAL SETTING OF THE KIZIL DAĞ

The Kızıl Dagh ophiolite massif represents the S-W end of the Amanos Range close to the border between Turkey and Syria. The massif is cut abruptly by a major fault running N-S which forms the shoreline. Rising to 1800 m above sea level, a very impressive outcrop is visible along the coast. The Kızıl Dagh was first mapped by DUBERTRET (1953) and described as a single huge submarine flow of Upper-Cretaceous age that has differentiated by gravity, with transition from fine grained basaltic rocks to coarse-grained ultramafics. The massif was later on interpreted as an ocean floor lithosphere by VUAGNAT et al. (1967). They recognized particularly the presence of a sheeted complex of dykes with chilled edges against screens of gabbros. Further studies have been done by ÇOĞULU (1973, 1974, 1976) and PARROT (1973).

The ophiolite suite can be characterized as follows:

- a) ultramafic tectonites (harzburgites and dunites) form the center of the massif,
- b) ultramafic cumulates formed by fractional crystallization of a magma produced by partial melt of the mantle (JACKSON, 1971). They outcrop mainly towards the SE edge of the massif,
- c) mafic rocks like olivine gabbros or amphibole gabbros showing a variety of banding and flow structures,
- d) a five kilometers wide sheeted complex formed of dolerites dykes with few gabbros screens,
- e) pillow lavas of basaltic composition outcrop to the NE.

The contacts between these members have often a tectonic character. One may also recall that a previous paleomagnetic study (WAGNER et al., 1976) did not show any magnetic field reversals; measured ages (DELALOYE et al., 1977) show that the Kızıl Dagh ophiolite seems to have acquired its sea floor magnetization during a normal polarity geomagnetic period.

## 3. ANALYTICAL METHODS

The determination of REE concentrations was made by neutron activation followed by high-resolution X-Ray spectrometry or  $\gamma$ -spectrometry.

For a number of years neutron activation analysis has been used for the determination of rare-earth elements in rocks. We propose a method for the simultaneous determination of eight to ten REE in the mafics of an ophiolite (gabbros, dolerites, basalts). Before neutron activation, the RE elements must be separated from the other elements present in rocks so as to obtain the same matrix for rocks and standards for the measurements of activity. This separation is done by gravimetry and column chromatography; the procedure is described by VOLDET and HAERDI (1976). The standards were prepared by precipitation of the oxalates of mixtures of RE elements in proportions suitable for mafic rocks.

Samples and standards were first irradiated for 30 min. in the AGN-201-P reactor of Geneva University at a thermal neutron flux of  $10^9 \text{n.cm}^{-2}.\text{s}^{-1}$  and Eu and Dy were determined. Then, the same samples and standards were irradiated for 10 hours in the Diorit reactor of the Swiss Federal Institute for Reactor Research, Würenlingen at a thermal neutron flux of about  $2,5.10^{12} \text{n.cm}^{-2}.\text{s}^{-1}$  and the following REE were determined;

- a) by high-resolution X-Ray spectrometry (crystal Ge):  
 $^{152\text{m}}\text{Eu}$  ( $t_{1/2}$ , 9.3 h),  $^{153}\text{Sm}$  ( $t_{1/2}$ , 46.5 h),  $^{166}\text{Ho}$  ( $t_{1/2}$ , 1.12 d),  $^{177}\text{Lu}$  ( $t_{1/2}$ , 6.75 d),  $^{166}\text{Yb}$  ( $t_{1/2}$ , 30.6 d),  $^{147}\text{Nd}$  ( $t_{1/2}$ , 11.1 d) and  $^{143}\text{Ce}$  ( $t_{1/2}$ , 33 h);
- b) by  $\gamma$ -spectrometry (crystal Ge[Li]):  
 $\text{Gd}$ ( $^{161}\text{Tb}$ ) ( $t_{1/2}$ , 7.2 d),  $^{177}\text{Lu}$  ( $t_{1/2}$ , 6.75 d),  $^{143}\text{Ce}$  ( $t_{1/2}$ , 33 h) and  $^{140}\text{La}$  ( $t_{1/2}$ , 40.3 h).

These activities are measured after a cooling time of about 36, 60 and 110 hours. Each sample is counted for 500 s with the Ge detector and 60 s with the Ge(Li) detector. For X-Ray spectrometry, the equipment used is described in our work (Loc. cit.), and for  $\gamma$ -spectrometry the following equipment was used:

Ge(Li) detector (Seforad, 50  $\text{cm}^3$ ), preamplifier Seforad SR-100, amplifier Ortec 472 and a 4096 channel Zoomax SEIN analyzer (VOLDET and HAERDI, 1978). The precision (of the order of 5 to 20%) is limited principally by the accuracy of the measurement method. No interference of other elements was observed; this confirms the validity of the separation method.

The method has been tested on several standard rocks; the following results are obtained for the U.S.G.S. standard rock BCR-1:

Element	La	Ce	Sm	Eu	Gd	Dy	Ho	Lu
Value in ppm	23.4	50.9	6.2	1.90	7.2	6.34	1.6	2.4

The values from La to Ho are in agreement with those published by several authors (MENZIES, 1976). One has to be careful with the value of Lu which in our case is 2.4 ppm and normally is given as around 0.6 ppm.

#### 4. RESULTS AND DISCUSSION

Several Eastern Mediterranean ophiolites such as the Troodos, Pindos and Othris have been studied for their REE concentrations (e.g. KAY et al., 1976, MENZIES, 1976). They exhibit a general REE pattern which can be summarized from the Troodos ophiolite:

- a) All rocks have the same trend showing a depletion in light REE (LREE).
- b) The concentration of REE increases by roughly a factor of 50 from the gabbros to the pillow lavas.

The general trend indicate that these rocks were derived from a uniform LREE depleted asthenosphere.

As our gabbros are from different type we have analyzed some specific samples such as cumulate gabbros, high level gabbros and a dyke of pegmatitic gabbro, the results are given in Table 1. All these data have been normalized with respect to the REE distribution for chondrites given by NAKAMURA (1974); the resulting ratios are presented in Table 2. Looking at those results one notices that with the exception of the La ratio for samples M44 and M69, all our gabbros have REE abundances higher than the chondrites in respect with some of the Troodos gabbros for example (KAY et al., 1976) who have abundances ratios lower than one.

Table 1. REE abundances (ppm) of different kinds of gabbros

Samples nb.	M44	M45	M48	M50	M52	M66	M68	M69	M70B	M81	M85	M103
La	0,2	0,7	3,2	0,5	n.d.	0,7	1,2	0,2	0,7	0,8	2,1	1,0
Ce	n.d.	< 15 *	< 15	< 15	< 15	< 15	< 15	< 15	< 15	< 15	< 15	< 15
Sm	0,5	0,7	1,7	0,7	0,7	0,5	0,9	0,3	0,7	0,9	1,7	0,9
Eu	0,2	0,3	1,5	0,4	0,2	0,5	0,7	0,2	0,5	0,5	1,0	0,6
Dy	1,0	1,5	8,2	1,8	1,5	2,6	3,3	1,0	2,7	1,8	5,5	3,1
Ho	0,5	0,5	0,3	0,6	0,5	0,3	0,6	0,1	0,7	0,8	1,2	1,0
Yb	n.d.	1,2	2,0	3,2	1,1	1,2	1,4	0,4	3,2	3,0	3,3	4,4
Lu	0,3	0,1	n.d.	0,3	0,1	0,2	0,2	0,1	0,3	0,4	0,5	0,6

\* for Ce values below the sensitivity of our method is quoted <15.

Table 2. Normalized REE ratios

	a	M44	M45	M48	M50	M52	M66	M68	M69	M70B	M81	M85	M103
57	La (0.329)	0.606	2.128	9.726	1.520	-	2.128	3.647	0.608	2.128	2.432	6.383	3.039
58	Ce (0.865)	-	-	-	-	-	-	-	-	-	-	-	-
62	Sm (0.203)	2.463	3.448	8.374	3.448	3.448	2.463	4.433	1.478	3.448	4.433	8.374	4.433
63	Eu (0.070)	2.597	3.896	19.480	5.945	2.597	6.493	9.091	2.597	6.493	6.493	12.987	7.792
66	Dy (0.343)	2.915	4.373	23.906	5.248	4.373	7.580	9.621	2.915	7.872	5.248	16.035	9.038
67	Ho (0.070) <sup>b</sup>	7.142	7.149	4.885	8.571	7.142	4.285	8.571	1.428	10.000	11.428	17.142	14.285
70	Yb (0.220)	-	5.454	9.091	14.545	5.000	5.454	6.364	1.818	9.999	13.636	14.999	20.000
71	Lu (0.034)	8.823	2.941	-	8.823	2.941	5.882	5.882	2.941	8.823	11.764	14.706	17.647

a. Chondrite values By Nakamura (1974) in ppm.

b. Value extrapolated from average chondrite plot.

The ratios of four typical gabbros are plotted on a logarithmic scale against REE in increasing atomic number in abscissa (Fig. 1). From this figure it can be seen that the REE patterns are more or less parallel to each other but large differences in concentrations exist. All samples exhibit a depletion in LREE and most of them have a positive Europium anomaly with respect to the neighbouring elements (in our case Sm and Dy). This anomaly is generally related to the cumulate plagioclase.

A crude analysis of the data shows that the high level gabbros sampled in screens in the sheeted complex (e.g. microgabbro M85) have the same REE




Fig. 1 Relative rare earth element patterns of four typical gabbros.

abundances as the dykes from the complex; similar observations have been made by KAY and SENECHAL (1976) in the Troodos, they concluded "... Thus the gabbros in the plutonic complex appear to be derived by crystal sorting from magmas with the same REE pattern as the dykes and lavas".

It is also very interesting to compare the REE behaviour in the cumulate transition zone where one finds interbedding of cumulate gabbro (sample M50) and cumulate peridotite (sample M52, wehrlite). In the latter the REE abundances are lower as expected from ultra-mafic material; La is not detected, Eu exhibits a negative anomaly. The only common point for both samples is the same ratio of Sm.

Another interesting point is that the pegmatitic gabbro sample M81 coming from a dyke in the peridotite has the same pattern with similar abundances as the cumulate gabbro M50.

Finally the olivine gabbro-norite M69 with olivine and two pyroxenes (opx, cpx) has the lowest general trend in agreement with the REE partition rules giving the relative REE content of the main minerals

$$\text{hbl} > \text{cpx} > \text{pl} > \text{opx} > \text{ol}$$

## CONCLUSION

As our REE results on gabbros represent only a step of our REE investigation of the Hatay ophiolite, it is difficult to draw a definite conclusion. Nevertheless one may emphasize that the cumulate and non cumulate gabbros are related to each other by an evolutionary process. It has been shown that the concentrations are varying from 0.6 to  $8.8 \times$  chondrites for a cumulate gabbro to 6.4 to 14.7 for a microgabbro of the top of the plutonic system.

## Acknowledgments

This work was supported by the grant No. 2.322-076 from the Swiss National Science Foundation. Technical assistance was supplied by M.T.A. Ankara for which we are grateful.

## Bibliography

- CANN, J. R., 1970. Rb, Sr, Y, Zr, Nb in some oceanfloor basaltic rocks. *Earth Planet. Sci. Letters* 10, 7-10.
- ÇOĞULU, E., 1973. New data on the petrology of Kızıl Dağ massif (Hatay, Turkey). *Proc. Congress Earth Sci. "50th Anniv. Turk. Rep."*. 409-423.
- ÇOĞULU, E., 1974. Ultrabasic tectonites and layered peridotites of the Hatay Area (Turkey) *Bull. MTA* 83, 139-147.

- ÇOĞULU, E., DELALOYE, M., VUAGNAT, M. and WAGNER, J.-J., 1976. Some geochemical, geochronological and petrophysical data on the ophiolitic massif from the Kızıl Dagh Hatay-Turkey. *C.R. Soc. Phys. Hist. Nat. Genève* 10, 141-150.
- DELALOYE, M., VUAGNAT, M. and WAGNER, J.-J., 1977. K-Ar ages from the Kızıl Dagh ophiolitic complex (Hatay, Turkey) and their interpretation. *Int. Symp. on the Structural history of the Mediterranean Basins. Split (Yugoslavia) oct. 76.* B. Biju-Duval and L. Montadert, Eds. Editions Technip, Paris, 73-78.
- DEWEY, J. F., PITMAN, W. C., RYAN, W. B. F. and BONIN, T., 1973. Plate tectonics and the evolution of the Alpine System. *Geol. Soc. Am. Bull.* 84, 3137-3180.
- DUBERTRET, L., 1953. Géologie des roches vertes du N-W de la Syrie et du Hatay (Turquie). *Notes et Mém. sur le Moyen-Orient*, 6, 227 p.
- GASS, I. G., 1968. Is the Troodos massif of Cyprus a fragment of mesozoic ocean floor? *Nature* 220, 39-42.
- JACKSON, E. D., 1971. The origine of ultramafic rocks by cumulus processes. *Fortsch. Miner. Stuttgart* 48, 128-147.
- KAY, R. W. and SENECHAL, R. G., 1976. The Rare Earth Geochemistry of the Troodos Ophiolite Complex. *J. Geoph. Res.* 81, 964-970.
- MENZIES, M., 1976. Rifting of a Tethyan Continent. Rare Earth evidence of an accreting plate margin. *Earth Plan. Sci. Letters* 28, 427-438.
- MENZIES, M., 1976. Rare earth geochemistry of fused ophiolitic and alpine lherzolites. I: Othris, Lanzo and Trodos. *Geochim. Cosmochim. Acta.* 40, 645-656.
- MOORES, E. M. and JACKSON, E. D., 1974. Ophiolites and the oceanic crust. *Nature* 250, 136-139.
- NAKAMURA, N., 1974. Determination of REE, Ba, Fe, Mg, Na and K in carbonaceous and ordinary chondrites. *Geochim. Cosmochim. Acta* 38, 757-775.
- PARROT, F., 1973. Pétrologie de la coupe du Djebel Moussa, massif basique ultrabasique du Kızıl Dagh (Hatay, Turquie). *Sci. de la Terre*, 18, 143-172.
- PEARCE, J. A., 1975. Basalt geochemistry used to investigate past tectonic environments on Cyprus. *Tectonophysics* 25, 41-67.
- SCHILLING, J.-G., 1975. Rare-earth variations across "normal segment" of the Reykjanes Ridge 60°-53°N, Mid-Atlantic Ridge, 29°S, and East Pacific Rise, 2°-19°S, and evidence on the composition of the underlying low velocity layer. *J. Geophys. Res.* 80, 1459-1473.
- VOLDET, P. and HAERDI, W., 1976. Determination of Eu and Dy in rocks by neutron activation and high-resolution X-ray spectrometry. *Anal. Chim. Acta* 87, 227-231.
- VOLDET, P. and HAERDI, W., 1978. Determination of rare-earth elements in rocks by neutron activation followed by high-resolution X-ray spectrometry or  $\gamma$ -spectrometry. *Anal. Chim. Acta.* 97, 185-189.
- Vuagnat, M. et ÇOĞULU, E., 1967. Quelques réflexions sur le massif basique-ultrabasique du Kızıl Dagh, Hatay, Turquie. *C.R. Soc. Phys. Hist. Nat. Genève* 2/3, 210-216.
- WAGNER, J.-J., DELALOYE, M., HEDLEY, I., VUAGNAT, M. and ÇOĞULU, E., 1976. Magnetic properties of a fragment of Tethyan lithosphere. EGS abstract in *EOS Am. Geoph. Union Trans.*, 57, 655.