Zeitschrift: Bollettino della Società ticinese di scienze naturali

Herausgeber: Società ticinese di scienze naturali

Band: 87 (1999)

Artikel: The use of mitochondrial DNA sequences in insect taxonomy:

examples from the scathophagidae (diptera, calyptratae)

Autor: Bernasconi, M.V. / Valsangiacomo, C, / Piffaretti, J.-C.

DOI: https://doi.org/10.5169/seals-1003285

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

The use of mitochondrial DNA sequences in insect taxonomy: examples from the *Scathophagidae* (Diptera, Calyptratae)

M.V. Bernasconi^{1,2}, C. Valsangiacomo², J.-C. Piffaretti² e P.I. Ward¹

Zoologisches Museum der Universität, Winterthurerstr. 190, CH-8057 Zürich,
Istituto Cantonale Batteriosierologico, Via Ospedale 6, CH-6904 Lugano

Key Words: Scathophagidae, Phylogeny, mitochondrial DNA (mtDNA), Subunit I of the cytochrome oxidase gene (COI)

INTRODUCTION

The use of mitochondrial DNA as molecular markers for a wide range of taxonomic, phylogenetic, population and evolutionary investigations in animals has become well established (AVISE et al. 1987, HILLIS 1987, MORITZ et al. 1987, PACKER 1989, QUICKE 1993, AVISE 1994, SIMON et al. 1994, HILLIS et al. 1996). The positive attributes of mitochondrial DNA can be summarised as follows: (i) Its small size is associated with relative manageability and ease of extraction. (ii) Most cells contain many mitochondria, so that a significant quantity of mtDNA is available even from small tissue samples. (iii) A general conservation of gene order and composition across metazoa. (iv) A range of mutational rates in different regions of the molecule. (v) Because of its maternal inheritance, differences that occur in mtDNA are due entirely to mutation and are not the result of independent assortment or recombination (Quicke 1993, Moritz et al. 1987, Wolstenholme 1992, AVISE et al. 1987).

The mtDNA in higher animals is a circular, double-stranded molecule. It comprises approximately 16'000 base-pairs encoding two rRNAs, 22 tRNAs and usually 13 other genes, mostly coding for proteins involved in the electron transport system located on the inner mitochondrial membrane. Animal mitochondrial genes lack introns and intergenic sequences are generally small or lacking. Also absent are repetitive DNA and pseudogenes. Further, the genetic code employed in mitochondria is slightly different from that of nuclear genes. (Clary & Wolstenholme 1985, Avise et al. 1987, Moritz et al. 1987, Wolstenholme 1992, Crozier & Crozier 1993).

Although mtDNA-sequence data have proven valuable in phylogenetic analysis, gene choice is of crucial importance (SIMON et al. 1994, LUNT et al. 1996). Among the coding genes in the mitochondria genome, the subunit I of the cytochrome oxidase (COI) gene possesses features making it particularly suitable for evolutionary studies (LUNT et al. 1996). First, it has been well-studied at the biochemical level and its size and structure appears to be conserved across all animals investigated. Then, because the COI gene is one of the largest protein-coding genes in

the animal mitochondrial genome, it is possible to amplify and sequence a large portion of DNA within the same functional complex. Furthermore, highly conserved and variable regions are closely associated.

Using examples taken from the Scathophagidae (Diptera), we illustrate the usefulness of mtDNA sequences in elucidating taxonomic and phylogenetic questions. Scathophagids flies, with more than 250 species, are mainly confined to the Holarctic region. Only about 5 species occur south of the equator, in South America and Africa, mostly at high altitudes, whereas in the Oriental region one species occurs in the Malay Peninsula (VOCKEROTH 1956, VOCKEROTH 1989). Individuals of most species feed on vertebrate dung or decomposing carcasses, performing the ecologically-important function of resource recycling. One species, Scathophaga stercoraria, has been used extensively to investigate questions in animal ecology and evolution (SIGURIONSDOTTIR & PARKER 1980, Amano 1983, Parker & Simmons 1991, Simmons & WARD 1991, WARD & SIMMONS 1991, SIMMONS & Parker 1992, Parker et al. 1993, Ward 1993, Ward & HAUSCHTECK-JUNGEN 1993; MÜHLHÄUSER et al. 1996, SIMMONS et al. 1996, BLANCKENHORN 1997, OTRONEN et al. 1997, Blanckenhorn 1998, Ward 1998). However, the taxonomy and phylogeny of this family still remain unclear. The use of morphological characters, including the structure of male genitalia which seems to be very variable for some species are usually enough for the correct identification of genera and species but are still insufficient to infer precise phylogeny. The integration of the new molecular data with previous morphological studies should provide therefore a powerful tool for the inference of correct phylogeny of this fly family.

MATERIALS AND METHODS

After DNA extraction and standard PCRs, a portion of 810 bp of the terminal region of the COI gene was sequenced in 40 Scathophagid species covering a wide geographic area, as well as a diverse spectrum of ecological habitats. Moreover, three other flies species of the *Mus*-

coidea superfamily, Musca domestica (family Muscidae), Lasiomma seminitidum (Anthomyiidae) and Fannia armata (Fanniidae) were also sequenced. Outgroup comparisons were made with published COI gene sequence of Drosophila yakuba (EMBL database access X03240), and translations to amino acid sequences used the Drosophila code (DE BRUIJN 1983). Sequence data were analysed by pairwise sequence alignment and by multi alignment with the Lasergene program Megalign (DNAstar 1994) and phylogenetic analysis were performed using MEGA (Molecular Evolutionary Genetics Analysis 1.01, KUMAR et al. 1993). Several Neighbour Joining trees were generated using different methods for distance estimation. Some Maximum Parsimony trees obtained with heuristic search were also examined.

RESULTS AND DISCUSSION

The topologies of the several phylogenetic trees generated are overall very similar (Fig. 1). In general, molecular data are in agreement with previous morphological information. But some peculiarities reveal discrepancies between the two approaches. Within the *Scathophagidae* two subfamilies are currently recognised: the *Delinae* (represented here by the species *Delina nigrita* and *Phrosia albilabris*) and the *Scathophaginae* (GORODKOV 1986, VOCKEROTH 1989). However, various authors have proposed a different number of subfamilies or phylogenetic groups (ranging from 2 to 5; SACK 1976). The molecular data do not confirm the separation in 2 subfamily-groups currently recognised and the relationships among the groups proposed in the past




Fig. 1 - Rooted Neighbour Joining tree for 40 Scathophagids species and 3 other Muscoidea species (*Musca domestica* belongs to the *Muscidae*, *Lasiomma seminitidum* to the *Anthomyüdae*, and *Fannia armata* to the *Fanniidae*) based on the Tamura-Nei model and gamma corrected distances (with default gamma parameter a=0.5). *Drosophila yakuba* was chosen as outgroup. Bootstrap values for 500 replicates are shown.

are not always confirmed. Only the Delina genus appears sometimes to be isolated relative to the other genera. Nevertheless, this information is not supported by all the trees generated and the bootstrap values are usually low. Concerning the genus *Phrosia*, its membership of the *Delinae* is controversial even from the morphological point of view. As a matter of fact some authors consider this genera as a Scathophaginae and cannot understand why Phrosia should be placed in the Delinae subfamily (F. Püchel, personal communication 1997). In particular, adult individuals of Phrosia albilabris are morphologically very similar to the species belonging to the genus Cordilura (R. Vockeroth, personal communication 1998). Molecular data not only support the idea that *Phrosia* should be considered as a true Scathophaginae but also place this species unequivocally within the Cordilura genus.

Pogonota, Lasioscelus and Oheniella were once treated at the genus rank and considered as belonging to the same phylogenetic group (HACKMAN 1956). Modern taxonomists have now placed Lasioscelus as a subgenera of Pogonota (GORODKOV 1986) The molecular data confirm the relationship among the 3 groups relative to the Scathophagidae. However, it seems not correct to consider Lasioscelus as a subgenera of Pogonota, since no tree supports this interpretation. Even the generally high bootstrap values indicate a more strict relationship between Pogonota and Oheniella than between Pogonota and Lasioscelus.

Scathophaga taeniopa and S. suilla, are morphologically very similar species. In particular, a detailed study based on traditional morphological characters and comparing also the genitalia of the two species concluded that S. taeniopa and S. suilla are identical (SIFNER 1975, 1995). The major problem related to these two species is the risk of error by morphological determination, perhaps because of the lack of clearly distinctive chaetotaxy which could be applied to all the individuals. In this context the molecular data are very useful, allowing a sure characterisation of the specimens. Even if the case is not to be considered closed, the molecular data unequivocally suggest that S. taeniopa and S. suilla are different and must therefore be considered as distinct taxa.

S. analis, even if present in classical works on Scathophagidae, is considered as a doubtful species in the Catalogue of Palaearctic Diptera (GORODKOV 1986). Molecular data seems to indicate that it is probably a synonym of S. inquinata. In the determination key, the discriminating point is represented by the anterior cross-vein which should be dark-shadowed in S. analis but not in S. inquinata (SACK 1976). It is possible that the cross-vein character used to separate the two fly types is not valid, even if analysing the specimens from a morphological point of view, some other showy differences appear, like for example the colour of the shoulders. Perhaps under the name S. analis and S. inquinata entomologists have separated individuals belonging to the same species. The only way to clarify the question is to get sequence information generated from specimens which have been previously identified without any doubt by morphological characters.

In conclusion, as shown by these examples, the inte-

gration of the molecular data derived from mtDNA sequences, together with previous morphological studies, provides a keystone for the inference of correct phylogeny and taxonomy of an insect family.

Acknowledgements

We would like to thank all the people who helped us in very different ways. In particular we thank F. Püchel (Bielefeld), F. Sifner (Praha), R. Vockeroth (Ottawa), J. Pawlowski (Geneva), P. Flook (Basel), B. Merz (Zurich), M. Nelson (Edinburgh), L. Greve Jensen (Bergen), V. Michelsen (Copenhagen), R. Meier (Copenhagen), M. v. Tschirnhaus (Bielefeld), Knut Rognes (Stavanger), Walter Hackman (Helsinki), R. Cuny (Alberta), R. Schmidt (Zurich), G. Bächli (Zurich) and M. Otronen (Turku).

REFERENCES

- AMANO K., 1983. Studies on the intraspecific competition in dung breeding flies. 1. Effects on larval density in yellow dung fly, *Scathophaga stercoraria* (Diptera: Scathophagidae). Jap J Sanit Zool 34:165-175.
- AVISE J.C., 1994. Molecular Markers, Natural History and Evolution. Chapman & Hall New York-London, xiv, 511.
- Avise J.C., Arnold J., Ball R.M., Bermingham E., Lamb T., Neigel J.E., Reeb C.A., Saunders N.C., 1987. Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. Ann Rev Ecol Syst 18: 489-522.
- BLANCKENHORN W.U., 1997. Effects of temperature on growth, development and diapause in the yellow dung fly-against all rules? Oecologia 111: 318-324.
- BLANCKENHORN W.U., 1998. Adaptive phenotypic plasticity in grouth, development, and body size in the yellow dung fly. Evolution 52: 1394.
- CLARY D.O. & WOLSTENHOLME D.R., 1985. The mitochondrial DNA molecule of *Drosophila yakuba*: Nucleotide sequence, gene organisation, and genetic code. J Mol Evol 22: 252-271.
- CROZIER R.H. & CROZIER Y.C., 1993. The mitochondrial genome of the honeybee *Apis mellifera*: complete sequence and genome organisation. Genetics 133: 97-117
- DE BRUIJN M.H., 1983. *Drosophila melanogaster* mitochondrial DNA, a novel organization and genetic code. Nature 304 (5923): 234-241.
- GORODKOV K.B., 1986. Catalogue of palaearctic Diptera, Scathophagidae-Hypodermatidae, 11: 11-41.
- HACKMAN W., 1956. The Scatophagidae (Dipt.) of eastern Fennoscandia. Helsingsforsiae, 1-67.
- HARRISON R.G., 1989. Animal Mitochondrial DNA as a genetic marker in population and evolutionary biology. Trend Ecol Evol 4: 6-11.
- HILLIS D.M.. 1987. Molecular versus morphological approaches to systematics. Ann Rev Ecol Syst 118: 23-42.
- HILLIS D.M., MORITZ C., MABLE B.K., 1996. Molecular systematics, Second edition, Sinauer Associates, Inc. Publishers, Sunderland Massachusetts USA. xvi, 655.
- KUMAR S., TAMURA K., NEI M., 1993. Molecular evolutionary genetics analysis, version 1.01. The Pennsylvania State University, University Park, PA 16802.

- LUNT D.H., ZHANG D.-X., SZYMURA J.M., HEWITT G.M., 1996. The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies. Insect Mol Biol 5(3): 153-165.
- MORITZ C., DOWLING T.E., BROWN W.M., 1987. Evolution of animal mitochondrial DNA: relevance for population biology and systematics. Ann Rev Ecol Syst 18: 269-292.
- MÜHLHÄUSER C, BLANCKENHORN W.U., WARD P.I., 1996. The genetic component of copula duration in the yellow dung fly. Anim Behav 51: 1401-1407.
- Otronen M., Reguera P., Ward P.I., 1997. Sperm storage in the yellow dung fly *Scathophaga stercoraria*: Identifying the sperm of competing males in separtate female spermathecae. Ethology 103, 844-854.
- PACKER C., 1989. Molecular Ecology, Trend Ecol Evol 4: 156-157. PARKER G.A. & SIMMONS L.W., 1991. A model of constant random sperm displacement during mating: evidence from *Scatophaga*. Proc R Soc Lond B 246: 107-115.
- Parker G.A., Simmons L.W., Ward P.I., 1993. Optimal copula duration in dungflies: effects of frequency dependence and female mating status. Behav Ecol Sociobiol 32:157-166.
- QUICKE D.L.J., 1993. Principles and techniques of contemporary taxonomy. Blackie Academic & Professional. xii, 311.
- SACK P., 1976. 62a Cordyluridae. In Lindner, E. (ed.) Die Fliegen der Palaearktischen Region, E. Schweizerbart'sche Verlagsbuchhandlung Stuttgart, 7: 1-103.
- SIFNER F., 1975. Scatophagidae (Diptera) de Mongolie. Annales Historico-Naturales Musei Nationalis Hungarici 67: 219-227.
- SIFNER F., 1995. Additions to the Catalogue of flies of the Palaearctic region (Diptera: Scatophagidae). Eur J Entomol 92: 513-515
- SIGURJONSDOTTIR H. & PARKER G.A., 1980. Dung fly struggles: evidence for assessment strategy. Behav Ecol Sociobiol 8: 219-230.

- SIMMONS L.W. & PARKER G.A., 1992. Individual variation in sperm competition success of yellow dung fly *Scathophaga ster-coraria* (L.). Evolution 46: 366-375.
- SIMMONS L.W. & WARD, P.I., 1991. The heritability of sexually dimorphic traits in the yellow dung fly *Scathophaga stercoraria* (L.). J Evol Biol 4: 593-601.
- SIMMONS L.W., STOCKLEY P., JACKSON P., PARKER G.A., 1996. Sperm competition or sperm selection: no evidence for female influence over paternity in yellow dung fly *Scathophaga sterco-raria*. Behav Ecol Sociobiol 38: 199-206.
- SIMON C., FRATI F., BECKENBACH A., CRESPI B., LIU H., FLOOK, P., 1994. Evolution, Weighting, and Phylogenetic Utility of Mitochondrial Gene Sequences and a Compilation of Conserved Polymerase Chain Reaction Primers. Ann Entomol Soc Am 87(6): 651-701.
- VOCKEROTH J.R., 1989. Scathophagidae. In McAlpine (ed) Manual of nearctic Diptera 2: 1085-1097.
- VOCKEROTH J.R., 1956. Distribution patterns of the Scatomyzinae (Diptera, Muscidae). Proceedings tenth international congress of entomology 1: 619-626.
- WARD P.I., 1993. Females influence sperm storage and use in the yellow dung fly Scathophaga stercoraria (L.). Behav Ecol Sociobiol 32: 313-319.
- WARD P.I., 1998. A possible explanation for cryptic female choice in the yellow dung fly, Scathophaga stercoraria (L.). Ethology 104: 97-110.
- WARD P.I. & HAUSCHTECK-JUNGEN E., 1993. Variation in sperm length in the yellow dung fly *Scathophaga stercoraria* (L.). J Insect Physiol 39 (7): 545-547.
- WARD P.I. & SIMMONS L.W. 1991. Copula duration and testes size in the yellow dung fly, *Scathophaga stercoraria* (L.): the effects of diet, body size, and mating history. Behav Ecol Sociobiol, 29: 77-85
- WOLSTENHOLME D.R., 1992. Animal mitochondrial DNA: structure and evolution. Int. Rev. Cytol. 141: 173-216.