

Zeitschrift: Jahrbuch der Schweizerischen Naturforschenden Gesellschaft.
Wissenschaftlicher und administrativer Teil = Annuaire de la Société
Helvétique des Sciences Naturelles. Partie scientifique et administrative

Herausgeber: Schweizerische Naturforschende Gesellschaft

Band: 162 (1982)

Artikel: Developmental regulation of tumor autonomy in plants

Autor: Meins, Frederick

DOI: <https://doi.org/10.5169/seals-90899>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Developmental regulation of tumor autonomy in plants

Frederick Meins, Jr.

Summary

Neoplastic diseases of plants have several proximal causes; for example, genetic transformation by plasmids of bacterial origin; transformation by double stranded RNA viruses; and, the combination of foreign genomes in interspecific plant hybrids. Regardless of proximal cause, autonomous growth in these diseases appears to result from the inappropriate production of growth factors by the tumor cells. There is evidence that genetic changes are neither necessary nor sufficient for expression of the tumor state in plants: autonomously growing cells can arise as a consequence of epigenetic changes. Tumor cells that contain foreign genes necessary for autonomous growth can differentiate, participate in organogenesis, and exhibit normal growth regulation. This suggests that even when transformation results from a genetic lesion, expression of the tumor state is ultimately regulated by epigenetic mechanisms of the type operating in normal development.

Introduction

As a result of neoplastic transformation, cells acquire the capacity for autonomous growth, which is then inherited by individual tumor cells. Plant neoplastic diseases provide experimental systems well suited for studying the transformation process since the physiological basis for autonomous growth is known and it is possible to establish the developmental and genetic potentialities of the tumor cell by regeneration of complete plants.

In this article I will briefly review the biology of plant tumors using crown gall disease as an example and then provide evidence to support the hypothesis that tumor transformation in plants is a form of inappropriate

cell differentiation which is precisely regulated by the same types of mechanisms that operate in normal development.

Tumor inception in crown-gall

Crown-gall tumors result when cells conditioned by wounding from a wide variety of dicotyledonous plant species and certain gymnosperms are exposed to a tumor inducing principle (TIP) elaborated by virulent strains of *Agrobacterium tumefaciens*. The tumors which form at the site of inoculation are true tumors since the cells continue to express their neoplastic character in culture and when grafted onto the host plant in the absence of the inciting bacterium. The type of tumor obtained depends on the plant host, the site of inoculation, and the strain of bacterium used (Braun 1978). For example, *Kalanchoë* or tobacco plants inoculated with the B6 strain of bacterium form large, unorganized tumors that produce the opine octopine, whereas plants inoculated with the T37 strain form complex, highly organized tumors, known as teratomas, that produce another opine, nopaline. Both the pattern of development and type of opine made persist in cloned cell lines serially transferred in culture, indicating that these traits are inherited at the cellular level.

There are several lines of evidence indicating that TIP is a DNA sequence derived from a large, tumor-inducing plasmid (Ti) present in virulent strains of the crown-gall bacterium. First, during tumor inception a portion of this plasmid, T-DNA, is transferred to the host cell (Chilton *et al.* 1977) where it is integrated into one or more sites in the nuclear DNA (Yadav *et al.* 1980). Second, the type of opine produced by the tumor cell is specified by T-DNA (Bomhoff *et al.* 1976), which, in the case of octopine plasmids, for

example, codes for the key enzyme in the synthesis of octopine (Schröder *et al.* 1981; Murai and Kemp 1982). Third, mutations at specific sites in the T-DNA affect the growth and development of tumors elicited with the mutant strains of plasmid (Hooykaas *et al.* 1982; Garfinkel *et al.* 1981). Finally, reversion of tumor cells is accompanied by the loss of all, or a major portion, of the T-DNA (Yang *et al.* 1980; Yang and Simpson 1981). Thus it appears that the maintenance of the tumor state in crown-gall depends on the presence of Ti-plasmid-DNA sequences that are transferred from the bacterium to the plant cell during tumor inception.

Growth-promoting substances and tumor autonomy

Comparison of the nutritional requirements of normal and tumor cells in culture provides strong evidence that the autonomous growth of plant tumors involves an alteration in the capacity of the cell to produce growth factors. Several growth factors are required for proliferation of higher plant cells in culture. The most important ones are the auxins, which promote DNA synthesis, mitosis and cell enlargement, and cell-division factors, such as the cytokinins, which promote cell division. For example, pith cells of tobacco exhibit an absolute requirement for exogenous auxin and cytokinin for continuous growth on an otherwise complete medium (Jablonski and Skoog 1954). As a result of transformation, these cells acquire the capacity to grow without the added factors and produce auxin and cytokinin in quantities sufficient to support the growth of normal cells (Braun 1958). Similar changes, although less well-documented, are also found in tumors arising in tumor-prone genetic hybrids (Bayer 1982) and in wound tumor disease, which is caused by a double-stranded RNA virus (Black 1982). These findings lead to the important conclusion that, regardless of the proximal cause for transformation, autonomous growth results from changes in the production of substances necessary for the proliferation of normal cells (Braun 1958; Meins 1974).

The details of how production of growth factors is regulated in tumor cells and the relationship between growth and production

of these factors is not known. The observation that crown-gall tumors on plants mimic the application of auxin has led to the hypothesis that tumor autonomy results from the overproduction of auxin (Link and Eggers 1941). This seems unlikely. Tumor tissues *in planta* and in culture contain auxin at concentrations in the range found in rapidly growing normal tissues of the same plant species (Weiler and Spanier 1981; Pengelly and Meins 1981). There is growing evidence that different genes in the T-DNA affect auxin and cytokinin production by the tumor cell (Garfinkel *et al.* 1981; Hooykaas *et al.* 1982; Morris *et al.* 1982). It is not known, however, whether these genes code for enzymes needed for the synthesis of these factors or act indirectly by altering the expression of host genes for these enzymes.

Developmental regulation of growth autonomy

Insight into the relative contribution of foreign and host genes in the regulation of growth autonomy is provided by the observation that heritable changes in auxin and cytokinin requirement similar to those encountered in transformation also occur in normal development. For example, leaf cells of tobacco require exogenous cytokinin for rapid growth in culture, whereas stem-cortex cells do not (Meins and Lutz 1979). The fact that both states are inherited by individual cells implies that heritable changes in cytokinin requirement occurred sometime in the development of the two tissue types. Direct evidence that this can result in autonomous growth comes from studies of habituation (Meins 1982a).

Tissues from a variety of plant species in culture sometimes lose their requirement for exogenous auxin, or for both auxin and cytokinin. This process, known as habituation, is progressive and generates autonomously proliferating cells, which, in some cases, form transplantable tumors when grafted onto the host plant. Therefore, it appears that habituation is a form of spontaneous neoplastic transformation that occurs in culture.

In spite of their extreme phenotypic stability when serially propagated in culture, cytokinin-habituated cells can be induced to lose




Fig. 1. Plant regeneration experiment showing that the cytokinin-habituated phenotype (C^+) reverts to the non-habituated phenotype (C^-) when cloned tobacco cells of pith origin are induced to form complete plants. From Meins 1974.

their autonomous character. Cloned lines of habituated tobacco cells, which are of pith origin, regularly form normal-looking, fertile tobacco plants when placed on a differentiation-inducing medium. Pith tissues from the regenerated plants exhibit the normal, non-habituated phenotype in culture (fig. 1). Results of this type obtained with many cloned lines and detailed measurements of the rates at which habituation is induced and reversed in culture indicate that habituation involves heritable, potentially

reversible, epigenetic changes rather than classical mutations (Meins 1982a). Conversion of cells to the cytokinin-autotrophic state is a key event in both tumor transformation and habituation, which has an epigenetic basis. Therefore, neither somatic mutation nor the introduction of foreign genes need be invoked to account for the stability of the tumor state.

Functions essential for autonomous growth can also be switched on and off in tumor cells containing foreign genes (Braun 1959;

Braun and Wood 1976; Turgeon *et al.* 1976). When cloned, crown-gall teratoma tissues of tobacco are grafted onto the cut stem of tobacco plants from which the auxillary buds have been removed, they give rise to shoots which gradually become more normal in appearance. Some of these shoots eventually flower and set fertile seed. These seeds develop into normal tobacco plants, which have completely lost their tumorous character and, as expected, no longer contain readily detectable T-DNA sequences (Yang *et al.* 1980). Reversion with the loss of Ti-plasmid genes also occurs spontaneously in cultured teratoma tissues, but at a low incidence (Yang and Simpson 1981; Meins 1982a).

Of particular interest, however, are those cases in which tumor cells mimic the normal state but still retain T-DNA sequences. Leaves near the top of teratoma-derived shoots are perfectly normal in gross morphology, physiology and histology. Nevertheless these leaves still produce nopaline (Wood *et al.* 1978), contain T-DNA (Yang *et al.* 1980), and, when placed in culture on a basal medium that does not support the growth of normal leaf tissue, give rise to typical teratoma tissues. It may be argued that the teratoma-derived shoots consist predominantly of revertant cells and that teratomatous tissues arise from a few tumor cells that still retain T-DNA. There is strong evidence against this. Protoplasts from highly differentiated mesophyll cells isolated from leaves on teratoma-derived shoots regularly form teratomatous clones when cultured (Binns *et al.* 1981). Similar results are obtained with single, highly specialized, epidermal hair cells (Huff and Turgeon 1981). Thus, teratoma-derived shoots consist of tumor cells that differentiate, participate in organogenesis, and exhibit normal growth regulation while retaining their tumorous character in a covert form.

Suppression of the tumorous state also occurs in plants regenerated from tumors of genetic hybrids. Carlson *et al.* (1972) have fused protoplasts from two species of *Nicotiana* that give tumor-prone hybrid plants. The dikaryotic clones which result, like crown-gall tissues, do not require auxin and cytokinin for growth in culture. Plants regenerated from these clones are tumor prone;

when wounded, typical teratomas arise at the wound site. This shows that cells genetically predisposed to grow autonomously do not express their tumor character in the intact plant, even though the cells retain this potential. Finally, there is a mutation of tobacco in which leaf cells exhibit the cytokinin-habituated phenotype in culture (Meins, 1982c). Plants regenerated from these cells develop normally and have normal appearing leaves, but the leaf tissue still expresses its autonomous character when placed in culture. Therefore, autonomous growth whether resulting from the incorporation of plasmid genes, mutation, or the hybridization of dissimilar genomes is phenotypically suppressed when plants or shoots are regenerated from the altered cells.

Conclusion

There is good evidence that the autonomous growth of plant tumor cells involves the inappropriate production of growth substances such as auxin and cytokinin. What is not clear, at present, is the molecular basis for these changes. It is still not known whether the primary lesion in different tumor diseases is the same or not, whether growth-factor requirement is regulated at the level of growth-factor production or growth-factor degradation, and how foreign genes regulate these processes.

Genetic events such as the integration of foreign genes, interspecific hybridization and somatic mutation appear to be neither necessary nor sufficient for the expression of growth autonomy. Autonomous tumor cells can arise by epigenetic changes that do not seem to involve permanent genetic changes. Studies of teratoma derived shoots and genetic hybrid tumors show that expression of the tumor state can be blocked even when transformation results from a genetic lesion. The picture that emerges is that transformation is a form of abnormal differentiation subject to regulation by the same types of epigenetic mechanisms involved in normal development.

Acknowledgements

I wish to thank J.-F. Conscience, P. King, and C. Moroni for their useful comments

and criticism. Work cited from the author's laboratory was supported in part by grant No. CA20053 from the U.S. Public Health Service, National Cancer Institute.

Updating note

Since this article was written in 1982, direct evidence has been obtained that genes in the T-DNA code for enzymes important in auxin and cytokinin production (Barry *et al.*, 1984; Inze *et al.*, 1983; Schroeder *et al.*, 1984). Comparison of the cytokinin and auxin contents of habituated and crown gall transformed cells has provided further support for the hypothesis that plant cells contain genes with functions similar to the oncogenic genes in the T-DNA (Pengelly and Meins, 1983; Hansen *et al.*, in press). An excellent review of the problem of tumor autonomy in plants has recently appeared (Binns, 1983).

References

- Barry, G.F., Rogers, S.G., Fraley, R.T., Brand, L. 1984. Identification of a cloned cytokinin biosynthetic gene. *Proc. Natl. Acad. Sci. USA.*, 81, 4776-4780.
- Bayer, M. 1982: Genetic tumors: Physiological aspects of tumor formation in interspecies hybrids. In: Kahl, G., Schell, J.S.: *Molecular biology of plant tumors*. Academic, New York and London.
- Binns, A., Wood, H.N., Braun, A.C. 1981: Suppression of the tumorous state in Crown gall teratomas of tobacco. A clonal analysis. *Differentiation* 19, 97-102.
- Binns, A.N. 1983. The biology and molecular biology of plant cells infected by *Agrobacterium tumefaciens*. In: Mifflin, B.J.: *Oxford Surveys of Plant Molecular and Cell Biology*, pp. 133-160, Clarendon, Oxford.
- Black, L.M. 1982: Wound tumor disease. In: Kahl, G., Schell, J.S.: *Molecular biology of plant tumors*. Academic, New York and London.
- Bomhoff, G., Klapwijk, P.M., Kester, H.C.M., Schilperoort, R.A., Hernalsteens, J.P., Schell, J. 1976: Octopine and nopaline synthesis genetically controlled by a plasmid of *Agrobacterium tumefaciens*. *Molec. and Gen. Genetics* 145, 177-181.
- Braun, A.C. 1959: A demonstration of the recovery of the crown-gall tumor cell with the use of complex tumors of single-cell origin. *Proc. Natl. Acad. Sci. U.S.A.* 45, 932-938.
- Braun, A.C. 1978: Plant tumors. *Biochim. Biophys. acta* 516, 167-191.
- Braun, A.C., Wood, H.N. 1976: Suppression of the neoplastic state with the acquisition of specialized functions in cells, tissues, and organs of crown-gall teratomas of tobacco. *Proc. Natl. Acad. Sci. U.S.A.* 73, 469-500.
- Carlson, P.S., Smith, H.H., Dearing, R.D. 1972: Paraxial interspecific plant hybridization. *Proc. Natl. Acad. Sci. U.S.A.* 69, 2292-2294.
- Chilton, M.-D., Drummond, M.H., Merlo, D.J., Sciaky, D., Montoya, A.L., Gordon, M.P., Nestor, E.W. 1977: Stable incorporation of plasmid DNA into higher plant cells: The molecular basis of Crown gall tumorigenesis. *Cell* 11, 263-271.
- Garfinkel, D.J., Simpson, R.B., Ream, L.W., White, F.F., Gordon, M.P., Nester, E.W. 1981: Genetic analysis of crown gall: Fine structure map of the T-DNA by site-directed mutagenesis. *Cell* 27, 143-153.
- Hansen, C.E., Meins, F., Jr., Milani, A. Clonal and physiological variation in the cytokinin content of tobacco-cell lines differing in cytokinin requirement and capacity for neoplastic growth. *Differentiation*, in press.
- Hooykaas, P.J.J., Ooms, G., Schilperoort, R.A. 1982: Tumors induced by different strains of *Agrobacterium tumefaciens*. In: Kahl, G., Schell, J.S.: *Molecular biology of plant tumors*. Academic, New York and London.
- Huff, M., Turgeon, R. 1981: Neoplastic potential of trichomes isolated from tobacco crown gall teratomas. *Differentiation* 19, 93.
- Inze, D., Follin, A., Van Lijsebettens, M., Simeons, C., Genetello, C., Van Montagu, M., Schell, J. 1984. Genetic analysis of the individual T-DNA genes of *Agrobacterium tumefaciens*; further evidence that two genes are involved in indole-3-acetic acid synthesis. *Molec. Gen. Genet.*, 194, 265-274.
- Jablonski, J.R., Skoog, F. 1954: Cell enlargement and cell division in excised tobacco pith tissue. *Physiol. Plant* 7, 16-24.
- Link, G.K.K., Eggers, V. 1941: Hyperauxiny in crown gall of tomato. *Bot. Gaz.* 103, 87-106.
- Meins, F., Jr. 1974: Mechanisms underlying tumor transformation and tumor reversal in crown-gall, a neoplastic disease of higher plants. *Symp. Soc. Devel. Biol.* 32, 23-39.
- Meins, F., Jr. 1982a: Habituation of cultured plant cells. In: Kahl, G., Schell, J.S.: *Molecular biology of plant tumors*. Academic, New York and London.
- Meins, F., Jr. 1982b: Tumor reversal in plants. *Prog. Clinical Biol. Rsch.* 85B, 567-576.
- Meins, F., Jr. 1982c: The nature of the cellular heritable change in cytokinin habituation. In: Earle, E., Demarly, Y.: *Variability in plants regenerated from tissue culture*. Praeger, New York.
- Meins, F., Jr., Lutz, J. 1979: Tissue-specific variation in the cytokinin habituation of cultured tobacco cells. *Differentiation* 15, 1-6.
- Morris, R.O., Akiyoshi, D.E., MacDonald, E.M.S., Morris, J.W., Regier, D.A., Zaerr, J.B. 1982: Cytokinin metabolism in relation to tumor induction by *Agrobacterium tumefaciens*. In: Wareing, P.F.: *Plant growth substances 1982*. Academic Press, London and New York.

- Murai, N., Kemp, J. 1982: Octopine synthase mRNA isolated from sunflower crown gall callus is homologous to the Ti plasmid of *Agrobacterium tumefaciens*. Proc. Natl. Acad. Sci. U.S.A. 79, 86-90.
- Pengelly, W.L., Meins, F., Jr. 1981: The relationship of indole-3-acetic acid content and growth of crown-gall tumor tissues of tobacco in culture. Differentiation. 21, 27-31.
- Pengelly, W.L., Meins, F., Jr. 1983. Growth, auxin requirement, and indole-3-acetic acid content of cultured crown-gall and habituated tissues of tobacco. Differentiation, 25, 101-105.
- Schröder, J., Schröder, G., Huisman, H., Schilperoort, R.A., Schell, J. 1981: The mRNA for lysopine dehydrogenase in plant tumor cells is complementary to a Ti-plasmid fragment. FEBS Letters 129, 166-168.
- Schroeder, G., Waffenschmidt, S., Weiler, E.W., Schroeder, J. 1984. The T-region of Ti plasmid codes for an enzyme synthesizing indole-3-acetic acid. Eur. J. Biochem., 138, 387-391.
- Turgeon, R., Wood, H.N., Braun, A.C. 1976: Studies on the recovery of Crown gall tumor cells. Proc. Natl. Acad. Sci. U.S.A. 73, 3562-3564.
- Weiler, E.W., Spanier, K. 1981: Phytohormones in the formation of crown gall tumors. Planta 153, 326-337.
- Wood, H.N., Binns, A.N., Braun, A.C. 1978: Differential expression of oncogenicity and nopoline synthesis in intact leaves derived from crown gall teratomas of tobacco. Differentiation. 11, 175-180.
- Yadav, N.S., Postle, K., Saiki, R.K., Thomashaw, M.F., Chilton, M.-D. 1980: T-DNA of a crown gall teratoma is covalently joined to host plant DNA. Nature. 287, 458-461.
- Yang, F.-M., Montoya, A., Merlo, D.J., Drummond, M.H., Chilton, M.-D., Nester, E.W., Gordon, M.P. 1980: Foreign DNA sequences in crown gall teratomas and their fate during the loss of the tumorous trait. Molec. Gen. Genetics. 177, 707-714.
- Yang, F.-M., Simpson, R. 1981: Revertant seedlings from crown gall tumors retain a portion of the bacterial T-DNA sequences. Proc. Natl. Acad. Sci. U.S.A. 78, 4151-4155.

Address of Author

Dr. Frederick Meins, Jr.
Friedrich Miescher-Institut
P.O. Box 2543
CH-4002 Basel (Switzerland)