Zeitschrift: The Swiss observer: the journal of the Federation of Swiss Societies in

the UK

Herausgeber: Federation of Swiss Societies in the United Kingdom

Band: - (1979)

Heft: 1758

Rubrik: For the gardener

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 27.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

widening of numerous footpaths and the provision of new ones, it is now possible to undertake even big walking tours in the Aletsch region dryshod nearly all the year round. A partially new footpath runs at a slight elevation across the left bank of the Rhône at a height of between 2,500 and 3,300 ft from Brig via Mörel and Bister to Grengiols and from there into the Binn Valley. On the other side of the Rotten, walking is possible on snow-free routes at a halfway height of between 3,300 ft and 4,500 ft from Mörel via Breiten to Ried-Greich-Goppisberg-Betten into the Fiesch Valley. In midsummer and autumn the attractions are the walking routes at the height of about 6,600 ft, such as the new path from Belalp across the Aletsch Glacier to Riederalp and Bettmeralp. Walking and natural science weeks are, incidentally, organised on the Riederalp and Bettmeralp.

BIRTHDAYS & ANNIVERSARIES

The following readers will be celebrating birthdays and anniversaries during the next few weeks:—

Mr. H. Humbel will be 74 on 5th October and Madame de Fischer, Berne, will celebrate her birthday on 10th. Mrs. Keufeler on the 18th and Mrs. H. Shepherd will be 77 on the 21st. Dr. E. M. Bircher (Bissone TI) will be 84 on the 22nd and Mrs. Theresa Furrer will be 89 on the 24th. Mr. H.J. Thomson will be 65 on the 25th October and Mr. A. Kunzler will be 83 on the 15th November.

Mr. and Mrs. Reutlinger (Basle) will have their 34th wedding anniversary on 9th October on the same date Mr. and Mrs. E. Bader will have been married for 64 years.

Congratulations and many happy returns to all of them and any other subscribers whose birthdays or anniversaries we are unaware.

FOR THE GARDENER

BLACK CURRANT

Young plants should be planted between October and at latest March, November is the ideal month. Plant around 4 ft to 6 ft apart and 1½ in to 2 in deeper than the soil mark as this encourages bottom growth. Immediately after planting cut down ALL shoots to second bud up from soil level and give good manure or compost mulch.

Ideal soil is well drained but moisture retaining into which manure or compost is worked before planting.

Every year in March give good mulch of manure or compost at the rate of two buckets per square yard. In late January feed with 1 oz (one tablespoon) sulphate of potash and every third year add 2 ozs (two tablespoons) of superphosphate. Water well in dry spells.


As plants are shallow rooting do not hoe or fork weeds but stifle with a good mulch, grass cuttings can be used for this.

Buds and fruit should be protected from birds, fruit should not be picked until fully ripe.

Fruit is produced in the main on previous year's growth so the aim of pruning is to remove some of the older darker wood to encourage new growth. Keep balance between extension growths and new shoots from the base.

Propagation can be by taking cuttings of half ripe wood about 9 in long. Cut just above a bud and trim unripe wood from tip. Root in prepared trench, planting about 6 in apart. After one year the cuttings should have rooted and be making some growth, dig up and plant in permanent position and treat as new plants.

Big Bud is the curse of black currants and bushes should be sprayed with lime sulphur when the flowers have turned into the 'grape' stage, that is when the flowers have finished blooming and small purple bunches of forming fruit appear.





TRAVEL WEEKLY TO AND FROM SWITZERLAND AND ENGLAND

Providing DOOR TO DOOR SERVICES with vans specially equipped to carry

HOUSEHOLD REMOVALSEXHIBITION GOODSWORKS OF ART — MACHINERY

For full information contact:—
LEP PACKING LTD., CORNEY ROAD, CHISWICK, LONDON W4
TEL: 01-995 1300 (Ext. 110) TELEX: 23944