

Geometric and algebraic intersection numbers.

Autor(en): **Kervaire, Michel A.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **39 (1964-1965)**

PDF erstellt am: **24.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-29886>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Geometric and algebraic intersection numbers

by MICHEL A. KERVAIRE, New York (USA)

Let M be a connected $2n$ -dimensional differential manifold, not necessarily compact. Let $x_0 \in M$ be a base point, and $\alpha \in \pi_n(M, x_0)$ a given homotopy class. It is well known that, unless M is simply connected, there need not exist any differentiable imbedding $\varphi: S^n \rightarrow M$ representing α .

Let \bar{M} be the universal cover of M provided with an arbitrary but fixed orientation, and let $a \in H_n(\bar{M})$ be the (integral) homology class of a lifting of α .

Theorem 1. *Assuming $n > 2$, the class $\alpha \in \pi_n(M, x_0)$ is representable by a differentiable imbedding $\varphi: S^n \rightarrow M^{2n}$ if and only if for every covering transformation $\tau \neq 1$ of \bar{M} the homology intersection number $a \cdot \tau(a)$ vanishes.*

If M is oriented, one can define a scalar product

$$H_q(\bar{M}) \otimes H_{m-q}(\bar{M}) \rightarrow \mathbb{Z}[\pi]$$

with values in the integral group ring of $\pi = \pi_1(M, x_0)$. (Cf. K. REIDEMEISTER [2] and J. MILNOR [1].) Here $m = \dim M = \dim \bar{M}$ need not be even, and we assume that the projection map $p: \bar{M} \rightarrow M$ is orientation preserving. The image of $x \otimes y$ under the above pairing will be denoted as in MILNOR [1] by $[x, y]$.

In terms of this scalar product Theorem 1 can be formulated as follows:

Theorem 1'. *Let M^{2n} be connected and oriented. Assuming that $n > 2$, the class $\alpha \in \pi_n(M, x_0)$ is representable by a differentiable imbedding $\varphi: S^n \rightarrow M^{2n}$ if and only if*

$$[a, a] - a \cdot a = 0.$$

The proof is given in §1 and §2. In §3 we give conditions under which two imbeddings $\varphi: X^q \rightarrow M^m$ and $\psi: Y^{m-q} \rightarrow M^m$ representing the homology classes α, β respectively are diffeotopic to imbeddings φ_0, ψ_0 such that the cardinality of the set $\varphi_0(X^q) \cap \psi_0(Y^{m-q})$ equals the absolute value $|\alpha \cdot \beta|$ of the homology intersection number $\alpha \cdot \beta$. (Cf. Theorem 2 below.)

The proofs of Theorem 1 and Theorem 2 depend on the following well known lemma, essentially due to H. WHITNEY.

Let B^r denote the open unit ball in R^r .

Lemma. *Let V^m be a differential manifold, not necessarily compact, and let $\varphi: B^q \rightarrow V^m$ and $\psi: B^{m-q} \rightarrow V^m$ be two differentiable imbeddings such that $\varphi(B^q)$ and $\psi(B^{m-q})$ intersect transversally at exactly two points $R = \varphi(P) = \psi(Q)$ and $R' = \varphi(P') = \psi(Q')$.*

Suppose that

- (i) both q and $m - q$ are larger than 2,
- (ii) if $u: I \rightarrow B^q$ and $v: I \rightarrow B^{m-q}$ are paths from P to P' and Q to Q' respectively, then the loop $\varphi(u) \cdot \psi(v^{-1})$ is freely homotopic in V to a constant loop,
- (iii) with respect to some orientation of a neighborhood of $\varphi(B^q) \cup \psi(B^{m-q})$ in V the intersection coefficients of $\varphi(B^q)$ and $\psi(B^{m-q})$ at R and R' are opposite, i. e. $\varphi(B^q) \cdot \psi(B^{m-q}) = 0$.

Then there exists a diffeotopy $\varphi_t: B^q \rightarrow V^m$ such that $\varphi_1 = \varphi$, $\varphi_t(x)$ is independent of t for $|x| > 1 - \varepsilon$ for some positive ε and $\varphi_0(B^q) \cap \psi(B^{m-q}) = \emptyset$.

For a proof, see [3] and [4].

§ 1. Proof of Theorem 1

It is easy to see that $a \cdot \tau(a) = 0$ for all $\tau \neq 1$ is a necessary condition for the existence of an imbedding $\varphi: S^n \rightarrow M^{2n}$ representing $\alpha \in \pi_n(M, x_0)$. (Recall that a denotes the homology class of a lifting of α in the universal cover \overline{M} of M .) For let $\varphi: S^n \rightarrow M^{2n}$ be a mapping representing α , and $f: S^n \rightarrow \overline{M}$ a lifting of φ . Let $\tau: \overline{M} \rightarrow \overline{M}$ be a covering transformation, $\tau \neq 1$. We show that if P is a point in $f(S^n) \cap \tau f(S^n)$ then φ is not an imbedding. Let $Q = \tau^{-1}P \in f(S^n)$. Since $\tau \neq 1$, we have $Q \neq P$. Choose $Q', P' \in S^n$ such that $f(Q') = Q$ and $f(P') = P$. Then $Q' \neq P'$ but $\varphi(Q') = \varphi(P')$ since f is a lifting of φ . Hence φ is not bijective. Now, if φ is an imbedding, it follows that $f(S^n) \cap \tau f(S^n) = \emptyset$ for every $\tau \neq 1$, and a fortiori $a \cdot \tau(a) = 0$.

Conversely, suppose that $a \cdot \tau(a) = 0$ for every $\tau \neq 1$. Since \overline{M} is simply connected and $n > 2$, WHITNEY's lemma (cf. introduction) implies that a can be represented by a differentiable imbedding $f: S^n \rightarrow \overline{M}$. The projection map $p: \overline{M} \rightarrow M$ is an immersion. Hence f projects to an immersion $p \circ f = \varphi: S^n \rightarrow M$. We may assume without loss of generality that $\varphi(S^n)$ intersects itself transver-

sally in a finite number of points where only two sheets of $\varphi(S^n)$ cross each other. In other words, we may assume that φ is a completely regular immersion in the sense of [4]. (This can be obtained by an arbitrarily close approximation to φ in the C^2 -topology so that the new φ still lifts to an imbedding.)

Let S_φ be the set of pairs of (distinct) points (P, Q) on S^n such that $\varphi(P) = \varphi(Q)$. If $S_\varphi \neq \emptyset$, pick a pair $(P, Q) \in S_\varphi$. *Claim:* There exists another pair $(P', Q') \in S_\varphi$ and φ is regularly homotopic to an immersion $\Phi: S^n \rightarrow M$ such that

- (i) any lifting $F: S^n \rightarrow \bar{M}$ of Φ is an imbedding,
- (ii) $S_\Phi = S_\varphi - \{(P, Q), (P', Q')\}$,
- (iii) Φ and φ coincide outside some neighborhood of a path joining P to P' on S^n .

This will prove the theorem by induction on the number of self-intersection points.

We now prove the claim. Since $\varphi(P) = \varphi(Q)$ and $f(P) \neq f(Q)$, there exists a covering transformation $\tau \neq 1$ such that $f(P) = \tau f(Q) = A$, say. The point A is a transversal intersection point of $f(S^n)$ and $\tau f(S^n)$. Let $\varepsilon (\varepsilon = \pm 1)$ be the intersection coefficient. Since $a \cdot \tau(a) = f(S^n) \cdot \tau f(S^n) = 0$ by assumption, there exists another intersection point A' of $f(S^n)$ and $\tau f(S^n)$ with intersection coefficient $-\varepsilon$. Let $P', Q' \in S^n$ be such that $f(P') = \tau f(Q') = A'$, and let $u: I \rightarrow S^n$ be a path on S^n from P to P' , and similarly $v: I \rightarrow S^n$ a path on S^n from Q to Q' such that $u(I) \cap v(I) = \emptyset$. We may assume moreover that $u(I)$ and $v(I)$ are disjoint from the points of the pairs in S_φ except for $u(0) = P, u(1) = P', v(0) = Q$ and $v(1) = Q'$. Then $\varphi u(I)$ and $\varphi v(I)$ intersect only at $\varphi u(0) = \varphi v(0)$ and $\varphi u(1) = \varphi v(1)$. Since \bar{M} is simply connected, $f u$ and $\tau f v$ are two homotopic paths from A to A' . Hence φu and φv are homotopic paths on M from $\varphi(P) = \varphi(Q) = R$ to $\varphi(P') = \varphi(Q') = R'$. Take disjoint open neighborhoods N_u and N_v of $u(I)$ and $v(I)$ respectively with diffeomorphisms $h_u: B^n \rightarrow N_u$ and $h_v: B^n \rightarrow N_v$, and such that P, Q, P', Q' are the only points from S_φ in $N_u \cup N_v$. Let $V = M - \varphi(S^n - N_u \cup N_v)$ and set $\varphi_1 = \varphi h_u|_{B^n}$ and $\psi = \varphi h_v|_{B^n}$. We are now in a position to apply WHITNEY's lemma. The loop $\varphi_1 h_u^{-1}(u) \cdot \psi h_v^{-1}(v^{-1})$ is homotopic to a constant loop in V because it is homotopic to a constant loop in M and the inclusion $V \rightarrow M$ induces an isomorphism $\pi_1 V \cong \pi_1 M$. Thus φ_1 is diffeotopic in V , relative to a neighborhood of the boundary of D^n , to an imbedding $\varphi_0: B^n \rightarrow V$ such that $\varphi_0(B^n) \cap \psi(B^n) = \emptyset$. Define the immersion $\Phi: S^n \rightarrow M$ by

$$\Phi(x) = \begin{cases} \varphi_0 h_u^{-1}(x) & \text{if } x \in N_u, \text{ and} \\ \varphi(x) & \text{if } x \in S^n - N_u. \end{cases}$$

It is easily checked that Φ satisfies the conditions (ii) and (iii) of the above claim. To see that condition (i) stating that Φ lifts to an imbedding $F: S^n \rightarrow \overline{M}$ is also satisfied, let A, B be points on S^n with $F(A) = F(B)$. A path w from A to B on S^n maps to a loop Fw , and the loop Φw is homotopic to a constant loop in M . Since $\Phi(A) = \Phi(B)$ we have $\varphi(A) = \varphi(B)$ because the self-intersection points of Φ are self-intersection points of φ . In fact φw is also homotopic to a constant loop in M . (We may assume $\varphi w = \Phi w$ because, unless $A = B$, we have $A, B \in S^n - N_u$ and we can take $w(I) \subset S^n - N_u$.) But then, this means that $f(A) = f(B)$, and since f is an imbedding by construction, it follows that $A = B$. So F is bijective, and hence an imbedding. The proof of Theorem 1 is thus complete.

§ 2. The scalar product

Let M be a connected, oriented, differential manifold of dimension m , not necessarily even. Suppose M is triangulated as a regular cell complex. The triangulation of M lifts to a triangulation of \overline{M} invariant under the covering transformations. We denote by $\xi_i^q, i = 1, \dots, \alpha_q$ the q -cells of M and choose for each i a lifting x_i^q of ξ_i^q . Let η_i^{m-q} be the dual cell to ξ_i^q . ($\xi_i^q \cdot \eta_i^{m-q} = 1$.) A lifting y_j^{m-q} of η_j^{m-q} is then determined by the condition $x_i^q \cdot y_j^{m-q} = \delta_{ij}$. When we change the lifting x_i^q of ξ_i^q we demand that y_j^{m-q} be changed accordingly so that $x_i^q \cdot y_j^{m-q} = \delta_{ij}$ remains valid.

Let $\pi = \pi_1(M, x_0)$ be the fundamental group of M at x_0 which we identify with the group of covering transformations of \overline{M} . A q -dimensional chain x of (the triangulation of) \overline{M} has a unique expression as $x = \sum_i \lambda_i x_i^q$, where $\lambda_i \in Z[\pi]$ and almost all λ_i 's are zero. If $y = \sum_j \mu_j y_j^{m-q}$, define

$$[x, y] = [\sum_i \lambda_i x_i^q, \sum_j \mu_j y_j^{m-q}] = \sum_i \lambda_i \bar{\mu}_i,$$

where $\mu \rightarrow \bar{\mu}$ is the anti-ringhomomorphism of $Z[\pi]$ onto itself determined by $\tau \rightarrow \tau^{-1}$ for $\tau \in \pi$.

Theorem. *The scalar product $[x, y]$ is independent of the choice of the liftings x_i^q and induces a pairing*

$$H_q \overline{M} \otimes H_{m-q} \overline{M} \rightarrow Z[\pi].$$

The first statement follows by an easy calculation, using $\overline{\lambda \cdot \mu} = \bar{\mu} \cdot \bar{\lambda}$. The bilinearity of the product is obvious, and the second statement follows from

the formula $[x, dy] = \pm [dx, y]$, where $\dim x + \dim y = m + 1$, and d is the boundary $Z[\pi]$ -homomorphism. (Compare [1].)

We now derive a formula which will yield a translation of the condition " $a \cdot \tau(a) = 0$ for all $\tau \neq 1$ " in terms of the scalar product.

Let $a_0 = \sum_i \alpha_i x_i^q$, $\alpha_i \in Z[\pi]$, be a representative q -chain for $a \in H_q \bar{M}$ and $b_0 = \sum_j \beta_j y_j^{m-q}$ a representative of $b \in H_{m-q} \bar{M}$ in terms of the dual subdivision of \bar{M} . Then,

$$[a_0, b_0] = \sum_i \alpha_i \bar{\beta}_i.$$

We calculate $a_0 \cdot \tau(b_0)$, writing x_i and y_j for x_i^q and y_j^{m-q} respectively to simplify the notation. We have $\alpha_i = \sum_{\rho \in \pi} a_{i,\rho} \rho$ and $\beta_j = \sum_{\sigma \in \pi} b_{j,\sigma} \sigma$, where $a_{i,\rho}, b_{j,\sigma} \in Z[\pi]$ are almost all 0. Then,

$$a_0 \cdot \tau(b_0) = (\sum_{i,\rho} a_{i,\rho} \rho x_i) \cdot \tau(\sum_{j,\sigma} b_{j,\sigma} \sigma y_j) = \sum_{i,\rho} a_{i,\rho} b_{i,\tau^{-1}\rho},$$

and

$$\sum_i \alpha_i \bar{\beta}_i = \sum_{i,\rho,\sigma} a_{i,\rho} b_{i,\sigma} \rho \sigma^{-1} = \sum_{\tau} (\sum_{i,\rho} a_{i,\rho} b_{i,\tau^{-1}\rho}) \tau.$$

In other words, the integer $a_0 \cdot \tau(b_0)$ is just the coefficient of τ in the scalar product $[a_0, b_0]$. In formula,

$$(*) \quad [a, b] = \sum_{\tau \in \pi} (a \cdot \tau(b)) \tau.$$

(Compare MILNOR [1] where (*) is taken as definition of $[a, b]$.)

Now, if $m = 2n, q = n$ and $a = b$, it follows that the condition " $a \cdot \tau(a) = 0$ for all $\tau \neq 1$ " is equivalent to

$$[a, a] - a \cdot a = 0.$$

This proves Theorem 1'.

Remarks. (1) If we replace a by τa in Theorem 1', $[a, a] - a \cdot a$ becomes $\tau([a, a] - a \cdot a) \tau^{-1}$. Hence, only the conjugacy class of $[a, a] - a \cdot a$ is well determined by α . It seems therefore adequate to choose once and for all a base point $z_0 \in \bar{M}$ above $x_0 \in M$ and require all liftings to be liftings at z_0 . We then have a function

$$Q : \pi_n(M, x_0) \rightarrow Z[\pi],$$

$Q(\alpha) = [a, a] - a \cdot a$, and α is representable by an imbedded n -sphere if and only if $Q(\alpha) = 0$. The function Q is somewhat like a quadratic form. Let

$\langle \alpha, \beta \rangle = [a, b] + (-1)^n \overline{[a, b]}$ and let λ_1 denote the coefficient of $1 \in \pi$ in $\lambda \in Z[\pi]$. Then

$$Q(\tau\alpha) = \tau Q(\alpha) \overline{\tau} \quad (\tau \in \pi),$$

$$Q(\alpha + \beta) = Q(\alpha) + Q(\beta) + B(\alpha, \beta),$$

where $B(\alpha, \beta) = \langle \alpha, \beta \rangle - \langle \alpha, \beta \rangle_1$ is bilinear and symmetric.

(2) We have proved a slightly stronger statement than Theorem 1'. A class $\alpha \in H_n M^{2n}$ is representable by an imbedded manifold $A^n \subset M^{2n}$ such that $\pi_1 A \rightarrow \pi_1 M$ is trivial if and only if α is the projection of a class $a \in H_n(\overline{M})$ which is representable by a submanifold $B^n \subset \overline{M}$ and $[a, a] - a \cdot a = 0$. This gives a hint for the study of the intersection of submanifolds of M in §3.

Example. Take M^{2n} to be the connected sum of $S^1 \times S^{2n-1}$ with k copies of $S^n \times S^n$. Let $x_i, y_i \in \pi_n(M, x_0)$ be represented by $S^n \times (\text{point})$ and $(\text{point}) \times S^n$ in the i -th copy of $S^n \times S^n$ (suitably joined to the base point x_0). Then, $\pi_n(M, x_0)$ is the free $Z[J]$ -module generated by $x_1, \dots, x_k, y_1, \dots, y_k$, where J denotes the (multiplicative) infinite cyclic group. If $\alpha = \sum_i \alpha_i x_i + \sum_j \beta_j y_j \in \pi_n(M, x_0)$ is a given homotopy class, where $\alpha_i, \beta_j \in Z[J]$, it follows from the formula (*) above that

$$Q(\alpha) = \sum_i (\alpha_i \overline{\beta_i} + (-1)^n \beta_i \overline{\alpha_i}) - \sum_i (\alpha_i \overline{\beta_i} + (-1)^n \beta_i \overline{\alpha_i})^0,$$

where λ^0 is the integer obtained by substituting the value 1 for every element of J in $\lambda \in Z[J]$.

For instance, if we let t denote a generator of J , the class $x + (t + t^{-1})y$ in $\pi_n M$, where $M = S^1 \times S^{2n-1} \# S^n \times S^n$, is representable by a differentiable imbedding of S^n into M if n is odd, but is not representable if n is even. The same statement holds for $x + (t - t^{-1})y$ interverting even and odd.

§ 3. Reducing the geometric intersection of submanifolds

Let $\varphi : X^q \rightarrow M^m$ and $\psi : Y^{m-q} \rightarrow M^m$ be differentiable immersions, resp. imbeddings, where X, Y, M are connected differential manifolds. We assume X, Y to be compact, without boundary.

Roughly speaking, the problem is to use a deformation of φ so as to reduce the intersection $\varphi(X) \cap \psi(Y)$ to consist of a number of points equal to the algebraic intersection number of $\varphi(X)$ and $\psi(Y)$.

We assume that M is oriented. There are then two cases:

Case 1. X and Y are oriented. Denoting by $\alpha \in H_q M$ and $\beta \in H_{m-q} M$ the integral homology classes represented by $\varphi: X^q \rightarrow M$ and $\psi: Y^{m-q} \rightarrow M$ respectively, the algebraic (homology) intersection number $\alpha \cdot \beta$ is then an integer.

Case 2. At least one of the manifolds X, Y is non-orientable. Then $\varphi: X^q \rightarrow M$ and $\psi: Y^{m-q} \rightarrow M$ still represent mod 2 homology classes $\alpha \in H_q(M; \mathbb{Z}_2)$ and $\beta \in H_{m-q}(M; \mathbb{Z}_2)$. In this case the intersection number $\alpha \cdot \beta$ is an integer modulo 2.

We use $|\alpha \cdot \beta|$ to mean the absolute value in Case 1. In Case 2, $|\alpha \cdot \beta|$ is the integer 0 or 1 depending on whether $\alpha \cdot \beta = 0 \in \mathbb{Z}_2$ or $\alpha \cdot \beta = 1 \in \mathbb{Z}_2$.

In either case we shall assume that φ and ψ satisfy the following hypothesis:

(H) *The induced homomorphisms $\varphi_*: \pi_1 X \rightarrow \pi_1 M$ and $\psi_*: \pi_1 Y \rightarrow \pi_1 M$ are trivial.*

It follows that φ and ψ can be lifted to differentiable immersions, resp. imbeddings $f: X^q \rightarrow \bar{M}$ and $g: Y^{m-q} \rightarrow \bar{M}$, where as before \bar{M} is the universal cover of M . We let a and b denote the homology classes represented by f and g . In Case 1, $a \in H_q \bar{M}$ and $b \in H_{m-q} \bar{M}$. In Case 2, $a \in H_q(\bar{M}; \mathbb{Z}_2)$ and $b \in H_{m-q}(\bar{M}; \mathbb{Z}_2)$.

We also assume

$$(H') \quad q > 2 \quad \text{and} \quad m - q > 2.$$

Finally, we use the following notation. If $\lambda \in Z[\pi]$, $\lambda = \sum_{\tau} n_{\tau} \tau$, set $w \lambda = \sum_{\tau} |n_{\tau}|$.

Theorem 2. *With the above notations and hypotheses, including (H) and (H'), the immersion, resp. imbedding $\varphi: X^q \rightarrow M^m$ is regularly homotopic, resp. diffeotopic to an immersion, resp. imbedding $\varphi_0: X^q \rightarrow M^m$ such that $\varphi_0(X) \cap \psi(Y)$ consists of $|\alpha \cdot \beta|$ points if and only if*

$$\text{in Case 1, } w[a, b] - |\alpha \cdot \beta| = 0,$$

$$\text{in Case 2, } [a, b] - a \cdot b = 0 \text{ in } \mathbb{Z}_2[\pi] \text{ for some liftings } a, b \text{ of } \alpha, \beta.$$

Remark. Observe that

$$\alpha \cdot \beta = \sum_{\tau \in \pi} a \cdot \tau(b).$$

Thus, in view of (*), the equation $w[a, b] - |\alpha \cdot \beta| = 0$ in Case 1 is equivalent to the statement that $a \cdot \tau(b)$ does not change sign as τ runs over π . (More precisely, $(a \cdot \sigma b)(a \cdot \tau b) \geq 0$ for all $\sigma, \tau \in \pi$.) Obviously the condition is independent of the choice of liftings.

An important special case in practice seems to be Case 1 when $|\alpha \cdot \beta| = 0$ or 1. Then, $w[a, b] - |\alpha \cdot \beta| = 0$ is equivalent to the nicer looking condition

$$[a, b] - a \cdot b = 0 \text{ for some liftings } a, b \text{ of } \alpha, \beta.$$

(However, in general this last condition is definitely stronger than the former.)

As an illustration to the theorem, let $M = S^1 \times S^{2n-1} \# S^n \times S^n$, and $x, y \in \pi_n(M, x_0)$ be the elements represented by $S^n \times Q$ and $P \times S^n$ respectively, with $x_0 = P \times Q$. Let $u, v \in \pi_n(M, x_0)$ be given elements and A be the 2 by 2 matrix over $Z[J]$ such that $\begin{pmatrix} u \\ v \end{pmatrix} = A \begin{pmatrix} x \\ y \end{pmatrix}$, where J is the multiplicative infinite cyclic group.

Suppose that $u \cdot v = 1$, and let I_n denote the matrix

$$I_n = \begin{pmatrix} 0 & 1 \\ (-1)^n & 0 \end{pmatrix}.$$

The classes u and v can be represented by imbedded spheres with just one intersection point if and only if

$$I_n A I_n^* A^* = \begin{pmatrix} t^r & 0 \\ 0 & t^{-r} \end{pmatrix}$$

for some integer r . (t denotes a generator of J and A^* is the conjugate transposed of $A = (a_{ij})$, i. e. $A^* = (a_{ij}^*)$, where $a_{ij}^* = \overline{a_{ji}}$.)

Proof of Theorem 2. We may assume that $\varphi(X)$ and $\psi(Y)$ intersect transversally in a finite number of points S_1, \dots, S_k and $\varphi^{-1}(S_i), \psi^{-1}(S_i)$ each consists of a single point for every $i = 1, \dots, k$. Let $f: X \rightarrow \overline{M}$ and $g: Y \rightarrow \overline{M}$ be arbitrary liftings of φ and ψ respectively. (Hypothesis (H).)

Case 1. The manifolds X and Y are oriented. If $\tau \in \pi$, we have $f(X) \cdot \tau g(Y) = \varepsilon_\tau |a \cdot \tau(b)|$ for some $\varepsilon_\tau = \pm 1$. For each $\tau \in \pi$ we can select intersection points $R_{\tau,j}, j = 1, \dots, |a \cdot \tau(b)|$, of $f(X)$ and $\tau g(Y)$ so that the intersection coefficient of $f(X)$ and $\tau g(Y)$ at $R_{\tau,j}$ is equal to ε_τ , and thus independent of j . (If $a \cdot \tau(b) = 0$, the set $\{R_{\tau,j}\}$ is empty.) Let $(P_{\tau,j}, Q_{\tau,j}) \in X \times Y$ be the uniquely determined pair such that $f(P_{\tau,j}) = \tau g(Q_{\tau,j}) = R_{\tau,j}$.

Now, let $(P, Q) \in X \times Y$ be such that $\varphi(P) = \psi(Q)$ and $P \neq P_{\tau,j}$ for all τ, j , if any such pair exists. Then, there exists a covering transformation $\sigma \in \pi$ such that $f(P) = \sigma g(Q) = R$, say, and $R \neq R_{\sigma,i}$ for all i . ($1 \leq i \leq |a \cdot \sigma b|$.) Actually $R \neq R_{\tau,j}$ for all τ, j . Since $f(X) \cdot \sigma g(Y) = \varepsilon_\sigma |a \cdot \sigma(b)|$, there must exist another pair $(P', Q') \in X \times Y$ with $f(P') = \sigma g(Q') = R'$, where $R' \neq R_{\sigma,i}$ for all i and the intersection coefficients of $f(X)$ and $\sigma g(Y)$ at

R and R' are opposite. Again, since $\psi^{-1}(pR')$ consists of a single point, we actually have $R' \neq R_{\tau,j}$ for all τ, j .

We choose a path $u : I \rightarrow X$ from P to P' such that $u(I)$ and some neighborhood N_u of $u(I)$ in X are disjoint from any other φ -preimage of an intersection point of $\varphi(X)$ and $\psi(Y)$. Using WHITNEY's lemma as in §1, we can eliminate the intersection points $\varphi(P) = \psi(Q)$ and $\varphi(P') = \psi(Q')$ by a diffeotopy of $\varphi|N_u$ keeping φ fixed near the boundary of N_u .

It follows that φ is always (i. e. without condition on $[a, b]$) regularly homotopic, resp. diffeotopic, to an immersion, resp. an imbedding φ_0 such that

$$|\varphi_0(X) \cap \psi(Y)| = \sum_{\tau} |a \cdot \tau(b)|,$$

where $|\varphi_0(X) \cap \psi(Y)|$ denotes the cardinality of the finite set $\varphi_0(X) \cap \psi(Y)$.

If $a \cdot \tau(b)$ does not change sign, we then have

$$|\varphi_0(X) \cap \psi(Y)| = |\sum_{\tau} a \cdot \tau(b)| = |\alpha \cdot \beta|.$$

Conversely, if $|\varphi_0(X) \cap \psi(Y)| = |\alpha \cdot \beta|$, then $|\sum_{\tau} a \cdot \tau(b)| = \sum_{\tau} |a \cdot \tau(b)|$, and it follows that $a \cdot \tau(b) = \varepsilon |a \cdot \tau(b)|$ for all τ , where $\varepsilon = \pm 1$ is independent of τ .

Case 2. The manifold X , say, is non-orientable. Then, $a \in H_q(\bar{M}; \mathbb{Z}_2)$ and we also regard b as a class in $H_{m-q}(\bar{M}; \mathbb{Z}_2)$. For those $\tau \in \pi$ such that $a \cdot \tau(b) \neq 0 \pmod{2}$, we choose an intersection point R_{τ} of $f(X)$ and $\tau g(Y)$, and let $(P_{\tau}, Q_{\tau}) \in X \times Y$ be the uniquely determined pair such that $f(P_{\tau}) = \tau g(Q_{\tau}) = R_{\tau}$. Let $(P, Q) \in X \times Y$ be a pair such that $\varphi(P) = \psi(Q)$ and $P \neq P_{\tau}$ for all $\tau \in \pi$. Then $f(P) = \sigma g(Q) = R$, say, for some $\sigma \in \pi$, and since either P_{σ} does not exist or $P \neq P_{\sigma}$, there exist another pair (P', Q') with $f(P') = \sigma g(Q') = R'$, say, where $P' \neq P_{\tau}$ for all $\tau \in \pi$. Let v be a path on Y from Q to Q' such that $v(I) \cap \{Q_{\tau}\} = \emptyset$. Choose an orientation of a neighborhood N_v of $v(I)$ in Y . Since X is non-orientable, there exists a path u in X from P to P' with $u(I) \cap \{P_{\tau}\} = \emptyset$, and an orientation of a neighborhood N_u of $u(I)$ in X such that the intersection number of $\varphi(N_u)$ and $\psi(N_v)$ is the integer 0. We can then use WHITNEY's lemma again, as in §1, and eliminate the pairs (P, Q) and (P', Q') from the set of intersection pairs by a diffeotopy of $\varphi|N_u$ keeping φ fixed near the boundary of N_u . Thus φ can be replaced by φ_0 such that

$$|\varphi_0(X) \cap \psi(Y)| = \sum_{\tau} |a \cdot \tau(b)|,$$

where $|a \cdot \tau(b)|$ is the integer 0 if $a \cdot \tau(b) = 0 \pmod{2}$ and the integer 1 if $a \cdot \tau(b) = 1 \pmod{2}$.

If $[a, b] = a \cdot b$ in $Z_2[\pi]$ for some liftings a, b , then $a \cdot \tau(b) = 0$ for $\tau \neq 1$, and therefore, using these liftings in the above argument, $|\varphi_0(X) \cap \psi(Y)|$ is equal to 0 or 1 depending on whether $a \cdot b = 0$ or 1 mod 2 respectively. In other words, $|\varphi_0(X) \cap \psi(Y)| = |a \cdot b|$. Since $a \cdot \tau(b) = 0$ for $\tau \neq 1$ implies $\alpha \cdot \beta = \sum_{\tau} a \cdot \tau(b) = a \cdot b$, we have $|\varphi_0(X) \cap \psi(Y)| = |\alpha \cdot \beta|$ as desired.

Conversely, if $\varphi(X) \cap \psi(Y) = \emptyset$, we obviously have $[a, b] - a \cdot b = 0$ for any liftings of φ and ψ since both terms are then 0. If $|\varphi(X) \cap \psi(Y)| = 1$, we take a, b to be the classes of liftings of φ and ψ whose images intersect each other. Then $a \cdot \tau(b) = \delta_{\tau, 1}$. Hence $[a, b] - a \cdot b = 0$ in this case too. This completes the proof of Theorem 2.

Courant Institute of Mathematical Sciences, New York University

REFERENCES

- [1] MILNOR J., *A duality theorem for Reidemeister torsion*. Ann. of Math., 76 (1962), 137–147.
- [2] REIDEMEISTER K., *Durchschnitt und Schnitt von Homotopieketten*. Monatsh. Math. 48, (1939), 226–239.
- [3] SHAPIRO A., *Obstructions to the imbedding of a complex in a euclidean space. I. The first obstruction*. Ann. of Math., 66 (1957), 256–269.
- [4] WHITNEY H., *The self-intersections of a smooth n -manifold in $2n$ -space*. Ann. of Math., 45 (1944), 220–246.

(Reçu le 20 octobre 1964)