

Reflections in abstract Coxeter groups

Autor(en): **Franzsen, W.N. / Howlett, R.B. / Mühlherr, B.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **81 (2006)**

PDF erstellt am: **26.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-1177>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Reflections in abstract Coxeter groups

W. N. Franzsen, R. B. Howlett and B. Mühlherr

Abstract. Let W be a Coxeter group and $r \in W$ a reflection. If the group of order 2 generated by r is the intersection of all the maximal finite subgroups of W that contain it, then any isomorphism from W to a Coxeter group W' must take r to a reflection in W' . The aim of this paper is to show how to determine, by inspection of the Coxeter graph, the intersection of the maximal finite subgroups containing r . In particular we show that the condition above is satisfied whenever W is infinite and irreducible, and has the property that all rank two parabolic subgroups are finite. So in this case all isomorphisms map reflections to reflections.

Mathematics Subject Classification (2000). 20F55; 51F15.

Keywords. Coxeter group, isomorphism problem, automorphism, reflection.

1. Introduction

The dihedral group of order 12 can be considered as Coxeter group of type $I_2(6)$ or as Coxeter group of type $A_1 \times I_2(3)$. This example shows that, in general, the set of reflections in a Coxeter system is not determined by the abstract group W alone, but does indeed depend on the choice of the Coxeter generating set R . However there are a lot of examples of Coxeter systems (W, R) where the abstract group does determine the set of reflections or even the set R up to W -conjugacy. The main motivation for the present paper is to show that the latter holds for infinite Coxeter groups having a finite, irreducible and 2-spherical Coxeter generating set, which is our Theorem 1 below.

In view of the main result of [5] it suffices to show that these Coxeter groups determine the set of reflections. In order to achieve this goal we provide a handy criterion for an involution in an abstract Coxeter group W to be a reflection with respect to any Coxeter generating set of W . Our principal observation is the following. Let (W, R) be a Coxeter system and let $w \in W$ be an involution. If $w \notin R^W$, then the centralizer of w in W contains a finite normal subgroup properly containing $\langle w \rangle$. This is an immediate consequence of Richardson's result in [16]. Thus, if $w \in W$ is an involution having the property that $\langle w \rangle$ is a maximal finite normal subgroup of its

centralizer in W , then w is a reflection with respect to any Coxeter generating set of W .

It turns out that it is more convenient to work with the *finite continuation* of an involution rather than to consider finite normal subgroups of its centralizer. The finite continuation of a finite order element w in a Coxeter group is defined to be the intersection of all maximal finite subgroups containing it; we write $\text{FC}(w)$ for the finite continuation of w . In this paper we restrict our attention to finitely generated Coxeter groups. For these it is a consequence of a result of Tits that every element of finite order is contained in some maximal finite subgroup; so $\text{FC}(w)$ is a finite subgroup of W (see Corollary 14 below). The main result of the present paper is a complete description of the finite continuation of a simple reflection in a Coxeter system of finite rank. Its proof constitutes the bulk of this paper.

Main Result. *Let (W, R) be a Coxeter system of finite rank. Then the following holds.*

- a) *For each $r \in R$ the finite continuation of r can be described.*
- b) *Given an involution $w \in W$ such that $\text{FC}(w) = \langle w \rangle$, then $w \in R^W$.*

Part a) of our main result is Theorem 7. Its precise statement requires some preparation. Part b) is Corollary 24.

The main result of this paper is in fact the first of two steps to reduce the isomorphism problem for Coxeter groups to its ‘reflection-preserving’ version. The second step is given in [12]. We refer to [15] for further information about the applications to the general isomorphism problem.

A special instance of the isomorphism problem for Coxeter groups is the question about their rigidity (see [3] for further information). In combination with the main result of [5] a consequence of our main result is the following rigidity result.

Theorem 1. *Let (W, R) be an irreducible, non-spherical Coxeter system such that R is finite and such that rr' has finite order for all $r, r' \in R$. Then the following assertions hold.*

- a) *For each $r \in R$ we have $\text{FC}(r) = \langle r \rangle$.*
- b) *If $S \subseteq W$ is such that (W, S) is a Coxeter system, then there exists $w \in W$ such that $S^w = R$.*
- c) *All automorphisms of W are inner-by-graph.*

In the language of [3], Part b) of the previous theorem means that an infinite, irreducible, 2-spherical Coxeter system is strongly rigid. Part c), which is an immediate consequence of Part b), improves the result of [13].

To conclude this introduction we remark that characterization results for reflections in even Coxeter groups have been obtained in [1]. Some of the results there can be deduced as corollaries of our main result as well.

Acknowledgements. The authors thank Frédéric Haglund for helpful discussions on the subject, and the Australian Research Council for support. The third author thanks the University of Sydney for hospitality when this research was undertaken.

2. Precise statement of the main result

Recall that a Coxeter group is a group with a presentation of the form

$$W = \text{gp}\langle \{r_a \mid a \in \Pi\} \mid (r_a r_b)^{m_{ab}} = 1 \text{ for all } a, b \in \Pi \rangle \quad (2.1)$$

where Π is some indexing set, whose cardinality is called the *rank* of W (relative to this presentation), and the m_{ab} satisfy the following conditions: $m_{ab} = m_{ba}$, each m_{ab} lies in the set $\{m \in \mathbb{Z} \mid m \geq 1\} \cup \{\infty\}$, and $m_{ab} = 1$ if and only if $a = b$. When $m_{ab} = \infty$ the relation $(r_a r_b)^{m_{ab}} = 1$ is interpreted as vacuous. We shall restrict attention to Coxeter groups of finite rank.

A *reduced expression* for an element $w \in W$ is a minimal length word expressing w as a product of elements of the distinguished generating set $\{r_a \mid a \in \Pi\}$. We define $\ell(w)$ to be the length of a reduced expression for w .

As is well known (and as we shall describe in Section 3 below), every Coxeter group W can be realized geometrically as a group generated by reflections. In this realization of W the reflections in W are the conjugates of the generators r_a .

The *Coxeter graph* associated with the presentation above is the graph with vertex set Π and edge set consisting of those pairs of vertices $\{a, b\}$ for which $m_{ab} \geq 3$. The edge $\{a, b\}$ is given the label m_{ab} . The *components* of Π are the connected components of the graph, and we say that W is *irreducible* if the graph is connected.

For each $I \subseteq \Pi$ we define W_I to be the subgroup of W generated by the set $\{r_a \mid a \in I\}$; we call these subgroups the *visible* subgroups of W . A *parabolic subgroup* of W is any conjugate of a visible subgroup. We say that $I \subseteq \Pi$ is *spherical* if W_I is finite, and we say that Π (or W) is *k-spherical* if all k -element subsets of Π are spherical.

The definitions given so far are fairly standard. In order to facilitate the precise statement of the main result, we introduce some nonstandard notation and terminology (in Definitions 2, 3, 4, 5 and 6 below).

Definition 2. If $w \in W$ has finite order, define the *finite continuation* of w , written $\text{FC}(w)$, to be the intersection of all maximal finite subgroups of W containing w .

Definition 3. The *odd graph* of W is the graph $\Omega(\Pi)$ obtained from the Coxeter graph by deleting the edges whose labels are infinite or even. For each $a \in \Pi$ we define $\text{Odd}(a)$ to be the connected component of $\Omega(\Pi)$ containing a . For each connected component M of $\Omega(\Pi)$ we define $E(M)$ to be the union of M with the set of all $b \in \Pi$ such that m_{cb} is even for some $c \in M$. We also abbreviate $E(\text{Odd}(a))$ to $E\text{Odd}(a)$.

In the discussions below, when we refer to the components of $E(M)$ we regard $E(M)$ as the full subgraph of the Coxeter graph spanned by the vertices in $E(M)$. In other words, the edges with even and infinite labels, deleted when forming the odd graph, are restored in $E(M)$.

Note that if $a \in L \subseteq \Pi$ and W_L is finite then $m_{ab} < \infty$ for all $b \in L$. Whether m_{ab} is odd or even it follows that $b \in E\text{Odd}(a)$. Thus $L \subseteq E\text{Odd}(a)$.

Definition 4. Let $M \subseteq \Pi$ be a connected component of $\Omega(\Pi)$. We call $b \in \Pi \setminus M$ a C_3 -neighbour of M if $m_{bc} \in \{2, 4\}$ for all $c \in E(M)$, the case $m_{bc} = 4$ occurring for at least one c , and for each $c \in E(M)$ with $m_{bc} = 4$ there is an $a \in M$ such that the following conditions are satisfied:

- (1) $m_{ba} = 2$ and $m_{ca} = 3$, and $m_{cd} = \infty$ for all $d \in M \setminus \{a, c\}$;
- (2) for all $e \in \Pi \setminus (M \cup \{b\})$, either $m_{ce} = \infty$ or $m_{ae} = m_{ce} = m_{be} = 2$.

Definition 5. Let $M \subseteq \Pi$ be a connected component of $\Omega(\Pi)$, and let $a \in M$ and $b \in \Pi \setminus M$. We call the pair (a, b) a *focus* of M in Π if the following conditions all hold.

- (1) All the edge labels of M are 3, and M is a tree.
- (2) For each $c \in M$, the set $C[b..c] \subseteq \Pi$ consisting of b and those elements of M that form the path from a to c in M constitutes a system of type C_k (for some k dependent on c).
- (3) If $c, d \in M \cup \{b\}$ with $c \notin C[b..d]$ and $d \notin C[b..c]$ then $m_{cd} = \infty$.
- (4) If $m_{ce} \neq \infty$ for some $c \in M$ and $e \in \Pi \setminus (M \cup \{b\})$, then $m_{ce} = 2 = m_{de}$ for all $d \in C[b..c]$.
- (5) The vertices of $M \cup \{b\}$ do not form a spherical component of $E(M)$.

Definition 6. Let $M \subseteq \Pi$ be a connected component of $\Omega(\Pi)$, and let $a, b \in M$. We call the two-element set $\{a, b\}$ a *half focus* of M in Π if $m_{ab} = 2$ and the following conditions all hold.

- (1) We have $m_{ac} = m_{bc} \in \{2, 3\}$ for all $c \in M \setminus \{a, b\}$, and $m_{ac} = m_{bc} \in \{2, \infty\}$ for all $c \in \Pi \setminus M$.
- (2) All the edge labels of $M \setminus \{b\}$ are 3, and $M \setminus \{b\}$ is a tree.

- (3) For each $c \in M \setminus \{a, b\}$, the set $D[a, b..c] \subseteq \Pi$ consisting of b and those elements of $M \setminus \{b\}$ that form the path from a to c constitutes a system of type D_k (for some k dependent on c).
- (4) If $c, d \in M \setminus \{a, b\}$ with $c \notin D[a, b..d]$ and $d \notin D[a, b..c]$ then $m_{cd} = \infty$.
- (5) If $m_{ce} \neq \infty$ for some $c \in M \setminus \{a, b\}$ and $e \in \Pi \setminus M$, then $m_{ce} = 2 = m_{de}$ for all $d \in D[a, b..c]$.
- (6) The vertices of M do not form a spherical component of $E(M)$.

We are now able to give a precise statement of Part a) of our main result.

Theorem 7. *For each connected component M of $\Omega(\Pi)$ there is at least one $a \in M$ such that $\text{FC}(r_a)$ is a visible subgroup of W . We have the following possibilities.*

Case A: Suppose that the component of $E(M)$ containing M is spherical, and let $a \in M$ be arbitrary. Then $\text{FC}(r_a) = W_J$, where J is the union of the spherical components of $E(M)$.

Case B: Suppose that the component of $E(M)$ containing M is not spherical, and M does not have any focus or half-focus in Π , and let J' be the union of the spherical components of $E(M)$ and the set of C_3 -neighbours of M . If $a \in M$ is not adjacent in Π to any C_3 -neighbour of M then $\text{FC}(r_a) = W_{J' \cup \{a\}}$, and if $a \in M$ is adjacent in Π to a C_3 -neighbour of M then $\text{FC}(r_a)$ is not visible.

Case C: Suppose that (a, b) is a focus of M . Then $\text{FC}(r_a) = W_J$ where J is the union of $\{a, b\}$ and the spherical components of $E(M)$, and $\text{FC}(r_c)$ is not visible for any $c \in M \setminus \{a\}$.

Case D: Suppose that $\{a, b\}$ is a half-focus of M . Then $\text{FC}(r_a) = \text{FC}(r_b) = W_J$, where J is the union of $\{a, b\}$ and the spherical components of $E(M)$, and $\text{FC}(r_c)$ is not visible for any $c \in M \setminus \{a, b\}$.

3. Reflections and root systems

Let \mathbb{R} be the real field, and V the vector space over \mathbb{R} with basis Π . Let B the bilinear form on V such that for all $a, b \in \Pi$,

$$B(a, b) = -\cos(\pi/m_{ab}).$$

To make our notation more compact we define $u \cdot v = B(u, v)$ for all $u, v \in V$. Note that $a \cdot a = 1$ for all $a \in \Pi$, since $m_{aa} = 1$.

For each $a \in V$ such that $a \cdot a = 1$, the *reflection along a* is the transformation of V given by $v \mapsto v - 2(a \cdot v)a$. It is well known (see, for example, Corollary 5.4 of [14]) that W has a faithful representation on V such that, for all $a \in \Pi$, the

element r_a acts as the reflection along a . We shall identify elements of W with their images in this representation. We also use the notation r_a for the reflection along a whenever $a \in V$ satisfies $a \cdot a = 1$. It is straightforward to show that each reflection r_a preserves the form B ; hence all elements of W preserve B . Furthermore, the equation $gr_ag^{-1} = r_{ga}$ holds for all $a \in V$ such that $a \cdot a = 1$ and all transformations g that preserve B .

We write $\text{Ref}(W)$ for the set of all reflections in W . It is immediate from the above comments that if $\Phi = \{wa \mid w \in W, a \in \Pi\}$ then $\{r_b \mid b \in \Phi\} \subseteq \text{Ref}(W)$.

The set Φ is called the *root system* of W , and elements of Φ are called *roots*. Elements of the basis Π are called *simple roots*, and the reflections r_a for $a \in \Pi$ are called *simple reflections*. A root is said to be *positive* if it has the form $\sum_{a \in \Pi} \lambda_a a$ with $\lambda_a \geq 0$ for all $a \in \Pi$, and *negative* otherwise. We write Φ^+ for the set of all positive roots and Φ^- for the set of all negative roots.

Lemma 8. *With the notation as above, the following statements hold.*

- (1) *Every negative root has the form $\sum_{a \in \Pi} \lambda_a a$ with $\lambda_a \leq 0$ for all $a \in \Pi$. Furthermore, $\Phi^- = \{-b \mid b \in \Phi^+\}$.*
- (2) *If $w \in W$ and $a \in \Pi$ then*

$$\ell(wr_a) = \begin{cases} \ell(w) + 1 & \text{if } wa \in \Phi^+, \\ \ell(w) - 1 & \text{if } wa \in \Phi^-. \end{cases}$$

- (3) *If $t \in \text{Ref}(W)$ then $t = r_b$ for some $b \in \Phi$.*
- (4) *The group W is finite if and only if the bilinear form B is positive definite.*
- (5) *The root system Φ is finite if and only if the group W is finite.*

Proof. Proofs of (1) and (2) can be found in [14, Section 5.4], Theorem 4.1 in [7] includes both (4) and (5), and (3) is [13, Lemma 2.2]. \square

The following result is well known.

Lemma 9. *Let $a \in \Pi$. Then $\text{Odd}(a) = \Pi \cap Wa$.*

For each $w \in W$ we define $N(w) = \{b \in \Phi^+ \mid wb \in \Phi^-\}$. By Part (2) of Lemma 8, if $w \neq 1$ then $N(w) \cap \Pi \neq \emptyset$. An easy induction shows that $N(w)$ has exactly $\ell(w)$ elements. In particular, $N(w)$ is a finite set. It is also easily shown that if Φ is finite then there is a unique $w \in W$ such that $N(w) = \Phi^+$. This element, which we denote by w_Π , is also the unique element of maximal length in W (which is a finite group). Furthermore, $w_\Pi \Pi = -\Pi$.

For each $\Gamma \subseteq \Phi$ the subgroup W_Γ generated by the set $\{r_a \mid a \in \Gamma\}$ is called a *reflection subgroup* of W . The set $\Phi_\Gamma = \{a \in \Phi \mid r_a \in W_\Gamma\}$ is called the *root subsystem* generated by Γ . Let $\Phi_\Gamma^+ = \Phi_\Gamma \cap \Phi^+$ and $\Phi_\Gamma^- = \Phi_\Gamma \cap \Phi^-$, and define

$$\Pi_\Gamma = \{a \in \Phi_\Gamma^+ \mid N(r_a) \cap \Phi_\Gamma = \{a\}\}.$$

The main theorem of Deodhar [8] and Theorem (3.3) of Dyer [9] yield the following result.

Theorem 10. *For each $\Gamma \subseteq \Phi$ the group W_Γ is a Coxeter group on the generating set $\{r_a \mid a \in \Pi_\Gamma\}$. The set $\{a \cdot b \mid a, b \in \Pi_\Gamma \text{ and } a \neq b\}$ is a subset of $\mathcal{C} = \{-\cos(\pi/m) \mid 2 \leq m \in \mathbb{Z}\} \cup (-\infty, -1]$. Moreover, if Δ is any subset of Φ^+ such that $\{a \cdot b \mid a, b \in \Delta \text{ and } a \neq b\} \subseteq \mathcal{C}$ then W_Δ is a Coxeter group on the generating set $\{r_a \mid a \in \Delta\}$.*

Note that the notation W_Γ introduced above is an extension of the notation for visible subgroups introduced in Section 2. However, if $\Gamma \not\subseteq \Pi$ then W_Γ need not be visible.

It is clear that if $I \subseteq \Pi$ then W_I preserves the subspace V_I of V spanned by I , and acts on this subspace as a Coxeter group with I as its set of simple roots. In this case $\Phi_I = \Phi \cap V_I$ and $\Pi_I = I$.

The following simple facts are well known.

Lemma 11. *In the above situation, $\Phi_I = \Phi \cap V_I$. Furthermore, $w \in W$ normalizes W_I if and only if $w\Phi_I = \Phi_I$. In particular, for all $a \in \Phi$, the reflection r_a normalizes W_I if and only if $a \in \Phi_I$ or $a \cdot b = 0$ for all $b \in I$.*

Suppose that $I \subseteq \Pi$ and $a \in \Pi \setminus I$, and let L be the component of (the Coxeter graph of) $I \cup \{a\}$ to which a belongs. If W_L is finite we define $v[a, I] = w_L w_{L \setminus \{a\}}$. It is easily seen that $v[a, I]I \subseteq I \cup \{a\}$, and that $v[a, I]b = b$ for all $b \in I \setminus L$. In particular, $v[a, I]I \in \mathcal{J} = \{J \subseteq \Pi \mid J = wI \text{ for some } w \in W\}$. It was proved in [11] (for finite Coxeter groups) and in [7] (in the general case) that every element $w \in W$ satisfying $wI \subseteq \Pi$ can be expressed as a product of elements of the form $v[a, I']$, with $I' \in \mathcal{J}$ and $a \in \Pi \setminus I'$. That is,

$$w = v[a_1, I_1]v[a_2, I_2] \dots v[a_n, I_n] \quad (3.1)$$

for some I_i, a_i such that (for each i) the component of $I_i \cup \{a_i\}$ containing a_i corresponds to a finite visible subgroup, $v[a_i, I_i]I_i = I_{i-1}$ for $1 < i \leq n$, and $I_n = I$. Furthermore, the following result holds.

Proposition 12. *Let $I, J \subseteq \Pi$. Then $\{w \in W \mid wW_Iw^{-1} = W_J\} = N(J, I)W_I$, where $N(J, I) = \{w \in W \mid wI = J\}$. Furthermore, for each $w \in N(J, I)$ and each $a \in \Pi \cap N(w)$ there is an expression for w of the form (3.1) above, with $(a_n, I_n) = (a, I)$ and $\ell(w) = \sum_{i=1}^n \ell(v[a_i, I_i])$.*

The following lemma, which appears in [2, Exercise 2d, p. 130], is fundamental to all of our arguments.

Lemma 13 (Tits). *If W is a Coxeter group and $H \leq W$ is finite, then H is contained in a finite parabolic subgroup of W .*

One immediate consequence of Lemma 13 is that every maximal finite subgroup of a Coxeter group is parabolic. Another consequence of the previous lemma is that each finite subgroup of W is contained in a maximal finite parabolic subgroup. (Remember that we always assume that W is finitely generated.) Thus the set of maximal finite subgroups of W containing a given finite order element of W is not empty, and hence we have the following fact.

Corollary 14. *If $w \in W$ has finite order, then $\text{FC}(w)$ is a well-defined finite subgroup of W .*

Lemma 15 (Kilmoyer). *Let $I, J \subseteq \Pi$. Then every (W_I, W_J) double coset in W contains a unique element of minimal length; moreover, if d is the minimal length element of $W_I d W_J$ then $W_I \cap d W_J d^{-1} = W_K$, where $K = I \cap dJ$.*

Proof. See [6, Theorem 2.7.4]. □

Corollary 16. *The intersection of a finite number of parabolic subgroups is a parabolic subgroup.*

The following consequence of Lemmas 13 and 15 is proved in [10, Lemma 11].

Lemma 17. *If J is a maximal spherical subset of Π then W_J is a maximal finite subgroup of W . Furthermore, W_J is not conjugate to any other visible subgroup of W .*

Another important tool in our analysis of automorphisms is the classification of involutions in Coxeter groups, due to Richardson [16].

Proposition 18. *Suppose that $w \in W$ is an involution. Then there is a $t \in W$ and a spherical $I \subseteq \Pi$ such that $w = t^{-1} w_I t$ with $\ell(w) = \ell(w_I) + 2\ell(t)$, and w_I is central in W_I .*

Proof. See [10, Proposition 5]. □

Definition 19. We say that $I \subseteq \Pi$ is of (-1) -type if W_I is finite and w_I is central in W_I .

The reason for the terminology is that I is of (-1) -type if and only if there is an element of W_I that acts on V_I as multiplication by -1 .

We need the following lemma.

Lemma 20. *Suppose that $I, J \subset \Pi$ with I of (-1) -type, and suppose that $t \in W$ has the property that $tw_I t^{-1} \in W_J$. Then $tW_I t^{-1} \subseteq W_J$.*

Proof. Let $a \in I$. Then $w_I(a) = -a$, and so $(tw_I t^{-1})(ta) = -ta$, whence it follows that either ta or $-ta$ is in the set $N(tw_I t^{-1})$. But $N(tw_I t^{-1}) \subseteq \Phi_J$; so $ta \in \Phi_J$, and therefore $tr_a t^{-1} = r_{ta} \in W_J$. Since W_I is generated by $\{r_a \mid a \in I\}$, the result follows. \square

In particular, it follows from Lemma 20 that if I, J are both of (-1) -type and $tw_I t^{-1} = w_J$ then $tW_I t^{-1} = W_J$. Conversely, suppose that $tW_I t^{-1} = W_J$, so that in fact $dW_I d^{-1} = W_J$ for all d in $W_J t W_I$ (which equals tW_I). Taking d to be the shortest element in tW_I , Lemma 15 yields that $dI = J$, and hence $x \mapsto dx d^{-1}$ is a length-preserving isomorphism $W_I \rightarrow W_J$; consequently $dw_I d^{-1} = w_J$. If w_I, w_J are central in W_I, W_J we deduce that $tw_I t^{-1} = w_J$. So we have proved the following result.

Lemma 21. *Suppose that I, J are subsets of Π that are both of (-1) -type. Then $\{t \in W \mid tw_I t^{-1} = w_J\} = \{t \in W \mid tW_I t^{-1} = W_J\}$.*

Proposition 22. *Let $I \subset \Pi$ be of (-1) -type. Then $W_I \subseteq \text{FC}(w_I)$.*

Proof. Let F be a maximal finite subgroup of W such that $w_I \in F$. By Lemma 13 there exist $t \in W$ and $J \subseteq \Pi$ such that $tFt^{-1} = W_J$. By Lemma 20 and the fact that $w_I \in F$ it follows that $tW_I t^{-1} \subseteq W_J$. Hence $W_I \subseteq t^{-1}W_J t = F$. \square

Proposition 23. *Let W, W' be Coxeter groups of finite rank and $\alpha: W \rightarrow W'$ an isomorphism. Let Π be the set of simple roots corresponding to the distinguished generating set of W , and let $a \in \Pi$. If r_a^α is not a reflection in W' then the intersection of all maximal finite subgroups of W containing r_a is a parabolic subgroup of order greater than 2.*

Proof. Write Π' for the set of simple roots of W' . Observe that Lemma 13 and Corollary 16 trivially imply that $\text{FC}(r_a)$ is a parabolic subgroup of W .

Since r_a^α is not a reflection it follows from Proposition 18 that $r_a^\alpha = tw_I t^{-1}$ for some $t \in W'$ and $I \subseteq \Pi'$ of (-1) -type and of rank at least 2. Clearly $\text{FC}(r_a)^\alpha = t\text{FC}(w_I)t^{-1}$, and by Proposition 22 we know that $W_I \subseteq \text{FC}(w_I)$. Therefore $(tW_I t^{-1})^{\alpha^{-1}} \subseteq \text{FC}(r_a)$, so that $\text{FC}(r_a)$ has order greater than 2, as required. \square

Corollary 24. *Let $w \in W$ be an involution such that $\text{FC}(w) = \langle w \rangle$ and let $S \subseteq W$ be such that (W, S) is a Coxeter system. Then $w \in S^W$.*

4. The finite continuation of a reflection

Let r be a reflection in W . Replacing r by wrw^{-1} replaces $\text{FC}(r)$ by $w\text{FC}(r)w^{-1}$, and so choosing w suitably enables us to assume that $\text{FC}(r) = W_J$, a visible parabolic subgroup. Furthermore, replacing r by trt^{-1} for suitable $t \in W_J$ enables us to assume that $r = r_a$ for some $a \in J$. (Note that these observations yield the first assertion of Theorem 7.)

Proposition 25. *Let $a \in J \subseteq \Pi$, and suppose that W_J is the intersection of all maximal finite subgroups of W containing r_a . Then $\{w \in W \mid wr_aw^{-1} \in W_J\}$ is a subset of the normalizer of W_J in W . Thus each W -conjugate of r_a in W_J is $N_W(W_J)$ -conjugate to r_a , and $C_W(r_a) \subseteq N_W(W_J)$. Moreover, if $b \in \Pi \setminus J$ is such that $W_{J \cup \{b\}}$ is infinite then $m_{bc} = \infty$ for all $c \in J$ such that r_c is conjugate to r_a in W .*

Proof. Let \mathcal{S} be the set of all maximal finite subgroups of W containing r_a , so that $W_J = \text{FC}(r_a) = \bigcap_{F \in \mathcal{S}} F$. Suppose that $w \in W$ satisfies $wr_aw^{-1} \in W_J$, and let $F \in \mathcal{S}$. Then $wr_aw^{-1} \in W_J \subseteq F$, and so $r_a \in w^{-1}Fw$. Thus $w^{-1}Fw$ is a maximal finite subgroup of W containing r_a , whence $w^{-1}Fw \in \mathcal{S}$. So

$$\bigcap_{F \in \mathcal{S}} F \subseteq \bigcap_{F \in \mathcal{S}} w^{-1}Fw$$

and so $W_J \subseteq w^{-1}W_Jw$. Since W_J is finite it follows that $w \in N_W(W_J)$.

Suppose that $c \in J$ with $r_c = wr_aw^{-1}$ for some $w \in W$. Clearly $F \mapsto wFw^{-1}$ is a bijection from the set of maximal finite subgroups of W containing r_a to the set of maximal finite subgroups of W containing r_c , and so $\text{FC}(r_c) = w\text{FC}(r_a)w^{-1}$. But $w\text{FC}(r_a)w^{-1} = wW_Jw^{-1} = W_J$ by the first part of the proof, and so $\text{FC}(r_c) = W_J$. Now suppose that $b \in \Pi \setminus J$ with $m_{cb} < \infty$. Then $W_{\{c,b\}}$ is finite, and so contained in a maximal finite subgroup F . Since $r_c \in F$ we must have $\text{FC}(r_c) \subseteq F$. It follows that the finite group F contains both W_J and r_b , and therefore $W_{J \cup \{b\}}$ is finite. \square

Assume, as in Proposition 25, that $a \in J \subseteq \Pi$ and $W_J = \text{FC}(r_a)$, and suppose now that $J \neq \{a\}$. Suppose that $L \subseteq \Pi$ is such that $J \subseteq L$ and W_L is finite. Then W_L is a finite Coxeter group possessing a visible parabolic subgroup W_J of rank greater than 1 that is normalized by the centralizer of some simple reflection $r_a \in W_J$. Indeed, W_J is normalized by all $w \in W_L$ such that $wr_aw^{-1} \in W_J$. Equivalently, by Lemma 8 (3), $\{w \in W_L \mid wa \in \Phi_J\} \subseteq N_W(W_J)$. This is a very restrictive condition, which we now proceed to examine with a case-by-case investigation of the different types of finite Coxeter groups. For the course of this investigation, we can (and shall) assume that $L = \Pi$.

So we assume for now that W is a finite Coxeter group of rank n , and our aim is to find all examples of the following phenomenon: there exist $\{a\} \subsetneq J \subseteq \Pi$ such that

the set $Q = \{w \in W \mid wa \in \Phi_J\}$ is a subset of $N_W(W_J)$. We assume that $J \neq \Pi$, since the condition is trivially satisfied otherwise.

If $K \subseteq \Pi$ is a component of the Coxeter graph such that $J \cap K = \emptyset$ then W_K is a direct factor of $N_W(W_J)$; moreover, $Q = (Q \cap W_{\Pi \setminus K})W_K$. So removing K from the graph will have no bearing on whether or not the condition $Q \subseteq N_W(W_J)$ holds. So we assume that there are no such components of Π . Exactly the same comments apply for a component K of Π such that $K \subseteq J$. So we also assume that there are none of these.

Assume that $\{a\} \subsetneq J \subsetneq \Pi$ and $Q \subseteq N_W(W_J)$. Suppose that $K \subseteq \Pi$ is a component of the Coxeter graph such that $a \notin K$. Then $r_b a = a$ for all $b \in K$; so $r_b \in Q \subseteq N_W(W_J)$, and it follows that $r_b c \in \Phi_J$ whenever $c \in J$. If $b \cdot c \neq 0$ then b is in the support of $r_b c$, and so $r_b c \in \Phi_J$ implies $b \in J$. Since K is connected it follows that if K contains any element of J then $K \subseteq J$. So either $K \cap J = \emptyset$ or $K \subseteq J$. But we have assumed that there are no such components. So the component of Π that contains a is the only component; that is, Π is irreducible.

Observe that the group $\text{Stab}(a) = \{w \in W \mid wa = a\}$ is a subset of Q and hence of $N_W(W_J)$. Note also that $N_W(W_J) = \{w \in W \mid w\Phi_J = \Phi_J\}$, which is also the stabilizer of the subspace V_J (since V_J is the subspace spanned by Φ_J and $\Phi_J = V_J \cap \Phi$). Now $\text{Stab}(a)$ is a parabolic subgroup of W whose root system is $\Phi \cap a^\perp$, and the following table gives the type of this root system in all cases.

W	$\text{Stab}(a)$	W	$\text{Stab}(a)$
A_n	A_{n-2}	E_7	D_6
C_n	$C_{n-2} + A_1$	E_8	E_7
C_n	C_{n-1}	H_3	$A_1 + A_1$
D_n	$D_{n-2} + A_1$	H_4	H_3
F_4	C_3	$I_2(2k)$	A_1
E_6	A_5	$I_2(2k+1)$	\emptyset

(For C_n there are two W orbits of roots, giving two possibilities for $\text{Stab}(a)$. For F_4 and $I_2(2k)$ there are also two W -orbits of roots, but $\text{Stab}(a)$ has the same type of root system whichever orbit a belongs to.) Since each irreducible constituent of its root system spans an irreducible $\text{Stab}(a)$ -submodule of V , the table shows that as a $\text{Stab}(a)$ -module, V has composition length two or three or (in one case only) four: a itself spans a trivial $\text{Stab}(a)$ -submodule of dimension 1, and a^\perp is either irreducible of dimension $n-1$ (for types F_4 , E_6 , E_7 , E_8 , H_4 , $I_2(2k)$ and one of the C_n possibilities), or the direct sum of irreducibles of dimensions 1 and $n-2$ (for types A_n , C_n , D_n when $n > 4$, H_3 and $I_2(2k+1)$), or the direct sum of three irreducibles of dimension 1 (for type D_4). Furthermore, the summands of a^\perp are pairwise nonisomorphic as $\text{Stab}(a)$ -modules, since even if they are of the same type their centralizers in $\text{Stab}(a)$ are different.

Since $\{a\} \subsetneq J \subsetneq \Pi$ and V_J is $\text{Stab}(a)$ -invariant, we see that $a^\perp = (V_J \cap a^\perp) \oplus V_J^\perp$, with both summands nonzero $\text{Stab}(a)$ -modules. So Π is of type A_n , C_n , D_n or H_3 . Furthermore, except in type D_4 , the two direct summands of a^\perp are irreducible and not isomorphic, and are therefore the only proper $\text{Stab}(a)$ -submodules of a^\perp . We conclude that V_J is spanned by a and one of the summands of a^\perp , while V_J^\perp is the other summand. In type D_4 we similarly deduce that V_J is spanned by a and one or two of the three 1-dimensional summands of a^\perp , and, correspondingly, V_J^\perp is of either of type $A_1 + A_1$ or of type A_1 .

If Π is of type A_n then one of the summands of a^\perp is of type A_{n-2} while the other is a trivial 1-dimensional $\text{Stab}(a)$ -module. If V_J^\perp is of type A_{n-2} then V_J must be of type A_1 , since the orthogonal complement of a subsystem of type A_{n-2} in A_n contains only a rank 1 root system. This contradicts the assumption that $\{a\} \subsetneq J$. So J is of type $A_1 + A_{n-2}$. Since W_J is visible, we deduce that a is an end node of the A_n diagram, and the node adjacent to a is the unique simple root not in J . However, if $n > 3$ then the maximal length element of W is in Q but not in the normalizer of W_J . So $n = 3$ and $J = \{a, c\}$, where c is the other end node. It is readily checked that Q has 8 elements and coincides with $N_W(W_J)$ (which is generated by W_J and an element that interchanges a and c).

If Π is of type C_n then one summand of a^\perp is of type C_{n-2} and the other of type A_1 . The roots in the A_1 summand are in the same W -orbit as a . If V_J^\perp is the A_1 component of a^\perp then $V_J = (V_J^\perp)^\perp$ is of type $C_{n-2} + A_1$. This determines J uniquely, since W_J is visible. If $n \geq 4$ and w is the longest element of the visible parabolic subgroup of type A_{n-1} , then $-wa \in \Pi \setminus \{b\} = J$, but $w \notin N_W(W_J)$. This contradicts the fact that $Q \subseteq N_W(W_J)$. So $n = 3$, and the elements of J are the end nodes a, c of the C_3 diagram, the middle node b being in the same W -orbit as a . Since $\Phi_J = \{\pm a, \pm c\}$ and c is not in the same W -orbit as a and b we deduce that $Q = \{w \in W \mid wa = \pm a\}$. Furthermore, of the 6 roots in the W -orbit of c , only c and $-c$ are orthogonal to a . So if $wa = \pm a$ then $wc = \pm c$. Thus if $w \in Q$ then $w\Phi_J = \Phi_J$, as required.

Continuing the discussion of C_n , suppose now that V_J^\perp is the C_{n-2} component of a^\perp . Then $V_J = (V_J^\perp)^\perp$ is of type C_2 . Writing $J = \{a, b\}$, the fact that $\text{Stab}(a)$ is of type $A_1 + C_{n-2}$ means that it is b rather than a that is the end node of the C_n diagram. If we put $c = r_b a$ then $\{\pm c\}$ is the component of $\Phi \cap a^\perp$ of type A_1 . It follows that $\{\pm a\} = \{\pm r_b c\}$ is the A_1 -component of $\Phi \cap (r_b a)^\perp = \Phi \cap c^\perp$. We see that $\text{Stab}(a) = \langle r_c \rangle \times W'$ and $\text{Stab}(c) = \langle r_a \rangle \times W'$, where W' is a parabolic (not visible) subgroup of W of type C_{n-2} . Indeed, the root system of W' is $\Phi \cap V_J^\perp$. The roots in Φ_J that are in the same W -orbit as a are $\pm a$ and $\pm c$, and so

$$Q = \{1, r_a, r_b, r_b r_a\} \text{Stab}(a) = \{1, r_a, r_b, r_b r_a\} \{1, r_c\} W' = W_J W'.$$

Hence our requirement that Q stabilizes $\Phi_J = \{\pm a, \pm b\}$ is indeed satisfied.

If Π is of type D_n with $n > 4$ then one summand of a^\perp is of type D_{n-2} and the other of type A_1 . The roots orthogonal to a D_{n-2} subsystem form a system of type $A_1 + A_1$. There are in fact two W -orbits of parabolic $A_1 + A_1$ subsystems, and the orthogonal complement of a D_{n-2} is perhaps better thought of as type D_2 , since the visible parabolic in this orbit corresponds to the two nodes of the diagram that form the fork. So if V_J^\perp is the D_{n-2} summand of a^\perp then $J = \{a, b\}$ consists of the two nodes of valency 1 that are adjacent to c , the node of valency 3. A similar statement applies for D_4 in the case that V_J^\perp is of type $A_1 + A_1$. In both cases the element $w = r_c r_a r_b r_c \in W$ satisfies $wa = b$ and $wb = a$, and since $\Phi_J = \{\pm a, \pm b\}$ we see that $Q = \{1, r_a, w, wr_a\} \text{Stab}(a)$. But $\text{Stab}(a) = \langle r_b \rangle \times W'$ and $\text{Stab}(b) = \langle r_a \rangle \times W'$, where W' is the parabolic subgroup corresponding to the subspace V_J^\perp , and it follows readily that Q stabilizes $\Phi_J = \{\pm a, \pm b\}$, as required.

Continuing the discussion of D_n , where $n \geq 4$, suppose now that V_J^\perp is an A_1 component of a^\perp . Then $V_J = (V_J^\perp)^\perp$ is of type $A_1 + D_{n-2}$. But the maximal length element of a visible A_{n-1} subsystem containing a takes a to an element of Φ_J but does not normalize W_J . So our requirement that $Q \subseteq N_W(W_J)$ is not met.

Finally, suppose that Π is of type H_3 , so that $\text{Stab}(a)$ is of type $A_1 + A_1$. Then V_J^\perp is of type A_1 , and hence J is of type $A_1 + A_1$. Let $J = \{a, c\}$, and note that $c = wa$ for some $w \in W$. Since $N_W(W_J)$ is generated by W_J and the central involution of W , we see that c is not in the $N_W(W_J)$ -orbit of a . Hence the element w above is in Q but not in $N_W(W_J)$, and so our requirements are not met.

We have thus established the following result.

Proposition 26. *Let Π be the set of simple roots for the finite irreducible Coxeter group W , and suppose that $a \in J \subseteq \Pi$. Then $\{w \in W \mid wa \in \Phi_J\}$ is a subset of $N_W(W_J)$ if and only if one of the following situations occurs:*

- (1) $J = \{a\}$;
- (2) $J = \Pi$;
- (3) $\Pi = \{a, b, c\}$ is of type C_3 , with $m_{ac} = 3$ and $m_{cb} = 4$, and $J = \{a, b\}$ of type $A_1 + A_1$;
- (4) Π is of type C_n and $J = \{a, b\}$ is of type C_2 , with b an end node of Π ;
- (5) Π is of type D_n or A_3 , and $J = \{a, b\}$, where a and b are end nodes that are both adjacent to some $c \in \Pi$.

We return now to investigation of an arbitrary finite rank Coxeter group W . The next proposition is an immediate consequence of Proposition 26 and the discussion preceding it.

Proposition 27. *Let $a \in J \subseteq L \subseteq \Pi$, and suppose that the group W_L is finite and that $W_J = \text{FC}(r_a)$. Let J_0 be the component of J containing a and L_0 the component*

of L containing J_0 . Then every component of J that is not contained in L_0 is a component of L . Furthermore, if $\{a\} \neq J \cap L_0 \neq L_0$ then $J \cap L_0 = \{a, b\}$ for some b , and one of the following alternatives occurs:

- (1) $L_0 = \{a, c, b\}$ is of type C_3 , with $m_{ac} = 3$ and $m_{cb} = 4$;
- (2) L_0 is of type C_n for some $n \geq 3$, with b an end node and $J_0 = \{a, b\}$ of type C_2 ;
- (3) L_0 is of type A_3 or type D_n for some $n \geq 4$, the nodes a and b having valency 1 and sharing a common neighbour.

One of the ingredients of alternative (2) of Proposition 27 is that the component of $\text{FC}(r_a)$ containing a is of type C_2 . We shall see that when this situation arises, $\text{Odd}(a)$ has a focus in Π .

Proposition 28. *Suppose that $a \in J \subseteq \Pi$ with $W_J = \text{FC}(r_a)$, and let J_0 be the component of J containing a . Suppose that $J_0 = \{a, b\}$ is of type C_2 . Then either $\text{Odd}(a) \cup \{b\}$ is a spherical component of $\text{EOdd}(a)$, or else (a, b) is a focus of $\text{Odd}(a)$ in Π .*

Proof. We use induction on k to prove that for all $k \geq 2$, if $b = c_1$, $a = c_2, c_3, \dots, c_k$ are simple roots satisfying

- (1) $2 < m_{c_i c_{i+1}} < \infty$ for all $i \in \{1, 2, \dots, k-1\}$, and
- (2) c_1, c_2, \dots, c_k are distinct from each other,

then $\{c_1, c_2, \dots, c_k\}$ forms a system of type C_k . The case $k = 2$ is immediately true.

Suppose that $k > 2$. The inductive hypothesis tells us that $\{c_1, c_2, \dots, c_{k-1}\}$ is of type C_{k-1} . The element $w = v[c_{k-1}, \{c_{k-2}\}] \dots v[c_4, \{c_3\}]v[c_3, \{c_2\}]$ has the property that $wa = wc_2 = c_{k-1}$, and so if we write $d = wb$ then

$$r_d = wr_b w^{-1} \in w \text{FC}(r_a) w^{-1} = \text{FC}(c_{k-1}),$$

since it is given that $b \in \text{FC}(r_a)$. But $W_{\{c_{k-1}, c_k\}}$ is finite, and so it follows that $\{r_d, r_{c_{k-1}}, r_{c_k}\}$ generates a finite group. Now $d \cdot c_{k-1} = b \cdot a = -\cos(\pi/4)$ and $c_{k-1} \cdot c_k = -\cos(\pi/m)$ for some $m > 2$. If $m \geq 4$ then

$$c_k \cdot d = c_k \cdot \left(b + \sqrt{2} \sum_{i=2}^{k-1} c_i \right) \leq -\sqrt{2}(c_k \cdot c_{k-1}) \leq -1,$$

whence the reflection subgroup $W_{\{r_d, r_{c_k}\}}$ is infinite (by Theorem 10), a contradiction. So $m = 3$. If $m_{c_i c_k} > 2$ for any $i \in \{1, 2, \dots, k-2\}$ then

$$c_k \cdot d \leq \sqrt{2}(c_k \cdot c_{k-1}) + c_k \cdot c_i < -1,$$

again giving a contradiction. So $m_{c_k c_i} = 2$ for all $i \in \{1, 2, \dots, k-2\}$ and $m_{c_k c_{k-1}} = 3$, and since $\{c_1, c_2, \dots, c_{k-1}\}$ is a system of type C_{k-1} it follows that $\{c_1, c_2, \dots, c_k\}$ is a system of type C_k , as claimed.

If there were $c, d \in \text{Odd}(a)$ with $3 < m_{cd} < \infty$ then b together with a minimal length odd-labelled path from a to $\{c, d\}$ would yield $c_1, c_2, \dots, c_k \in \Pi$ satisfying (1) and (2) above and not forming a system of type C_k , contradicting the result proved above. The same argument yields a contradiction if $c \in \text{Odd}(a)$ and $d \in \Pi \setminus \text{Odd}(a)$ with $3 < m_{cd} < \infty$, unless $\{c, d\} = \{a, b\}$. So all edge labels in $\text{Odd}(a)$ are 3, if $c, d \in \text{Odd}(a)$ are not adjacent in $\text{Odd}(a)$ then $m_{cd} \in \{2, \infty\}$, and if $c \in \text{Odd}(a)$ and $d \in \Pi \setminus \text{Odd}(a)$ then $m_{cd} \in \{2, \infty\}$ unless $\{c, d\} = \{a, b\}$. Furthermore, any circuit in $\text{Odd}(a)$ would similarly yield a contradiction (by combining the circuit with a minimal finite-labelled path connecting it to b). So $\text{Odd}(a)$ is tree.

For each $c \in \text{Odd}(a)$ let $C[b..c] \subseteq \Pi$ consist of b and the unique path from a to c in $\text{Odd}(a)$. The discussion above shows that $C[b..c]$ is always of type C . Now suppose that $c \in \text{Odd}(a)$ and $e \in \Pi \setminus C[b..c]$ with $m_{ce} = 2$. Write $C[b..c] = \{c_1, c_2, \dots, c_k\}$, with $c_1 = b$ and $c_k = c$, and let $d = b + \sqrt{2} \sum_{i=2}^k c_i$. An argument similar to one used above shows that $r_d \in \text{FC}(c)$, and hence $W_{\{d, c, e\}}$ is finite. So $d \cdot e > -1$. If $c_i \cdot e \neq 0$ then $c_i \cdot e \leq -1/2$; so it follows that there is at most one i with $c_i \cdot e \neq 0$. Suppose, for a contradiction, that there is exactly one such i . If $i > 1$ then $d \cdot e = \sqrt{2}(c_i \cdot e)$, and so $c_i \cdot e > -1/\sqrt{2}$. Hence $m_{c_i e} = 3$, and $d \cdot e = -1/\sqrt{2}$. But this means that the edges $\{c, d\}$ and $\{d, e\}$ of the Coxeter graph of $\{d, c, e\}$ are both labelled 4, contradicting the fact that $W_{\{d, c, e\}}$ is finite. So we must have $i = 1$, and finiteness of $W_{\{c, d, e\}}$ forces $b \cdot e = d \cdot e = -1/2$. But now if we put $L = \{e\} \cup J$ then, in the notation of Proposition 27, we have that $L_0 = \{e, b, a\}$ is of type C_3 with $J \cap L_0 = \{b, a\}$ of type C_2 , and Proposition 27 shows that this is not possible. We conclude that if $e \in \Pi$ has the property that $m_{ce} = 2$ for some $c \in \text{Odd}(a)$ then $m_{de} = 2$ for all $d \in C[b..c]$. In particular, if $e \in \Pi \setminus (\text{Odd}(a) \cup \{b\})$ and $m_{ce} \neq \infty$ for some $c \in \text{Odd}(a)$ then $m_{ce} = 2$, as shown above, and so $m_{de} = 2$ for all $d \in C[b..c]$.

All that remains to prove now is that if $c, d \in \text{Odd}(a)$ with $c \notin C[b..d]$ and $d \notin C[b..c]$, then $m_{cd} = \infty$. Since c and d are not adjacent in $\text{Odd}(a)$ the only alternative is that $m_{cd} = 2$; so suppose, for a contradiction, that this holds. Choose the vertex $e \in \text{Odd}(a)$ on the (unique) path from c to d such that the distance from e to a is minimal. Let c', d' be the neighbours of e in the path from c to d , with c' between e and c and d' between e and d . Then $c' \in C[b..c]$, and since $m_{cd} = 2$ it follows that $m_{c'd} = 2$. Now since $d' \in C[b..d]$ and $m_{c'd} = 2$ it follows that $m_{c'd'} = 2$. Thus the set $L \subseteq \Pi$ consisting of c' and d' and the vertices on the path from a to e form a system of type D (or A_3 if $e = a$). So L is spherical, and since $b \in \text{FC}(r_a)$ it follows that $L \cup \{b\}$ is spherical also. But this is impossible since $L \cup \{b\}$ is connected, has an edge labelled 4 (namely, $\{b, a\}$), and has a vertex of valency 3 (namely e). \square

The situation of alternative (3) of Proposition 27 is very similar to that of alternative (2), and in this case it turns out that $\text{Odd}(a)$ has a half-focus in Π .

Proposition 29. *Suppose that $a \in J \subseteq \Pi$ with $W_J = \text{FC}(r_a)$ and $\{a\}$ a component of J , and suppose that $J \cap \text{Odd}(a) \neq \{a\}$. Then either $\text{Odd}(a)$ is a spherical component of $\text{EOdd}(a)$, or else there exists an element $b \in \text{Odd}(a)$ such that $\{a, b\}$ is a half-focus of $\text{Odd}(a)$ in Π .*

Proof. Let $b \in (J \cap \text{Odd}(a)) \setminus \{a\}$, and let $w \in W$ with $wa = b$. Then $w \in N_W(W_J)$, by Proposition 25, and so

$$\text{FC}(r_b) = \text{FC}(wr_a w^{-1}) = w \text{FC}(r_a) w^{-1} = w W_J w^{-1} = W_J.$$

Moreover, $w\Phi_J = \Phi_J$, and since $a \cdot c = 0$ for all $c \in \Phi_J \setminus \{a\}$, it follows that $wa \cdot d = 0$ for all $d \in \Phi_J \setminus \{wa\}$. So $\{b\}$ is a component of J . Note that $m_{ab} = 2$, since a and b are in different components of J .

Let $c \in \Pi \setminus \{a, b\}$, and suppose first of all that $2 < m_{bc} < \infty$. Since $\{b, c\}$ is spherical and $W_J = \text{FC}(r_b)$ it follows that $J \cup \{c\}$ is spherical. Let $L = J \cup \{c\}$ and let L_0 be the component of L containing a . By Proposition 27, every component of J that is not contained in L_0 is a component of L . But b is adjacent to c in L ; so $\{b\}$ is not a component of L , and it follows that $b \in L_0$. Now $\{a\} \neq J \cap L_0$, since $b \in J \cap L_0$, and $J \cap L_0 \neq L_0$, since $c \in L_0$ and $c \notin J$ (since $\{b\}$ is a component of J). Furthermore, the conditions of alternative (2) of Proposition 27 are not satisfied, since a and b are not adjacent in J . So either alternative (1) or alternative (3) must hold, and since c is the only element of L not in J it follows that $L_0 = \{a, c, b\}$, with $m_{ac} = 3$. But a symmetrical argument, with the roles of a and b reversed, shows that every $d \in \Pi$ with $2 < m_{ad} < \infty$ has the property that $m_{bd} = 3$. So $m_{ac} = m_{bc} = 3$, and $\{a, c, b\}$ is of type A_3 .

Now suppose that $m_{bc} = 2$. Again since $\{b, c\}$ is spherical it follows that $J \cup \{c\}$ is spherical, and so $m_{ac} < \infty$. If $m_{ac} > 2$ then, as we have just observed, it follows that $m_{bc} = 3$, contrary to our assumption that $m_{bc} = 2$. So $m_{ac} = m_{bc} = 2$, and we have now shown that whenever $m_{bc} < \infty$ we have $m_{ac} = m_{bc} \in \{2, 3\}$. Since a symmetrical argument gives the same conclusion whenever $m_{ac} < \infty$, we conclude also that $m_{ac} = \infty$ if and only if $m_{bc} = \infty$.

We now use induction on k to prove that for all $k \geq 3$, if $b = c_1$, $a = c_2, c_3, \dots, c_k$ are simple roots satisfying

- (1) $2 < m_{c_i c_{i+1}} < \infty$ for all $i \in \{2, 3, \dots, k-1\}$, and
- (2) c_1, c_2, \dots, c_k are distinct from each other,

then $\{c_1, c_2, \dots, c_k\}$ forms a system of type D_k or A_3 . The case $k = 3$ follows from what we have proved above.

Suppose that $k > 3$. The inductive hypothesis tells us that $\{c_1, c_2, \dots, c_{k-1}\}$ is of type D_{k-1} (or A_3 if $k = 4$). The element $w = v[c_{k-1}, \{c_{k-2}\}] \dots v[c_4, \{c_3\}]v[c_3, \{c_2\}]$ has the property that $wa = wc_2 = c_{k-1}$, and so if we write $d = wb$ then

$$r_d = wr_{c_1}w^{-1} \in w\text{FC}(r_a)w^{-1} = \text{FC}(c_{k-1}),$$

since it is given that $b \in \text{FC}(r_a)$. But $W_{\{c_{k-1}, c_k\}}$ is finite, and so it follows that $\{r_d, r_{c_{k-1}}, r_{c_k}\}$ generates a finite group. Now $d \cdot c_{k-1} = b \cdot a = 0$ and $c_{k-1} \cdot c_k = -\cos(\pi/m)$ for some $m > 2$. If $c_k \cdot c_i \neq 0$ for some $i \in \{1, 2, \dots, k-2\}$ then

$$c_k \cdot d = c_k \cdot \left(c_1 + c_2 + c_{k-1} + 2 \sum_{i=3}^{k-2} c_i \right) \leq -\left(\frac{1}{2} + \cos \frac{\pi}{m} \right) \leq -1,$$

whence the reflection subgroup $W_{\{r_d, r_{c_k}\}}$ is infinite (by Theorem 10), a contradiction. So $c_k \cdot d = c_{k-1} \cdot c_k = -\cos \frac{\pi}{m}$. Since the reflection subgroup generated by $\{r_d, r_{c_{k-1}}, r_{c_k}\}$ is finite it follows that $m = 3$. So we have shown that $m_{c_k c_{k-1}} = 3$ and $m_{c_k c_i} = 2$ for $i < k-1$, and since $\{c_1, c_2, \dots, c_{k-1}\}$ is a system of type D_{k-1} it follows that $\{c_1, c_2, \dots, c_k\}$ is a system of type D_k , as claimed.

Note that $\text{Odd}(a) \setminus \{b\}$ and $\text{Odd}(b) \setminus \{a\}$ are both connected, since each element $c \in \text{Odd}(a)$ that is adjacent to a is also adjacent to b , and vice versa. If there were $c, d \in \text{Odd}(a) \setminus \{b\}$ with $3 < m_{cd} < \infty$ then b together with a minimal length odd-labelled path from a to $\{c, d\}$ would yield $c_1, c_2, \dots, c_k \in \Pi$ satisfying (1) and (2) above and not forming a system of type D_k , contradicting the result proved above. The same argument yields a contradiction whenever $c \in \text{Odd}(a) \setminus \{a, b\}$ and $d \in \Pi \setminus \text{Odd}(a)$ with $3 < m_{cd} < \infty$. So all edge labels in $\text{Odd}(a)$ are 3, if $c, d \in \text{Odd}(a)$ are not adjacent in $\text{Odd}(a)$ then $m_{cd} \in \{2, \infty\}$, and if $c \in \text{Odd}(a)$ and $d \in \Pi \setminus \text{Odd}(a)$ then $m_{cd} \in \{2, \infty\}$. Furthermore, any circuit in $\text{Odd}(a) \setminus \{b\}$ would similarly yield a contradiction (by combining the circuit with a minimal finite-labelled path connecting it to b). So $\text{Odd}(a) \setminus \{b\}$ is tree. Of course, $\text{Odd}(b) \setminus \{a\}$ is also a tree, by the same argument.

For each $c \in \text{Odd}(a) \setminus \{a, b\}$ let $D[a, b..c] \subseteq \Pi$ consist of b and the unique path from a to c in $\text{Odd}(a) \setminus \{b\}$. The discussion above shows that $D[a, b..c]$ is of type D . Now suppose that $c \in \text{Odd}(a) \setminus \{a, b\}$ and $e \in \Pi \setminus D[a, b..c]$ with $m_{ce} = 2$. Write $D[a, b..c] = \{c_1, c_2, \dots, c_k\}$, with $c_1 = b, c_2 = a$ and $c_k = c$, and let $d = c_1 + c_2 + c_k + 2 \sum_{i=3}^{k-1} c_i$. An argument similar to one used above shows that $r_d \in \text{FC}(c)$, and hence $W_{\{d, c, e\}}$ is finite. So $d \cdot e > -1$. If $c_i \cdot e \neq 0$ then $c_i \cdot e \leq -1/2$; so it follows that $\{i \mid c_i \cdot e \neq 0\}$ is a subset of $\{1, 2, k\}$ with at most one element. But $c_k \cdot e = 0$ since $m_{ce} = 2$, and $c_1 \cdot e = c_2 \cdot e$ since $m_{af} = m_{bf}$ for all $f \in \Pi$. So $c_i \cdot e = 0$ for all $i \in \{1, 2, \dots, k\}$. In particular, if $e \in \Pi \setminus \text{Odd}(a)$ and $m_{ce} \neq \infty$ for some $c \in \text{Odd}(a)$ then $m_{ce} = 2$, as shown above, and so $m_{de} = 2$ for all $d \in D[a, b..c]$.

All that remains to prove now is that if $c, d \in \text{Odd}(a) \setminus \{a, b\}$ with $c \notin D[a, b..d]$ and $d \notin D[a, b..c]$, then $m_{cd} = \infty$. Since c and d are not adjacent in $\text{Odd}(a)$ the only alternative is that $m_{cd} = 2$; so suppose, for a contradiction, that this holds. Choose the vertex $e \in \text{Odd}(a) \setminus \{b\}$ on the (unique) path from c to d such that the distance from e to a is minimal. Let c', d' be the neighbours of e in the path from c to d , with c' between e and c and d' between e and d . Then $c' \in C[b..c]$, and since $m_{cd} = 2$ it follows that $m_{c'd} = 2$. Now since $d' \in C[b..d]$ and $m_{c'd} = 2$ it follows that $m_{c'd'} = 2$. Thus the set $L \subseteq \Pi$ consisting of c' and d' and the vertices on the path from a to e form a system of type D_k , or A_3 if $e = a$. So L is spherical, and since $b \in \text{FC}(r_a)$ it follows that $L \cup \{b\}$ is spherical also. If $L = A_3$ then $m_{ac'} = m_{ad'} = 3$, and since $m_{bc'} = m_{ac'}$ and $m_{bd'} = m_{ad'}$ we see that $L \cup \{b\}$ is of type \tilde{A}_3 , contradicting the fact that $L \cup \{b\}$ is spherical. Similarly if L is of type D_k then $L \cup \{b\}$ is of type \tilde{D}_k , again giving a contradiction. \square

We also need to obtain further information about the situation of alternative (1) of Proposition 27. So for the next three lemmas we assume that $a \in J \subseteq L \subseteq \Pi$ with L spherical and $W_J = \text{FC}(r_a)$, and there exist $b \in J$ and $c \in L \setminus J$ such that $L_0 = \{a, c, b\}$ is a component of L of type C_3 , with $m_{ac} = 3$ and $m_{cb} = 4$.

Lemma 30. *For all $e \in \Pi \setminus \{a, c, b\}$, either $m_{ce} = m_{ae} = m_{be} = 2$ or $m_{ce} = \infty$. Moreover, $J \cap \text{Odd}(a) = \{a\}$.*

Proof. If $J \cap \text{Odd}(a) \neq \{a\}$ then, since $\{a\}$ is a component of J , Proposition 29 applies, and it follows in particular that no vertex in $\text{Odd}(a)$ lies on an edge with finite label different from 3. This contradicts $m_{bc} = 4$. So $J \cap \text{Odd}(a) = \{a\}$.

Suppose that $e \in \Pi \setminus \{a, c, b\}$ with $m_{ce} < \infty$. The group $r_c r_a W_{\{c,e\}} r_a r_c$ is finite and contains $r_c r_a W_{\{c\}} r_a r_c = W_{\{a\}}$; so there exists a maximal finite subgroup G of W containing r_a and the reflection along $(r_c r_a)e$. Since $r_b \in \text{FC}(r_a) \subseteq G$ it follows that $W_{\{b, (r_c r_a)e\}}$ is finite, and hence so is $W_{\{(r_a r_c)b, e\}} = r_a r_c W_{\{b, (r_c r_a)e\}} r_a r_c$. Hence

$$(b + \sqrt{2}c + \sqrt{2}a) \cdot e = (r_a r_c)b \cdot e > -1. \quad (4.1)$$

Assume, for a contradiction, that $m_{ce} \neq 2$. Then $c \cdot e \leq -1/2 < -1/2\sqrt{2}$, and so $(b + \sqrt{2}a) \cdot e > -1/2$, giving a contradiction if either $m_{be} \neq 2$ or $m_{ae} \neq 2$. So $b \cdot e = a \cdot e = 0$, and the inequality 4.1 above gives $c \cdot e > -1/\sqrt{2}$. So $m_{ce} = 3$. But now $W_{\{a,c,e\}}$ is of type A_3 , hence finite, and hence contained in a maximal finite subgroup that also contains $\text{FC}(r_a) = W_J$. Since $b \in J$ it follows that $\{a, c, e, b\}$ is spherical, which is false since it is of type \tilde{B}_3 . So $m_{ce} = 2$, and it remains to show that $m_{ae} = m_{be} = 2$.

Since $c \cdot e = 0$ we deduce from 4.1 that $(b + \sqrt{2}a) \cdot e > -1$, and in particular it follows that m_{ae} is 2 or 3. In either case $\{e, a, c\}$ is spherical (of type A_3 or $A_1 + A_2$), and so $\{e, a, c, b\}$ is also spherical (since $r_b \in \text{FC}(r_a)$). If either $m_{ae} \neq 2$ or $m_{be} \neq 2$

then applying Proposition 27 with $L'' = J \cup \{e, c\}$ in place of L yields a contradiction, since if L''_0 is the component of L'' containing a then $\{a, b\} \subseteq L''_0 \cap J \neq L''_0$ (since $c \notin J$), but L''_0 is not of type C_3 or D_n since it contains $\{a, c, b, e\}$. So $m_{ae} = m_{be} = 2$, as required. \square

Lemma 31. *Let $J' = J \setminus \{a\}$ and let $d \in \text{Odd}(a)$. Then $m_{cd} \neq 2$. If $m_{db'} \neq 2$ for some $b' \in J'$ then $m_{db'} = 4$, and there is a unique a' adjacent to d in $\text{Odd}(a)$; moreover, $\{a', d, b'\}$ is of type C_3 , and $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$. On the other hand, if $m_{db'} = 2$ for all $b' \in J'$ then $\text{FC}(r_d) = W_{J' \cup \{d\}}$.*

Proof. We use induction on the distance from d to a in $\text{Odd}(a)$. Observe that if $d = a$ then $m_{db'} = 2$ for all $b' \in J'$, since $\{a\}$ is a component of J , and we have $\text{FC}(r_d) = W_J = W_{J' \cup \{d\}}$ and $m_{cd} = 3 \neq 2$, as required.

Suppose now that $d \neq a$, and let $a = d_1, d_2, \dots, d_k = d$ be a minimal length path from a to d in $\text{Odd}(a)$. If $2 \leq i \leq k-1$ then d_i does not have valency 1 in $\text{Odd}(a)$, and so $m_{d_i b'} = 2$ for all $b' \in J'$, by the inductive hypothesis. The same is true for $i = 1$, since $\{a\}$ is a component of J .

We prove first that $m_{cd} \neq 2$. Assuming, for a contradiction, that $m_{cd} = 2$, then clearly $d \notin \{a, c, b\}$, and Lemma 30 tells us that $m_{ad} = 2$ and $m_{bd} = 2$. It follows that $m_{bf} = 2$ for all f in the set $M = \{d_1, d_2, \dots, d_k\}$, since $\{d_i\}$ is a component of $J' \cup \{d_i\}$ when $1 \leq i \leq k-1$. So $wb = b$ for all $w \in W_M$. Furthermore, since d and a lie in the same connected component of $\Omega(\Pi)$, we can choose $w \in W_M$ such that $wd = a$. Now since $wc \cdot a = c \cdot d = 0$ we see that the reflection r_{wc} centralizes r_a , and hence normalizes $\text{FC}(r_a) = W_J$. By Lemma 11 it follows that either $wc \in \Phi_J$ or $wc \cdot e = 0$ for all $e \in J$. But $wc \cdot b = c \cdot w^{-1}b = c \cdot b \neq 0$; so we must have $wc \in \Phi_J$, and hence $c \in w^{-1}\Phi_J \subseteq \Phi_{J \cup M}$. So $c \in M$, contradicting $m_{bf} = 2$ for all $f \in M$. So $m_{cd} \neq 2$.

Write $a' = d_{k-1}$ and $\tilde{J} = J' \cup \{a'\}$. Note that $\{a'\}$ is a component of \tilde{J} , and $\text{FC}(r_{a'}) = W_{\tilde{J}}$ (by the inductive hypothesis). Now since $\{d, a'\}$ is spherical, $\tilde{L} = \tilde{J} \cup \{d\}$ is spherical also. Let \tilde{L}_0 be the component of \tilde{L} containing a' .

Consider first the case that $m_{db'} = 2$ for all $b' \in J'$. Since also $m_{a'b'} = 2$ for all $b' \in J'$, it follows that r_d and $r_{a'}$ both fix all elements of J' . Since $v = v[d, \{a'\}] \in W_{\{a', d\}}$ satisfies $va' = d$, we conclude that

$$\text{FC}(r_d) = v \text{FC}(r_{a'}) v^{-1} = v W_{J' \cup \{a'\}} v^{-1} = W_{vJ' \cup \{va'\}} = W_{J' \cup \{d\}},$$

as required.

Now suppose that $m_{db'} \neq 2$ for some $b' \in J'$. Then $b' \in \tilde{J} \cap \tilde{L}_0$, and so $\{a'\} \neq \tilde{J} \cap \tilde{L}_0 \neq \tilde{L}_0$. Applying Proposition 27, we see that the situation of alternative (1) must hold: alternative (2) is ruled out since a' is a component of \tilde{J} , and alternative (3) is ruled out since $J' \cap \text{Odd}(a) = \emptyset$. Hence $\tilde{L}_0 = \{a', d, b'\}$ is of type C_3 , with $m_{db'} = 4$ and $m_{da'} = 3$. Furthermore, Lemma 30 tells us that $m_{de} \in \{2, \infty\}$ for all $e \in \Pi \setminus \{a', d, b'\}$; so a' is the unique neighbour of d in $\text{Odd}(a)$, as required. \square

Lemma 32. *Let $e \in \text{EOdd}(a) \setminus \text{Odd}(a)$ with $e \neq b$. Then $m_{be} = 2$.*

Proof. If $e \in J$ then it is clear that $m_{be} = 2$, since b is a component of J . So we may assume that $e \notin J$.

As above, write $J' = J \setminus \{a\}$. Since $e \in \text{EOdd}(a)$ there exists a $d \in \text{Odd}(a)$ with m_{de} even. If d is adjacent to some $b' \in J'$ then, by Lemma 31, there is a unique $a' \in \text{Odd}(a)$ adjacent to d ; furthermore, $\{a', d, b'\}$ is of type C_3 , and $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$. By Lemma 30, since $m_{de} \neq \infty$ it follows that $m_{a'e} = m_{de} = 2$. On the other hand, if d is not adjacent to any element of J' then Lemma 31 tells us that $\text{FC}(r_d) = W_{J' \cup \{d\}}$. So in either case there is an $a' \in \text{Odd}(a)$ with $m_{a'e}$ even and $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$.

Choose such an a' . Since $m_{a'e}$ is even, $v[e, \{a'\}]a' = a'$; moreover $v[e, \{a'\}]$ is the reflection along some root $f = \lambda e + \mu a'$. Note that $f \cdot a' = 0$, and hence $\lambda \neq 0$. Since $e \notin J$ it follows that $f \notin \Phi_J$. But r_f centralizes $r_{a'}$, and hence normalizes $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$. By Lemma 11 it follows that $f \cdot b = 0$. But also $a' \cdot b = 0$, since $\{a'\}$ and $\{b\}$ are distinct components of $J' \cup \{a'\}$; so it follows that $e \cdot b = 0$. Thus $m_{be} = 2$, as required. \square

Lemmas 30, 31 and 32 combine to yield the following result.

Proposition 33. *Suppose that $a \in J \subseteq L \subseteq \Pi$, with L spherical and $\text{FC}(r_a) = W_J$, and let L_0 be the component of L containing a . Suppose that $L_0 = \{a, c, b\}$ is of type C_3 , with $m_{ac} = 3$ and $m_{bc} = 4$, and $J \cap L_0 = \{a, b\}$. Then b is a C_3 -neighbour of $\text{Odd}(a)$. Furthermore, $J \cap \text{Odd}(a) = \{a\}$, and if $a' \in \text{Odd}(a)$ is not adjacent to any C_3 -neighbour of $\text{Odd}(a)$ then $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$, where $J' = J \setminus \{a\}$.*

Proof. If $m_{bd} \neq 2$ for some $d \in \text{Odd}(a)$, then $m_{db} = 4$, by Lemma 31. There is at least one $d \in \text{Odd}(a)$ such that $m_{bd} = 4$, namely $d = c$. Lemma 31 tells us that for each $d \in \text{Odd}(a)$ with $m_{bd} = 4$ there is an $a' \in \text{Odd}(a)$ such that $\{a', d, b\}$ is a system of type C_3 . Moreover, by Lemma 30, if $e \in \Pi \setminus (\text{Odd}(a) \cup \{b\})$ then either $m_{de} = \infty$ or $m_{ae} = m_{be} = m_{de} = 2$, while if $e \in \text{Odd}(a) \setminus \{a, c\}$ then $m_{de} = \infty$, since $m_{de} \neq 2$ by Lemma 31. And if $e \in \text{EOdd}(a) \setminus (\text{Odd}(a) \cup \{b\})$ then $m_{be} = 2$, by Lemma 32. So b satisfies all the requirements of a C_3 -neighbour of $M = \text{Odd}(a)$, as specified in Definition 4.

It now follows from Lemma 31 that if $a' \in \text{Odd}(a)$ is adjacent to some $b' \in J'$ then b' is a C_3 -neighbour of $\text{Odd}(a)$, and if a' is not adjacent to any such b' then $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$. Finally, $J \cap \text{Odd}(a) = \{a\}$, by Lemma 30. \square

Let $a, a' \in \Pi$, and suppose that $w \in W$ has the property that $wa = a'$. By Proposition 12 there exist $a_i \in \text{Odd}(a)$ and $c_i \in \Pi$ such that

- (i) $a_1 = a$ and $a_{k+1} = a'$,
- (ii) $m_{c_i a_i} \neq \infty$ and $v[c_i, \{a_i\}]a_i = a_{i+1}$, for all $i \in \{1, 2, \dots, k\}$,
- (iii) $w = v[c_k, \{a_k\}] \dots v[c_2, \{a_2\}]v[c_1, \{a_1\}]$.

Now let b be a C_3 -neighbour of $\text{Odd}(a)$. For each $c \in \text{Odd}(a)$ that is adjacent to b , define $X(c) = b + \sqrt{2}c + \sqrt{2}\tilde{a}$, where \tilde{a} is the unique neighbour of c in $\text{Odd}(a)$, and for each $c \in \text{Odd}(a)$ that is not adjacent to b , define $X(c) = b$. We show that $v[c_i, \{a_i\}]X(a_i) = X(a_{i+1})$ for all $i \in \{1, 2, \dots, k\}$.

Suppose first that neither c_i nor a_i is adjacent to b . Then $X(a_i) = b$, and since $a_{i+1} \in \{a_i, c_i\}$ we have that $X(a_{i+1}) = b$ also. Since r_{a_i} and r_{c_i} both fix b , and $v[c_i, \{a_i\}] \in W_{\{a_i, c_i\}}$, it follows that

$$v[c_i, \{a_i\}]X(a_i) = v[c_i, \{a_i\}]b = b = X(a_{i+1}),$$

as required.

Next, suppose that c_i is adjacent to b , but a_i is not adjacent to b . Since $m_{c_i a_i} \neq \infty$ and $a \in \text{Odd}(a)$ it follows that $c_i \in \text{EOdd}(a)$. Since b is a C_3 -neighbour of $\text{Odd}(a)$, it is not adjacent to any element of $\text{EOdd}(a) \setminus \text{Odd}(a)$; so $c_i \in \text{Odd}(a)$, and, moreover, $m_{c_i e} = \infty$ for all $e \in \text{Odd}(a) \setminus \{c_i\}$ apart from the unique neighbour of c_i in $\text{Odd}(a)$. So a_i is this unique neighbour, $m_{c_i a_i} = 3$, and $a_{i+1} = v[c_i, \{a_i\}]a_i = c_i$. Moreover, $m_{c_i b} = 4$ and $m_{a_i b} = 2$. So

$$v[c_i, \{a_i\}]X(a_i) = r_{a_i} r_{c_i} b = b + \sqrt{2}c_i + \sqrt{2}a_i = X(c_i) = X(a_{i+1})$$

as required.

Now suppose that a_i is adjacent to b , and let \tilde{a} be the unique neighbour of a_i in $\text{Odd}(a)$. Since $m_{a_i e} = \infty$ for all $e \in \text{Odd}(a) \setminus \{a_i, \tilde{a}\}$, if $c_i \in \text{Odd}(a)$ then $c_i = \tilde{a}$. In this case we see that

$$v[c_i, \{a_i\}]X(a_i) = r_{a_i} r_{c_i} (b + \sqrt{2}c_i + \sqrt{2}a_i) = b = X(c_i) = X(a_{i+1}),$$

since $a_{i+1} = v[c_i, \{a_i\}]a_i = c_i$. If $c_i = b$ then $v[c_i, \{a_i\}] = r_b r_{a_i} r_b$, which fixes both a_i and $X(a_i) = b + \sqrt{2}a_i + \sqrt{2}\tilde{a}$. So $v[c_i, \{a_i\}]X(a_i) = X(a_{i+1})$ in this case too. Finally, suppose that $c_i \notin \text{Odd}(a) \cup \{b\}$. Since $m_{c_i a_i} \neq \infty$ we must have $m_{c_i \tilde{a}} = m_{c_i a_i} = m_{c_i b} = 2$, (by the definition of a C_3 -neighbour). So

$$v[c_i, \{a_i\}]X(a_i) = r_{c_i} (b + \sqrt{2}c_i + \sqrt{2}\tilde{a}) = b + \sqrt{2}c_i + \sqrt{2}\tilde{a} = X(a_{i+1})$$

since $a_{i+1} = r_{c_i} a_i = a_i$.

We have now covered all cases, and shown that $v[c_i, \{a_i\}]X(a_i) = X(a_{i+1})$ for all $i \in \{1, 2, \dots, k\}$. By a trivial induction it follows that $X(a_{k+1}) = wX(a_1)$.

Thus we have established the following result.

Lemma 34. *Let $a \in \Pi$ and $w \in W$ such that $wa \in \Pi$. Suppose that b is a C_3 -neighbour of $\text{Odd}(a)$ that is not adjacent to a . Then*

$$wb = \begin{cases} b & \text{if } wa \text{ is not adjacent to } b, \\ b + \sqrt{2}wa + \sqrt{2}\tilde{a} & \text{if } wa \text{ is adjacent to } b, \end{cases}$$

where \tilde{a} is adjacent to wa in $\text{Odd}(a)$.

We are now able to give a detailed description of the components of J whenever W_J is the finite continuation of a simple reflection.

Proposition 35. *Suppose that $a \in J \subseteq \Pi$ with $W_J = \text{FC}(r_a)$, and suppose that K is a component of J . Then one of the following alternatives holds.*

- (a) $K = \{a\} = J \cap \text{Odd}(a)$.
- (b) $K = \{a, b\}$ is of type C_2 , and $J \cap \text{Odd}(a) = \{a\}$.
- (c) $K = \{a\}$ or $K = \{b\}$, where $\{a, b\} = J \cap \text{Odd}(a)$ is of type $A_1 + A_1$.
- (d) $K = \{b\} \not\subseteq \text{Odd}(a)$, and b is a C_3 -neighbour of $\text{Odd}(a)$.
- (e) $\text{Odd}(a) \subseteq K$, and K is a component of $\text{EOdd}(a)$.
- (f) $K \cap \text{Odd}(a) = \emptyset$, and K is a component of $\text{EOdd}(a)$.

Proof. We consider first the case that $K \cap \text{Odd}(a) \neq \emptyset$, and start by supposing that there exists a spherical $L \subseteq \Pi$ with $J \subseteq L$ and K not a component of L .

Choose such an L , and let L_0 be the component of L containing a . By Proposition 27, since K is not a component of L we must have $K \subseteq L_0$. So either $K = \{a\}$, in which case (a) above holds, or else $\{a\} \subsetneq \{a\} \cup K \subseteq J \cap L_0$. Furthermore, $J \cap L_0 \neq L_0$, since $K \neq L_0$. So if (a) does not hold then $\{a\} \neq J \cap L_0 \neq L_0$, and so one of the alternatives (1), (2) or (3) of Proposition 27 must hold.

Suppose that alternative (2) holds, so that $K = \{a, b\} = J \cap L_0$ for some b , and $\{a, b\}$ is of type C_2 . By Proposition 28 we see that each $c \in \text{Odd}(a) \setminus \{a\}$ lies in a type C spherical subset L' of Π containing $\{a, b\}$. Since $J \cap L' = \{a, b\}$ (by Proposition 27) it follows that $c \notin J$. So $J \cap \text{Odd}(a) = \{a\}$, and (b) above is satisfied.

Suppose that alternative (3) of Proposition 27 holds, so that $J \cap L_0 = \{a, b\}$ is of type $A_1 + A_1$, and $b \in \text{Odd}(a)$. Proposition 29 immediately yields that $J \cap \text{Odd}(a) = \{a, b\}$, and so (c) above is satisfied.

Suppose that alternative (1) of Proposition 27 holds, so that $L_0 = \{a, c, b\}$ with $m_{ac} = 3$ and $m_{cb} = 4$, and $J \cap L_0 = \{a, b\}$. By Lemma 30 we know that $b \notin \text{Odd}(a)$, and since we have assumed that $K \cap \text{Odd}(a) \neq \emptyset$, it follows that $K = \{a\} = J \cap \text{Odd}(a)$. So (a) holds.

We have now dealt with all cases that arise if there is a spherical $L \subseteq \Pi$ with $J \subseteq L$ and K not a component of L . So assume that K is a component of every spherical L containing J . We show that in this case $\text{Odd}(a) \subseteq K$, and K is a component of $\text{EOdd}(a)$; that is, (e) above holds.

To show that $\text{Odd}(a) \subseteq K$ it is clearly sufficient to show that if $a' \in K \cap \text{Odd}(a)$ and b is adjacent to a' in $\text{Odd}(a)$ then $b \in K$. Note that since $a' \in \text{Odd}(a)$ there exists $w \in W$ with $a' = wa$, and since Proposition 25 yields that $w \in N_W(W_J)$ it follows that $\text{FC}(r_{a'}) = W_J$. Now the assumption that b and a' are adjacent in $\text{Odd}(a)$ implies that $\{a', b\}$ is spherical, and therefore $J \cup \{b\}$ is spherical. But K is a component of

every spherical subset of Π containing J ; so it is a component of $J \cup \{b\}$. But $a' \in K$ and b is adjacent to a' ; so $b \in K$, as required.

Since $K \subseteq J \subseteq \text{EOdd}(a)$ and K is connected, saying that K is a component of $\text{EOdd}(a)$ is equivalent to saying that $m_{bc} = 2$ whenever $b \in K$ and $c \in \text{EOdd}(a) \setminus K$. So suppose that $c \in \text{EOdd}(a) \setminus K$. Then there exists an $a' \in \text{Odd}(a)$ such that $m_{a'c}$ is even. Thus $\{a', c\}$ is spherical, and as above it follows that $J \cup \{c\}$ is spherical. So K must be a component of $J \cup \{c\}$, and since $c \notin K$ it follows that $m_{bc} = 2$ for all $b \in K$, as required.

It remains to consider the case that $K \cap \text{Odd}(a) = \emptyset$; we must show that either (f) or (d) holds. We start by supposing that there exists a spherical $L \subseteq \Pi$ and a $w \in W$ with $wJ \subseteq L$ and wK not a component of L .

Choose such L and w , and let L_0 be the component of L containing wa . By Proposition 27, since wK is not a component of L we must have $wK \subseteq L_0$. Now $wJ \cap L_0 \neq L_0$ since $wK \neq L_0$, and $\{wa\} \neq wJ \cap L_0$ since $wa \notin wK$. So one of the alternatives (1), (2) or (3) of Proposition 27 must hold. Alternative (3) can be ruled out, since in that case $wJ \cap L_0 \subseteq \text{Odd}(wa)$, which is impossible since $K \cap \text{Odd}(a) = \emptyset$. If alternative (2) holds then $wK = wJ \cap L_0$ contains wa and is of type C_2 , whence K contains a and is of type C_2 , and (b) is satisfied. If alternative (1) holds then since $wK \neq \{wa\}$ it follows from Proposition 32 that $wK = \{b\}$, with b a C_3 -neighbour of $\text{Odd}(a)$. Since wa is not adjacent to b , it follows from Lemma 34 that $w^{-1}b = b$, unless a is adjacent to b , in which case $w^{-1}b = b + \sqrt{2}a + \sqrt{2}\tilde{a}$ for some \tilde{a} in $\text{Odd}(a)$. But this latter case cannot occur, since $w^{-1}b \in K \subseteq \Pi$. So $K = wK = \{b\}$, with b a C_3 -neighbour of $\text{Odd}(a)$, and (d) holds.

Finally, suppose that wK is a component of every spherical $L \subseteq \Pi$ such that $wJ \subseteq L$ for some $w \in W$. For each $c \in \text{EOdd}(a) \setminus K$ there is then a sequence $a = a_0, a_1, \dots, a_k = c$ in Π such that $m_{a_{i-1}a_i}$ finite for all $i \in \{1, 2, \dots, k\}$ and odd for all $i \in \{1, 2, \dots, k-1\}$. We shall show that, for every such sequence, $m_{ba_i} = 2$ for all $b \in K$ and $i \in \{0, 1, \dots, k\}$; in particular, this will show that $m_{bc} = 2$ whenever $b \in K$ and $c \in \text{EOdd}(a) \setminus K$, enabling us to conclude that K is a component of $\text{EOdd}(a)$.

The case $k = 0$ is clear, since $a \in J \setminus K$ and K is a component of J . Proceeding by induction, we may assume that $k > 0$ and $m_{ba_i} = 2$ for all $i \in \{1, 2, \dots, k-1\}$ and all $b \in K$. We see that the element $u = v[a_{k-1}, \{a_{k-2}\}]v[a_{k-2}, \{a_{k-3}\}] \dots v[a_1, \{a_0\}]$ centralizes W_K and has the property that $ua = a_{k-1}$, since the labels in the path from a to a_{k-1} are all odd. The group $u^{-1}W_{\{a_{k-1}, a_k\}}u$ is finite and contains $u^{-1}r_{a_{k-1}}u = r_a$, and so there is a maximal finite subgroup G of W containing this group and also containing W_J .

Note that $W_J \cup \{u^{-1}r_c u\} \subseteq G = w^{-1}W_L w$, for some $w \in W$ and spherical $L \subseteq \Pi$, the element u being in the centralizer of W_K . We may choose w to be the minimal length element of $W_L w = W_L w W_J$, and it then follows from Lemma 15 that $wJ \subseteq L$. Hence wK is a component of L . Furthermore, since $wu^{-1}r_c u w^{-1} \in W_L$

we see that the root $wu^{-1}c$ is in Φ_L and not in $\Phi_{wK} = wu^{-1}\Phi_K$ (since $c \notin \Phi_K$). So $wu^{-1}c \cdot wu^{-1}b = 0$ for all $b \in K$. So $c \cdot b = 0$, or (equivalently) $m_{bc} = 2$ for all $b \in K$, as required. \square

To complement the results we have obtained so far, our next task is to find conditions that ensure that a visible subgroup W_K is contained in $\text{FC}(r_a)$.

Lemma 36. *Let $a \in \Pi$ and K a component of $\text{EOdd}(a)$ such that W_K is finite. Then $W_K \subseteq \text{FC}(r_a)$.*

Proof. Let F be a maximal finite subgroup of W with $r_a \in F$, and choose $w \in W$ such that $wFw^{-1} = W_L$ for some $L \subseteq \Pi$. We may replace w by the minimal length element in the double coset $W_L w W_{\{a\}}$, since this does not affect the condition $wFw^{-1} = W_L$. So we have that $w^{-1}L \subseteq \Phi^+$, and, moreover, $r_a \in w^{-1}Lw \cap W_{\{a\}} = W_{w^{-1}L \cap \{a\}}$ by Lemma 15. So $wa \in L \subseteq \Pi$, and by Lemma 12 we see that w is a product of factors of the form $v[d, \{c\}]$, with $c, d \in \text{EOdd}(a)$. Since K is a component of $\text{EOdd}(a)$ it follows that each $v[d, \{c\}]$ normalizes W_K , and therefore w normalizes W_K . Moreover, since $wa \in L$ and L is spherical, it follows that $L \subseteq \text{EOdd}(wa) = \text{EOdd}(a)$. So W_L normalizes W_K . But W_K is finite, by hypothesis, and W_L is a maximal finite subgroup of W . So $W_K \subseteq W_L$, and $W_K = w^{-1}W_Kw \subseteq w^{-1}W_Lw = F$. Thus W_K is contained in all maximal finite subgroups of W containing r_a , as required. \square

Lemma 37. *Let $a \in \Pi$ and let b be a C_3 -neighbour of $\text{Odd}(a)$. If a and b are not adjacent in Π then $r_b \in \text{FC}(r_a)$.*

Proof. Let F be a maximal finite subgroup of W with $r_a \in F$. As in the proof of Lemma 36 there exist a $w \in W$ and a maximal spherical $L \subseteq \Pi$ with $wa = a' \in L$ and $F = w^{-1}W_Lw$. Since L is spherical, $L \subseteq \text{EOdd}(a)$.

Suppose first that a' is not adjacent to b . Then $m_{ca'} = \infty$ for every $c \in \text{Odd}(a)$ that is adjacent to b , and since $a' \in L$ it follows that no such c is in L . Thus $m_{be} = 2$ for all $e \in L \cap \text{Odd}(a)$. But since also $m_{be} = 2$ for all $e \in \text{EOdd}(a) \setminus (\text{Odd}(a) \cup \{b\})$, it follows that $m_{be} = 2$ for all $e \in L \setminus \{b\}$. Thus $\{b\}$ is a component of $L \cup \{b\}$, and since L is spherical it follows that $L \cup \{b\}$ is spherical. Maximality of L tells us that $b \in L$. Moreover, Lemma 34 gives $wb = b$, and so $r_b = w^{-1}r_bw \in w^{-1}W_Lw = F$.

On the other hand, suppose that a' is adjacent to b . In this case Lemma 34 gives $wb = b + \sqrt{2}a' + \sqrt{2}\tilde{a}$, where \tilde{a} is the unique neighbour of a' in $\text{Odd}(a)$. Furthermore, since $m_{a'e} \in \{2, \infty\}$ for all $e \in \Pi \setminus \{\tilde{a}, a', b\}$, we see that $m_{a'e} = 2$ for all $e \in L \setminus \{\tilde{a}, a', b\}$ (since L is spherical). But the definition of a C_3 -vertex also requires that $m_{\tilde{a}e} = m_{be} = 2$ whenever $m_{a'e} = 2$; so it follows that $\{\tilde{a}, a', b\}$ is a component of $L \cup \{\tilde{a}, a', b\}$, which is therefore spherical since L and $\{\tilde{a}, a', b\}$ are both spherical. Maximality of L tells us that $\{\tilde{a}, a', b\} \subseteq L$; so $wb = b + \sqrt{2}a' + \sqrt{2}\tilde{a} \in \Phi_L$, and $r_b = w^{-1}r_bw \in w^{-1}W_Lw = F$.

So $r_b \in F$ in all cases, and so r_b is contained in all maximal finite subgroups of W containing r_a , as required. \square

We now prove the converse to Proposition 28.

Proposition 38. *Let $a \in \Pi$ and $b \in \Pi \setminus \text{Odd}(a)$, and suppose that (a, b) is a focus of $\text{Odd}(a)$ in Π . Then $\text{FC}(r_a) = W_J$, where J is the union of $\{a, b\}$ and the spherical components of $\text{EOdd}(a)$. Moreover, $\text{FC}(r_{a'})$ is not visible for any $a' \in \text{Odd}(a) \setminus \{a\}$.*

Proof. For each $c \in \text{Odd}(a)$ let $X(c) = b + \sqrt{2} \sum_{i=1}^m c_i$ and $Y(c) = b + \sqrt{2} \sum_{i=1}^{m-1} c_i$, where $c_1 = a, c_2, \dots, c_m = c$ is the unique path from a to c in $\text{Odd}(a)$, noting that $X(c)$ and $Y(c)$ are roots in $\Phi_{C[b..c]}$. We remark, for later use, that $X(c)$ and $Y(c)$ are fixed by the reflections $r_b, r_{c_1}, \dots, r_{c_{m-2}}$.

Let $F = w^{-1}W_L w$ be a maximal finite subgroup of W containing r_a , with $L \subseteq \Pi$ and w of minimal length in $W_L w W_a$. Then $wa = a' \in L$, by Lemma 15. Put $L_0 = L \cap \text{Odd}(a)$.

Choose $c \in L_0$ with $C[b..c]$ of maximal cardinality. If $d \in L_0$ then $m_{cd} \neq \infty$ (since L_0 is spherical), whence $d \in C[b..c]$ by condition (3) of Definition 5. So $L_0 \subseteq C[b..c]$. Now if $e \in L \setminus L_0$ is arbitrary then $e \notin \text{Odd}(a)$ (since $e \notin L \cap \text{Odd}(a)$) and $m_{ce} < \infty$ (since $c, e \in L$ and L is spherical). By condition (4) of Definition 5 it follows that $m_{de} = 2$ for all $d \in C[b..c]$. Since this holds for all $e \in L \setminus L_0$, and $C[b..c]$ and $L \setminus L_0$ are both spherical, it follows that $C[b..c] \cup (L \setminus L_0)$ is spherical. But this set contains L (since $L_0 \subseteq C[b..c]$) and since L is a maximal spherical subset of Π we conclude that $L = C[b..c] \cup (L \setminus L_0)$.

By Proposition 12 and Lemma 9 there exist simple roots e_1, e_2, \dots, e_k and $d_1 = a, d_2, \dots, d_{k+1} = a' \in \text{Odd}(a)$ with $w = v[e_k, \{d_k\}] \dots v[e_2, \{d_2\}] v[e_1, \{d_1\}]$ and $v[e_i, \{d_i\}]d_i = d_{i+1}$ for all $i \in \{1, 2, \dots, k\}$. Moreover, $m_{e_i d_i} < \infty$ for all i . Let $w_0 = 1$ and $w_i = v[e_i, \{d_i\}]w_{i-1}$; we will show that

$$\begin{aligned} &\{w_i b, -w_i b, w_i(b + \sqrt{2}a), -w_i(b + \sqrt{2}a)\} \\ &= \{X(d_{i+1}), -X(d_{i+1}), Y(d_{i+1}), -Y(d_{i+1})\} \end{aligned}$$

for all $i \in \{0, 1, \dots, k\}$. The case $i = 0$ is trivial.

Proceeding inductively, suppose that $i > 1$ and

$$\{\pm w_{i-1} b, \pm w_{i-1}(b + \sqrt{2}a)\} = \{\pm X(d_i), \pm Y(d_i)\}.$$

It will be sufficient to show that $v[e_i, \{d_i\}]X(d_i)$ and $v[e_i, \{d_i\}]Y(d_i)$ both lie in the set $\{\pm X(d_{i+1}), \pm Y(d_{i+1})\}$.

Suppose first that $d_i = a$. Then $X(d_i) = b + \sqrt{2}a$ and $Y(d_i) = b$. If $e_i \notin \text{Odd}(a)$ then $m_{e_i d_i}$ is even, and $d_{i+1} = d_i = a$. Furthermore, by condition (4) of Definition 5 we have either $\{e_i, d_i\} = \{b, a\}$ or $m_{e_i b} = m_{e_i a} = 2$. In the former case

$v[e_i, \{d_i\}] = v[b, \{a\}] = r_b r_a r_b$, giving $v[e_i, \{d_i\}]b = -b - \sqrt{2}a = -X(a)$ and $v[e_i, \{d_i\}](b + \sqrt{2}a) = -b = -Y(a)$; in the latter case $v[e_i, \{d_i\}] = v[e_i, \{a\}] = r_{e_i}$, giving $v[e_i, \{d_i\}]b = b = Y(a)$ and $v[e_i, \{d_i\}](b + \sqrt{2}a) = b + \sqrt{2}a = X(a)$. If $e_i \in \text{Odd}(a)$ then $a \in C[b..e_i]$, and by condition (2) of Definition 5 we have $m_{e_i b} = 2$ and either $m_{e_i a} = 2$ or $m_{e_i a} = 3$. If $m_{e_i a} = 2$ then $d_{i+1} = d_i = a$, while if $m_{e_i a} = 3$ then $d_{i+1} = e_i$. Furthermore, in former case we find that $v[e_i, \{d_i\}]b = r_{e_i} b = b = Y(a)$ and $v[e_i, \{d_i\}](b + \sqrt{2}a) = b + \sqrt{2}a = X(a)$, while in the latter case we find that $v[e_i, \{d_i\}]b = r_a r_{e_i} b = b + \sqrt{2}a = Y(e_i)$ and $v[e_i, \{d_i\}](b + \sqrt{2}a + \sqrt{2}e_i) = X(e_i)$.

Now suppose that $d_i \neq a$. If $e_i \notin \text{Odd}(a) \cup \{b\}$ then $m_{e_i d_i} = 2$ and $d_{i+1} = d_i$. Moreover, $m_{e_i d} = 2$ for all $d \in C[b..d_i]$, and so $v[e_i, \{d_i\}] = r_{e_i}$ fixes all the roots in $\Phi_{C[b..d_i]}$, including $X(d_i) = X(d_{i+1})$ and $Y(d_i) = Y(d_{i+1})$. If $e_i \in \text{Odd}(a) \cup \{b\}$ and $\{e_i, d_i\}$ is not an edge of $\text{Odd}(a)$ then we again have $d_{i+1} = d_i$ and $v[e_i, \{d_i\}] = r_{e_i}$. By condition (3) of Definition 5 we either have $d_i \in C[b..e_i]$ or $e_i \in C[b..d_i]$. In the former case we have $m_{e_i d} = 2$ for all $d \in C[b..d_i]$, and as above we see that r_{e_i} fixes $X(d_i)$ and $Y(d_i)$. In the latter case the remark made at the start of the proof implies that it is still true that r_{e_i} fixes $X(d_i)$ and $Y(d_i)$. So we have shown that when $m_{e_i d_i} = 2$ it is true that $v[e_i, \{d_i\}]X(d_i)$ and $v[e_i, \{d_i\}]Y(d_i)$ both lie in the set $\{\pm X(d_{i+1}), \pm Y(d_{i+1})\}$, and it remains only to consider the case that e_i and d_i are adjacent in $\text{Odd}(a)$. Note that in this case $d_{i+1} = e_i$.

Let $C[b..d_i] = \{b, c_1, \dots, c_m\}$ with $c_1 = a$ and $c_m = d_i$, and suppose that $e_i = c_{m-1}$ is the vertex adjacent to d_i in $C[b..d_i]$. Then

$$\begin{aligned} v[e_i, \{d_i\}]X(d_i) &= r_{c_m} r_{c_{m-1}} \left(b + \sqrt{2} \sum_{j=1}^m c_j \right) \\ &= r_{c_m} \left(b + \sqrt{2} \sum_{j=1}^m c_j \right) = b + \sqrt{2} \sum_{j=1}^{m-1} c_j = X(e_i), \end{aligned}$$

and similarly

$$\begin{aligned} v[e_i, \{d_i\}]Y(d_i) &= r_{c_m} r_{c_{m-1}} \left(b + \sqrt{2} \sum_{j=1}^{m-1} c_j \right) \\ &= r_{c_m} \left(b + \sqrt{2} \sum_{j=1}^{m-2} c_j \right) = b + \sqrt{2} \sum_{j=1}^{m-2} c_j = Y(e_i). \end{aligned}$$

The alternative possibility is that d_i is adjacent to e_i in $C[b..e_i]$. Exactly the same calculations show that $v[e_i, \{d_i\}]X(d_i) = X(e_i)$ and $v[e_i, \{d_i\}]Y(d_i) = Y(e_i)$ in this case also.

The induction is now complete, and it follows in particular that $wb = w_kb$ is one of $\pm X(a')$ or $\pm Y(a')$. Hence

$$wb \in \Phi_{C[b..a']} \subseteq \Phi_{C[b..c]} \subseteq \Phi_L.$$

Thus $wr_bw^{-1} \in W_L$, and so $r_b \in w^{-1}W_Lw = F$. Since F was an arbitrary maximal finite subgroup of W containing r_a , this shows that $r_b \in \text{FC}(r_a)$.

Let \tilde{M} be the component of $\text{EOdd}(a)$ containing $\text{Odd}(a)$, and suppose, for a contradiction, that \tilde{M} is spherical. Clearly $b \in \tilde{M}$, as $m_{ba} = 4$, but $\tilde{M} = \text{Odd}(a) \cup \{b\}$ is not permitted, in view of condition (5) of Definition 5. So $\tilde{M} \setminus (\text{Odd}(a) \cup \{b\}) \neq \emptyset$. But for $e \in \tilde{M} \setminus (\text{Odd}(a) \cup \{b\})$ and $c \in \text{Odd}(a)$ we have $m_{ce} \neq \infty$, since \tilde{M} is spherical, and by condition (4) of Definition 5 it follows that $m_{be} = m_{ce} = 2$ for all $c \in \text{Odd}(a)$. This contradicts the fact that \tilde{M} is connected.

Now suppose that $a' \in \text{Odd}(a)$ is such that $\text{FC}(r_{a'}) = W_J$ for some $J \subseteq \Pi$, and let J_0 be the component of J containing a' . Since $J_0 \neq \tilde{M}$ it follows from Proposition 35 that J_0 has rank at most 2. Now since there exists $w \in W_{C[b..a']}$ such that $wa = -a'$ and $wb = X(a')$, and since $r_b \in \text{FC}(r_a)$, it follows that $r_{wb} = wr_bw^{-1} \in \text{FC}(wr_ar_a^{-1}) = \text{FC}(r_{a'})$. Thus $X(a') \in \Phi_J$, and so $C[b..a'] \subseteq J$. Since J_0 has rank at most 2, this means that $a' = a$ and $J_0 = \{a, b\}$.

It remains to prove that J is the union of J_0 and the spherical components of $\text{EOdd}(a)$. By Lemma 36 we know that all these components are contained in J . But if K is any other component of J such that $K \cap \text{Odd}(a) = \emptyset$, then by Proposition 35 we see that $K = \{b'\}$, with b' a C_3 -neighbour of $\text{Odd}(a)$. Since b is the only element of Π such that $m_{bc} \in \{2, 4\}$ for all $c \in \text{Odd}(a)$, we must have $b' = b$, contradicting the fact that the component of J containing b is $J_0 = \{a, b\}$. \square

Next, we have the converse to Proposition 29.

Proposition 39. *Let $a \in \Pi$ and suppose that there exists a $b \in \text{Odd}(a)$ such that $\{a, b\}$ is a half-focus of $\text{Odd}(a)$ in Π . Suppose also that the vertices $\text{Odd}(a)$ do not comprise a spherical subset of Π . Then $\text{FC}(r_a) = W_J$, where J is the union of $\{a, b\}$ and the spherical components of $\text{EOdd}(a)$. Moreover, $\text{FC}(r_{a'})$ is not visible for any $a' \in \text{Odd}(a) \setminus \{a, b\}$.*

Proof. For each $c \in \text{Odd}(a) \setminus \{a, b\}$, define

$$X(c) = b + a + c + 2 \sum_{i=2}^{m-1} c_i$$

where $c_1 = a$, $c_2, \dots, c_m = c$ is the unique path from a to c in $\text{Odd}(a) \setminus \{b\}$. Then $X(c)$ is a root in $\Phi_{D[a,b..c]}$ and is fixed by the reflections $r_b, r_{c_1}, \dots, r_{c_{m-2}}$ and r_{c_m} . Define also $X(a) = b$ and $X(b) = a$.

Let $F = w^{-1}W_L w$ be a maximal finite subgroup of W containing r_a , with $L \subseteq \Pi$ and w of minimal length in $W_L w W_a$. Then $wa = a' \in L$, by Lemma 15. Put $L_0 = L \cap \text{Odd}(a)$.

Choose $c \in L_0$ with $D[a, b..c]$ of maximal cardinality. If $d \in L_0$ then $m_{cd} \neq \infty$ (since L_0 is spherical), whence $d \in D[a, b..c]$ by condition (4) of Definition 6. So $L_0 \subseteq D[a, b..c]$. Now if $e \in L \setminus L_0$ is arbitrary then $e \notin \text{Odd}(a)$ (since $e \notin L \cap \text{Odd}(a)$) and $m_{ce} < \infty$ (since $c, e \in L$ and L is spherical). By condition (5) of Definition 6 it follows that $m_{de} = 2$ for all $d \in D[a, b..c]$. Since this holds for all $e \in L \setminus L_0$, and $D[a, b..c]$ and $L \setminus L_0$ are both spherical, it follows that $D[a, b..c] \cup (L \setminus L_0)$ is spherical. But this set contains L (since $L_0 \subseteq D[a, b..c]$) and since L is a maximal spherical subset of Π we conclude that $L = D[a, b..c] \cup (L \setminus L_0)$.

By Proposition 12 and Lemma 9 there exist simple roots e_1, e_2, \dots, e_k and $d_1 = a, d_2, \dots, d_{k+1} = a' \in \text{Odd}(a)$ with $w = v[e_k, \{d_k\}] \dots v[e_2, \{d_2\}]v[e_1, \{d_1\}]$ and $v[e_i, \{d_i\}]d_i = d_{i+1}$ for all $i \in \{1, 2, \dots, k\}$. Furthermore, we have $m_{e_i d_i} < \infty$ for all i . Let $w_0 = 1$, and $w_i = v[e_i, \{d_i\}]w_{i-1}$ for $i \geq 1$. We will show that

$$\{w_i b, -w_i b\} = \{X(d_{i+1}), -X(d_{i+1})\}$$

for all $i \in \{0, 1, \dots, k\}$.

The case $i = 0$ is trivial. Proceeding inductively, suppose that $i > 1$ and $w_{i-1}b = \pm X(d_i)$. It will be sufficient to show that $v[e_i, \{d_i\}]X(d_i) = \pm X(d_{i+1})$.

Suppose first that $d_i = a$, so that $X(d_i) = b$. If $e_i \neq b$ then $m_{e_i b} = m_{e_i a} \in \{2, 3\}$, since $m_{e_i a} = m_{e_i d_i} \neq \infty$. We also have $m_{e_i a} = 2$ if $e_i = b$. In the case $m_{e_i a} = 3$ we have $v[e_i, \{d_i\}] = r_a r_{e_i}$, and $d_{i+1} = r_a r_{e_i} a = e_i$. Furthermore,

$$v[e_i, \{d_i\}]X(d_i) = r_a r_{e_i} b = a + b + e_i = X(e_i) = X(d_{i+1}),$$

as required. In the case $m_{e_i a} = 2$ we have $v[e_i, \{d_i\}] = r_{e_i}$, giving $d_{i+1} = r_{e_i} a = a$, and

$$v[e_i, \{d_i\}]X(d_i) = r_{e_i} b = \pm b = \pm X(d_{i+1}),$$

since either $e_i = b$ or $m_{e_i b} = 2$.

The case $d_i = b$ is the same as the case $d_i = a$ with a and b interchanged; so suppose that $d_i \notin \{a, b\}$. If $e_i \notin \text{Odd}(a)$ then $m_{e_i d_i} = 2$ and $d_{i+1} = d_i$. Moreover, $m_{e_i d} = 2$ for all $d \in D[a, b..d_i]$, and so $v[e_i, \{d_i\}] = r_{e_i}$ fixes all the roots in $\Phi_{D[a, b..d_i]}$, including $X(d_i) = X(d_{i+1})$. If $e_i \in \text{Odd}(a)$ and $\{e_i, d_i\}$ is not an edge of $\text{Odd}(a)$ then we again have $d_{i+1} = d_i$ and $v[e_i, \{d_i\}] = r_{e_i}$. By condition (4) of Definition 6 we either have $d_i \in D[a, b..e_i]$ or $e_i \in D[a, b..d_i]$. In the former case we have $m_{e_i d} = 2$ for all $d \in C[b..d_i]$, and as above we see that r_{e_i} fixes $X(d_i)$. In the latter case it is still true that r_{e_i} fixes $X(d_i)$, since the only simple reflection of $D[a, b..d_i]$ that does not fix $X(d_i)$ is the one corresponding to the vertex adjacent to d_i . So we have shown that when $m_{e_i d_i} = 2$ it is true that $v[e_i, \{d_i\}]X(d_i)$ and

$v[e_i, \{d_i\}]Y(d_i)$ both lie in the set $\{\pm X(d_{i+1}), \pm Y(d_{i+1})\}$, and it remains to consider the case that e_i and d_i are adjacent in $\text{Odd}(a)$. Note that in this case $d_{i+1} = e_i$.

Let $D[a, b..d_i] = \{b, c_1, \dots, c_m\}$ with $c_1 = a$ and $c_m = d_i$. Suppose first that $m > 2$, and suppose that $e_i = c_{m-1}$ is the vertex adjacent to d_i in $D[a, b..d_i]$. Then

$$\begin{aligned} v[e_i, \{d_i\}]X(d_i) &= r_{c_m}r_{c_{m-1}}\left(b + a + c_m + 2\sum_{j=2}^{m-1} c_j\right) \\ &= r_{c_m}\left(b + a + c_m + c_{m-1} + 2\sum_{j=1}^{m-2} c_j\right) \\ &= b + a + c_{m-1} + 2\sum_{j=1}^{m-2} c_j = X(e_i). \end{aligned}$$

If $m = 2$ and $e_i = b$ then

$$v[e_i, \{d_i\}]X(d_i) = r_b r_{d_i}(a + b + d_i) = a = X(b) = X(e_i),$$

and if $e_i = a$ then similarly

$$v[e_i, \{d_i\}]X(d_i) = r_a r_{d_i}(a + b + d_i) = b = X(a) = X(e_i).$$

The alternative possibility is that $d_i = c_{m-1}$ is the vertex adjacent to $e_i = c_m$ in $D[a, b..e_i] = \{b, c_1, \dots, c_m\}$. We calculate that

$$\begin{aligned} v[e_i, \{d_i\}]X(d_i) &= r_{c_{m-1}}r_{c_m}\left(b + a + c_{m-1} + 2\sum_{j=2}^{m-2} c_j\right) \\ &= r_{c_{m-1}}\left(b + a + c_m + c_{m-1} + 2\sum_{j=1}^{m-2} c_j\right) \\ &= b + a + c_m + 2\sum_{j=1}^{m-1} c_j = X(e_i), \end{aligned}$$

as required.

The induction is now complete, and it follows that $wb = w_k b = \pm X(a')$. Hence

$$wb \in \Phi_{D[a, b..a']} \subseteq \Phi_{D[a, b..c]} \subseteq \Phi_L.$$

Thus $wr_b w^{-1} \in W_L$, and so $r_b \in w^{-1}W_L w = F$. Since F was an arbitrary maximal finite subgroup of W containing r_a , this shows that $r_b \in \text{FC}(r_a)$.

Note that since W has a graph automorphism that swaps r_a and r_b and fixes all the other simple reflections, it must also be true that $r_a \in \text{FC}(r_b)$.

Let \tilde{M} be the component of $\text{EOdd}(a)$ containing $\text{Odd}(a)$, and suppose, for a contradiction, that \tilde{M} is spherical. Note that $\tilde{M} \neq \text{Odd}(a)$, in view of condition (5) of Definition 5. So $\tilde{M} \setminus \text{Odd}(a) \neq \emptyset$. But for all $e \in \tilde{M} \setminus \text{Odd}(a)$ and $c \in \text{Odd}(a)$ we have $m_{ce} \neq \infty$, since \tilde{M} is spherical, and by conditions (1) and (5) of Definition 5 it follows that $m_{ce} = 2$ for all $c \in \text{Odd}(a)$. This contradicts the fact that \tilde{M} is connected.

Suppose that $a' \in \text{Odd}(a)$ is such that $\text{FC}(r_{a'}) = W_J$ for some $J \subseteq \Pi$, and let J_0 be the component of J containing a' . Since $J_0 \neq \tilde{M}$ it follows from Proposition 35 that $J \cap \text{Odd}(a)$ has rank at most 2. Now suppose, for a contradiction, that $a' \notin \{a, b\}$. Since there exists an element $w \in W_{D[a, b..a']}$ such that $wa = a'$ and $wb = X(a')$, it follows that $r_{wb} = wr_bw^{-1} \in \text{FC}(wr_aw^{-1}) = \text{FC}(r_{a'})$. Thus $X(a') \in \Phi_J$, and so $D[a, b..a'] \subseteq J$, contradicting the fact that the rank of $J \cap \text{Odd}(a)$ is at most 2. So we deduce that $a' = b$ or $a' = a$. Moreover, in either case we know that $\{a, b\} \subseteq J \cap \text{Odd}(a)$, and since $J \cap \text{Odd}(a)$ has rank at most 2 it follows that $J \cap \text{Odd}(a) = \{a, b\}$.

By Lemma 36 we know that all spherical components of $\text{EOdd}(a)$ are components of J , and by Proposition 35 all other components of J that intersect $\text{Odd}(a)$ trivially correspond to C_3 -neighbours of $\text{Odd}(a)$. But clearly the conditions of Definition 6 imply that $\text{Odd}(a)$ has no C_3 -neighbours. So we conclude that J is the union of $\{a, b\}$ and the spherical components of $\text{EOdd}(a)$, as required. \square

Proof of Theorem 7. Let M be a connected component of $\Omega(\Pi)$, and write \tilde{M} for the component of $\text{E}(M)$ containing M .

Suppose first that \tilde{M} is spherical, so that the conditions of Case A of Theorem 7 are satisfied, and let $a \in M$ be arbitrary. Observe that all C_3 -neighbours of M are contained in \tilde{M} . Choose $a' \in M$ such that $\text{FC}(r_{a'})$ is visible, and let $\text{FC}(r_{a'}) = W_J$. By Lemma 36 we know that \tilde{M} is contained in J , and hence $a \in J$. So by Proposition 25 it follows that $\text{FC}(r_a) = W_J$ also. By Proposition 35 the only possible components of J apart from J_0 are the other spherical components of $\text{E}(M)$, and by Lemma 36 all of these are indeed components of J . So J consists of the spherical components of $\text{E}(M)$, as required.

Now suppose that \tilde{M} is not spherical. If there exists a $b \in \Pi \setminus M$ such that (a, b) is a focus of M then it follows from Proposition 38 that $\text{FC}(r_a) = W_J$, where J is composed of $\{a, b\}$ and the spherical components of $\text{E}(M)$, and $\text{FC}(r_{a'})$ is not visible for any $a' \in \text{Odd}(a) \setminus \{a\}$. Similarly, if there exists a $b \in M$ such that $\{a, b\}$ is a half-focus of M , then it follows from Proposition 39 that $\text{FC}(r_a) = \text{FC}(r_b) = W_J$, where J is composed of $\{a, b\}$ and the spherical components of $\text{E}(M)$, and $\text{FC}(r_{a'})$ is not visible for any $a' \in \text{Odd}(a) \setminus \{a, b\}$.

Finally, suppose that \tilde{M} is not spherical and M does not have a focus or a half focus. Suppose that $a \in M$ is such that $\text{FC}(r_a) = W_J$ for some $J \subseteq \Pi$, and let K be the component of J containing a .

Suppose first that alternative (b) of Proposition 35 holds, so that $K = \{a, b\}$ is of type C_2 , and $J \cap M = \{a\}$. Since M does not have any focus in Π , it follows from Proposition 28 that $M \cup \{b\}$ is a spherical component of $E(M)$. But the component of $E(M)$ containing M is \tilde{M} , which, by our assumptions, is not spherical. So this case does not arise.

Alternative (c) of Proposition 35 is similarly impossible, by Proposition 29, and alternative (e) is also incompatible with the assumption that \tilde{M} is not spherical. So we conclude that alternative (a) holds: $K = \{a\} = J \cap \text{Odd}(a)$. Note that all spherical components of $E(M)$ are components of J , and by Proposition 35 the only other possible components are the sets $\{b\}$ such that b is a C_3 -neighbour of M .

Suppose that b is a C_3 -neighbour of M that is adjacent to a . Let \tilde{a} be the unique neighbour of a in M . By Lemma 37 we know that $r_b \in \text{FC}(r_{\tilde{a}})$, and so it follows that $\text{FC}(r_a) = r_{\tilde{a}}r_a \text{FC}(r_{\tilde{a}})r_ar_{\tilde{a}}$ contains the reflection along the root $r_{\tilde{a}}r_ab = b + \sqrt{2}a + \sqrt{2}\tilde{a}$. Since $\text{FC}(r_a) = W_J$, it follows that both b and \tilde{a} are in J . But this contradicts the fact that the component of J containing a is just $\{a\}$.

This reasoning has shown that if $a \in M$ is adjacent in Π to a C_3 -neighbour of M then $\text{FC}(r_a)$ is not visible. On the other hand, we know that there is at least one $a \in M$ such that $\text{FC}(r_a)$ is visible. So we may choose an $a \in M$ such that $\text{FC}(r_a) = W_J$ for some $J \subseteq \Pi$. Since a is not adjacent to any C_3 -neighbour of M it follows by Lemma 37 that all C_3 -neighbours of M are in J . So we conclude that $J = J' \cup \{a\}$, where J' is the union of the spherical components of $E(M)$ and the C_3 -neighbours of M .

It remains to prove that if a' is any other element of M that is not adjacent to any C_3 -neighbour of M then $\text{FC}(r_{a'}) = W_{J' \cup \{a'\}}$. Given such an a' , since a' lies in $M = \text{Odd}(a)$, we may choose $w \in W$ such that $wa = a'$. By Proposition 12 we see that $w \in W_{\tilde{M}}$, and so w fixes all other components of $E(M)$. And w fixes all C_3 -neighbours of M , by Lemma 34. So w fixes J' , and it follows that

$$\text{FC}(r_{a'}) = w \text{FC}(r_a)w^{-1} = wW_{J' \cup \{a\}}w^{-1} = W_{wJ' \cup \{wa\}} = W_{J' \cup \{a'\}},$$

as required. This completes the proof of Theorem 7.

Proof of Theorem 1. Let $a \in \Pi$ and $M = \text{Odd}(a)$. As Π is 2-spherical it follows that $\Pi = E(M)$, and, as Π is non-spherical, it follows that Case A of Theorem 7 does not hold for M . As there are no ∞ -labels in the Coxeter graph of Π , Cases C and D do not hold either. Hence we are in Case B. Since there are no ∞ -labels in the Coxeter graph of Π , there are no C_3 neighbors of M . As $E(M) = \Pi$ is irreducible, there are no spherical components of $E(M)$. It follows now from Theorem 7 that there is an $a' \in \text{Odd}(a)$ such that $\text{FC}(r_{a'}) = \langle r_{a'} \rangle$. As r_a and $r_{a'}$ are W -conjugate we have $\text{FC}(r_a) = \langle r_a \rangle$ as well, and this completes the proof of Part a) of Theorem 1.

Let $S \subseteq W$ be such that (W, S) is a Coxeter system. It follows from Part a) and Corollary 24 that $r_a \in S^W$ for each $a \in \Pi$, and hence $\{r_a \mid a \in \Pi\} \subseteq S^W$. As Π

is assumed to be non-spherical, irreducible and 2-spherical, it follows now from the main result of [5] that there is an element $w \in W$ such that $\{r_a \mid a \in \Pi\} = S^w$. This completes the proof of Part b) of Theorem 1. As Part c) is an immediate consequence of Part b) we are done.

References

- [1] P. Bahls and M. Mihalik, Reflection Independence in even Coxeter Groups. *Geom. Dedicata* **110** (2005), 63–80. Zbl 1072.20043 MR 2136020
- [2] N. Bourbaki, *Groupes et algèbres de Lie*, Chapitres 4, 5 et 6. Actualités Scientifiques et Industrielles 1337, Hermann, Paris 1968. Zbl 0186.33001 MR 0240238
- [3] N. Brady, J. McCammond, B. Mühlherr and W. Neumann, Rigidity of Coxeter Groups and Artin Groups. *Geom. Dedicata* **94** (2002), 91–109. Zbl 1031.20035 MR 1950875
- [4] B. Brink, On centralizers of reflections in Coxeter groups. *Bull. London Math. Soc.* **28** (1996), 465–470. Zbl 0852.20033 MR 1396145
- [5] P. E. Caprace and B. Mühlherr, Reflection rigidity of 2-spherical Coxeter groups. Preprint 2003, 24pp, to appear in *Proc. London Math. Soc.*.
- [6] Roger W. Carter, *Finite Groups of Lie Type: Conjugacy Classes and Complex Characters*. J. Wiley & Sons, 1985. Zbl 0567.20023 MR 0794307
- [7] V. V. Deodhar, On the root system of a Coxeter group. *Comm. Algebra* **10** (1982), 611–630. Zbl 0491.20032 MR 0647210
- [8] V. V. Deodhar, A note on subgroups generated by reflections in Coxeter groups. *Arch. Math. (Basel)* **53** (1989), 543–546. Zbl 0688.20028 MR 1023969
- [9] M. Dyer, Reflection subgroups of Coxeter systems. *J. Algebra* **135** (1990), 57–73. Zbl 0712.20026 MR 1076077
- [10] W. N. Franzsen and R. B. Howlett, Automorphisms of nearly finite Coxeter groups. *Adv. Geom.* **3** (2003), 301–338. Zbl 1044.20022 MR 1997410
- [11] R. B. Howlett, Normalizers of parabolic subgroups of reflection groups. *J. London Math. Soc. (2)* **21** (1980), 62–80. Zbl 0427.20040 MR 0576184
- [12] R. B. Howlett and B. Mühlherr, Isomorphisms of Coxeter groups which do not preserve reflections. Preprint 2004, 18pp.
- [13] R. B. Howlett, P. J. Rowley and D. E. Taylor, On outer automorphism groups of Coxeter groups. *Manuscripta Math.* **93** (1997), 499–513. Zbl 0888.20023 MR 1465894
- [14] James E. Humphreys, *Reflection Groups and Coxeter Groups*. Stud. in Adv. Math. 29, Cambridge University Press, Cambridge 1990. Zbl 0768.20016 MR 1066460
- [15] B. Mühlherr, The isomorphism problem for Coxeter groups. Preprint 2005, 15pp, to appear in *The Coxeter Legacy: Reflections and Projections*, Fields Institute Communications.
- [16] R. W. Richardson, Conjugacy classes of involutions in Coxeter groups. *Bull. Austral. Math. Soc.* **26** (1982), 1–15. Zbl 0531.20017 MR 0679916

Received February 23, 2005

W. N. Franzsen, Australian Catholic University, 25A Barker Rd, Strathfield NSW 2135, Australia

E-mail: b.franzsen@mary.acu.edu.au

R. B. Howlett, School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia

E-mail: R.Howlett@maths.usyd.edu.au

B. Mühlherr, Département de Mathématiques, ULB C.P. 216, Bd. du Triomphe, 1050 Bruxelles, Belgium

E-mail: bernhard.muhlher@ulb.ac.be

Leere Seite
Blank page
Page vide