

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 83 (2008)

Artikel: Generalized Property R and the Schoenflies Conjecture
Autor: Scharlemann, Martin
DOI: <https://doi.org/10.5169/seals-99036>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 04.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Generalized Property R and the Schoenflies Conjecture

Martin Scharlemann*

Abstract. There is a relation between the generalized Property R Conjecture and the Schoenflies Conjecture that suggests a new line of attack on the latter. The new approach gives a quick proof of the genus 2 Schoenflies Conjecture and suffices to prove the genus 3 case, even in the absence of new progress on the generalized Property R Conjecture.

Mathematics Subject Classification (2000). 57Q25, 57N13.

Keywords. Schoenflies Conjecture, Heegaard splitting, framed surgery, critical level embedding, Property R.

1. Introduction and preliminaries

The Schoenflies Conjecture asks whether every PL (or, equivalently, smooth) 3-sphere in S^4 divides the 4-sphere into two PL balls. The appeal of the conjecture is at least 3-fold:

- The topological version (for locally flat embeddings) is known to be true in every dimension. Both the PL and the smooth versions (when properly phrased, to avoid problems with exotic structures) are known to be true in every *other* dimension.
- If the Schoenflies Conjecture is false, then there is no hope for a PL prime decomposition theorem for 4-manifolds, for it would imply that there are PL 4-manifolds X and Y , not themselves 4-spheres, so that $X \# Y \cong S^4$.
- The Schoenflies Conjecture is weaker than the still unsolved 4-dimensional PL Poincaré Conjecture, and so might be more accessible.

Little explicit progress has been made on the Schoenflies Conjecture for several decades, a time which has nonetheless seen rapid progress in our understanding of both 3- and 4-dimensional manifolds. Here we outline how the Schoenflies Conjecture is connected to another important problem on the border between classical 3- and

*Research partially supported by a National Science Foundation grant. Thanks also to Catalonia's Centre Recerca Matemàtica for their extraordinary hospitality while this work was being completed

4-dimensional topology, namely the generalized Property R Conjecture. We show how at least some of the last two decades of progress in combinatorial 3-dimensional topology, particularly sutured manifold theory, can be used to extend the proof of the Schoenflies Conjecture from what are called genus 2 embeddings of S^3 in S^4 to genus 3 embeddings. In some sense this is a small advance, but it has some philosophical interest: genus 2 surfaces have long been known to have special properties (e.g. the hyperelliptic involution) that are not shared by higher genus surfaces. That this approach works for genus 3 suggests that the special properties of genus 2 surfaces are not needed and so are not a barrier to success for arbitrary, higher genus embeddings.

We work in the PL category throughout. All manifolds discussed are orientable.

2. Generalized Property R

Recall the famous Property R theorem, proven in a somewhat stronger form by David Gabai [Ga2]:

Theorem 2.1 (Property R). *If 0-framed surgery on a knot $K \subset S^3$ yields $S^1 \times S^2$ then K is the unknot.*

It is well known (indeed it is perhaps the original motivation for the Property R Conjecture) that Property R has an immediate consequence for the handlebody structure of 4-manifolds:

Corollary 2.2. *Suppose U^4 is a homology 4-sphere and has a handle structure containing exactly one 2-handle and no 3-handles. Then U is the 4-sphere.*

Remark. It is immediate that U^4 is in fact a homotopy 4-sphere, since the dual handle structure would have no 1-handles and U^4 is simply connected. But this property plays no role in the proof.

Proof. Since U has a handle structure with no 3-handles, dually it has a handle structure with no 1-handles. In order for U to be connected, this dual handle structure must then have exactly one 0-handle, so the original handle structure has a single 4-handle.

Let $U_- \subset U$ be the union of all 0- and 1-handles of U and $M = \partial U_-$. U_- can be thought of as the regular neighborhood of a graph or, collapsing a maximal tree in that graph, as the regular neighborhood of a bouquet of circles. The 4-dimensional regular neighborhood of a circle in an orientable 4-manifold is $S^1 \times D^3$, so U_- is the boundary connected sum $\natural_n(S^1 \times D^3)$, some $n \geq 0$. (Explicitly, the number of summands n is one more than the difference between the number of 1-handles and 0-handles, i.e. $1 - \chi$, where χ is the Euler characteristic of the graph.) It follows

that $M = \#_n(S^1 \times S^2)$ and, in particular, $H_2(M) \cong \mathbb{Z}^n$. Now consider the closed complement U_+ of U_- in U . Via the dual handle structure, U_+ is obtained by attaching a single 2-handle to B^4 , so it deformation retracts to a 2-sphere and, in particular, $H_2(U_+) \cong \mathbb{Z}$. Since U is a homology 4-sphere and $H_2(U_-) = 0$, it follows from the Mayer–Vietoris sequence

$$H_3(U) = 0 \rightarrow H_2(M) \rightarrow H_2(U_+) \oplus H_2(U_-) \rightarrow H_2(U) = 0$$

that $\mathbb{Z} \cong H_2(U_+) \cong \mathbb{Z}^n$, so $n = 1$ and $M = S^1 \times S^2$.

On the other hand, U_+ , whose handle structure (dual to that from U) consists of a 0-handle and a 2-handle, is visibly the trace of surgery on a knot in S^3 , namely the attaching map of the 2-handle. The framing of the surgery is zero, since the generator of $H_2(U_+)$ is represented by $* \times S^2 \subset S^1 \times S^2 \cong M$ and this class visibly has trivial self-intersection. Since the result of 0-framed surgery on the knot is $M = S^1 \times S^2$, the knot is trivial by Property R (Theorem 2.1) so U_+ is simply $S^2 \times D^2$.

Hence U is the boundary union $S^1 \times D^3 \cup_{\partial} S^2 \times D^2$. Of course the same is true of S^4 , since the closed complement of a neighborhood of the standard 2-sphere in S^4 is $S^1 \times D^3$. So we see that U can be obtained from S^4 by removing the standard $S^1 \times D^3$ and pasting it back in, perhaps differently. But it is well known (and is usefully extended to all 4-dimensional handlebodies by Laudenbach and Poenaru [LP]) that any automorphism of $S^1 \times S^2$ extends to an automorphism of $S^1 \times D^3$, so the gluing homeomorphism extends across $S^1 \times D^3$ to give a homeomorphism of U with S^4 . \square

The generalized Property R conjecture (cf. Kirby Problem 1.82) says this:

Conjecture 1 (Generalized Property R). Suppose L is a framed link of $n \geq 1$ components in S^3 , and surgery on L via the specified framing yields $\#_n(S^1 \times S^2)$. Then there is a sequence of handle slides on L (cf. [Ki]) that converts L into a 0-framed unlink.

In the case $n = 1$ no slides are possible, so Conjecture 1 does indeed directly generalize Theorem 2.1. On the other hand, for $n > 1$ it is certainly necessary to include the possibility of handle slides. For if one starts with the 0-framed unlink of n -components and does a series of possibly complicated handle-slides, the result will be a possibly complicated framed link L of n -components. The result of doing the specified framed surgery on L will necessarily be the same (cf. [Ki]) as for the original unlink, namely $\#_n(S^1 \times S^2)$, but L itself is no longer the unlink. The example L is still consistent with Conjecture 1 since simply reversing the sequence of handle slides will convert L back to the framed unlink. So in some sense Conjecture 1 is the broadest plausible generalization of Theorem 2.1.

The generalized Property R Conjecture naturally leads to a generalized Corollary 2.2:

Proposition 2.3. *Suppose Conjecture 1 is true. Then any homology 4-sphere U with a handle structure containing no 3-handles is S^4 .*

Proof. Again focus on the 3-manifold M that separates U_- (the manifold after the 0 and 1-handles are attached) from its closed complement U_+ in U . The dual handle structure on U shows that U_+ is constructed by attaching some 2-handles to B^4 . On the other hand, the original handle structure shows that U_- is the regular neighborhood of a graph, so, as before for some n , $U_- \cong \natural_n(S^1 \times D^3)$ and $M \cong \#_n(S^1 \times S^2)$. In particular $H_2(M) \cong \mathbb{Z}^n$. Since U is a homology 4-sphere and $H_2(U_-) = 0$, it follows as before from the Mayer–Vietoris sequence that $H_2(U_+) \cong \mathbb{Z}^n$. Hence U_+ must be obtained from B^4 by attaching exactly n 2-handles. Then the generalized Property R conjecture would imply that $U_+ \cong \natural_n(S^2 \times D^2)$. It is shown in [LP] that any automorphism of $\#_n(S^1 \times S^2) = \partial \natural_n(S^1 \times D^3)$ extends to an automorphism of $\natural_n(S^1 \times D^3)$. (This is not quite stated explicitly in [LP] beyond the observation on p. 342, “mark that no diffeomorphism of X^p was needed here!”). Hence the only manifold that can be obtained by gluing U_+ to U_- along M is S^4 . \square

The proposition suggests this possibly weaker conjecture:

Conjecture 2 (Weak generalized Property R conjecture). Suppose attaching n 2-handles to B^4 gives a manifold W whose boundary is $\#_n(S^1 \times S^2)$. Then $W \cong \natural_n(S^2 \times D^2)$.

We have then:

Proposition 2.4. *The weak generalized Property R conjecture (Conjecture 2) is equivalent to the conjecture that any homology 4-sphere U with a handle structure containing no 3-handles is S^4 .*

Proof. The proof of Proposition 2.3 really required only Conjecture 2, so only the converse needs to be proved.

Suppose we know that any homology 4-sphere with a handle structure containing no 3-handles is S^4 . Suppose W is a 4-manifold constructed by attaching n 2-handles to B^4 and ∂W is $\#_n(S^1 \times S^2)$. Consider the exact sequence of the pair $(W, \partial W)$:

$$\begin{aligned} 0 = H_3(W, \partial W) &\rightarrow H_2(\partial W) \rightarrow H_2(W) \\ &\rightarrow H_2(W, \partial W) \rightarrow H_1(\partial W) \rightarrow H_1(W) = 0. \end{aligned}$$

Since the last two non-trivial terms are both \mathbb{Z}^n , the inclusion induces an isomorphism of the first two non-trivial terms, $H_2(\partial W) \rightarrow H_2(W) \cong \mathbb{Z}^n$. Attach $V = \natural_n(S^1 \times D^3)$ to W by a homeomorphism of their boundaries and call the result U . (There is an

obvious homeomorphism of boundaries, and any other one will give the same 4-manifold, per [LP]). Then the Mayer–Vietoris sequence for the pair (W, V) shows that U is a homology 4-sphere hence, under our assumption, $U = S^4$.

$V \subset U$ is just a regular neighborhood of the wedge of n circles Γ . Since U is simply connected, Γ is homotopic to a standard (i.e. planar) wedge of circles in U whose complement is $\natural_n(S^2 \times D^2)$. In dimension 4, homotopy of 1-complexes implies isotopy (apply general position to the level-preserving map $\Gamma \times I \rightarrow U \times I$) so in fact $W \cong \natural_n(S^2 \times D^2)$ as required. \square

Setting aside conjecture, here is a concrete extension of Property R:

Proposition 2.5. *Suppose a 2-handle is attached to a genus n 4-dimensional handlebody $N = \natural_n(S^1 \times D^3)$ and the resulting 4-manifold N_- has boundary $\#_{n-1}(S^1 \times S^2)$. Then $N_- \cong \natural_{n-1}(S^1 \times D^3)$.*

Proof. The proof is by induction on n ; when $n = 1$ this is Property R. Suppose then that $n > 1$ and let $K \subset \partial(\natural_n(S^1 \times D^3)) \cong \#_n(S^1 \times S^2)$ be the attaching map for the 4-dimensional 2-handle. The hypothesis is then that surgery on K yields $\#_{n-1}(S^1 \times S^2)$, a reducible manifold. But examining the possibilities in [Sch2] we see that this is possible only if $\#_n(S^1 \times S^2) - K$ is itself reducible, so in particular one of the non-separating 2-spheres $\{*\} \times S^2$ is disjoint from K . Following [LP], this 2-sphere bounds a 3-ball in N . Split N along this 3-ball, converting N to $\natural_{n-1}(S^1 \times D^3)$ and ∂N_- to $\#_{n-2}(S^1 \times S^2)$. By inductive hypothesis, the split open N_- is $\natural_{n-2}(S^1 \times D^3)$ so originally $N_- \cong \natural_{n-1}(S^1 \times D^3)$. \square

Remark. Experts will note that, rather than use [Sch2], one can substitute the somewhat simpler [Ga1]: If $n > 1$ then $H_2(\#_n(S^1 \times S^2) - \eta(K)) \neq 0$. Since both ∞ - and 0-framed surgery on K (or on a companion solid torus if $\#_n(S^1 \times S^2) - \eta(K)$ is toroidal) yield reducible (hence non-taut) 3-manifolds, from [Ga1] it follows that $\#_n(S^1 \times S^2) - \eta(K)$ is itself not taut, hence is reducible.

3. Application: Heegaard unions

Let $H^n = \natural_n(S^1 \times D^2)$ denote a 3-dimensional genus n orientable handlebody and $J^n = \natural_n(S^1 \times D^3)$ denote a 4-dimensional genus n orientable handlebody. H^n and J^n can also be thought of as regular neighborhoods in, respectively, R^3 and R^4 of any graph Γ with Euler characteristic $\chi(\Gamma) = 1 - n$.

Definition 3.1. Suppose, for some $\rho_0, \rho_1, \rho_2 \in \mathbb{N}$, H^{ρ_0} is embedded into both ∂J^{ρ_1} and ∂J^{ρ_2} so that its complement in each ∂J^{ρ_i} , $i = 1, 2$ is also a handlebody. Then the 4-manifold $W = J^{\rho_1} \cup_{H^{\rho_0}} J^{\rho_2}$ is called the *Heegaard union* of the J^{ρ_i} along H^{ρ_0} . See Figure 1.




Figure 1

The term Heegaard union comes from the fact that H^{ρ_0} is half of a Heegaard splitting of both ∂J^{ρ_1} and ∂J^{ρ_2} . Moreover, if W is such a Heegaard union, then $(\partial J^{\rho_1} - H^{\rho_0}) \cup_{\partial H^{\rho_0}} (\partial J^{\rho_2} - H^{\rho_0})$ is a Heegaard splitting of ∂W . The construction here is tangentially related to the construction in [BC, 2.4] of a 4-dimensional cobordism between three Heegaard-split 3-manifolds. Indeed, if two of the three 3-manifolds in the Birman–Craggs construction are of the form $\#_i(S^1 \times S^2)$ and are then filled in with copies of $\#_i(S^1 \times D^3)$ the result is a Heegaard union.

Lemma 3.2. *If a Heegaard union $W = J^{\rho_1} \cup J^{\rho_2}$ is a rational homology ball, then $\rho_0 = \rho_1 + \rho_2$.*

Proof. The first and second homology groups (rational coefficients) of W are trivial, so the result follows from the Mayer–Vietoris sequence of $W = J^{\rho_1} \cup_{H^{\rho_0}} J^{\rho_2}$:

$$H_2(W) = 0 \rightarrow H_1(H^{\rho_0}) \rightarrow H_1(J^{\rho_1}) \oplus H_1(J^{\rho_2}) \rightarrow H_1(W) = 0. \quad \square$$

Proposition 3.3. *Suppose a Heegaard union $W = J^{\rho_1} \cup_{H^{\rho_0}} J^{\rho_2}$ is a homology ball and $\partial W \cong S^3$. If the weak generalized Property R conjecture (Conjecture 2) is true for $\min\{\rho_1, \rho_2\}$ components, then $W = B^4$.*

Proof. Suppose with no loss of generality that $\rho_1 \leq \rho_2$. Let J_i denote J^{ρ_i} , $i = 1, 2$ and H_0 denote H^{ρ_0} . Consider the genus ρ_0 Heegaard splitting of ∂J_2 given by $H_0 \cup_{\partial H_0} (\partial J_2 - H_0)$. According to Waldhausen [Wa] there is only one such Heegaard splitting of ∂J_2 up to homeomorphism, obtained as follows: Regard J_2 as the product of the interval with a genus ρ_2 3-dimensional handlebody H . Then $H \times \{0\} \subset \partial(H \times I) = \partial J_2$ and $\partial J_2 - (H \times \{0\})$ are both 3-dimensional handlebodies. The resulting Heegaard splitting of ∂J_2 is called the *product splitting*. It can be regarded as the natural Heegaard splitting of $\partial J_2 \cong \#_{\rho_2}(S^1 \times S^2)$. Any other Heegaard splitting (e.g. the genus ρ_0 splitting at hand) is homeomorphic to a stabilization of this standard splitting.

As proven in [LP] and noted above, any automorphism of ∂J_2 extends over J_2 itself, so we may as well assume that the Heegaard splitting $H_0 \cup_{\partial H_0} (\partial J_2 - H_0)$

actually is a stabilization of the product splitting. In particular, and most dramatically, if $\rho_0 = \rho_2$ then no stabilization is required, so J_2 is just $H_0 \times I$ and $W \cong J_1$. Much the same is true if $\rho_0 = \rho_2 + 1$: most of H_0 is just H , so its attachment to J_1 has no effect on the topology of J_1 . The single stabilization changes the picture slightly, and is best conveyed by considering what the effect would be of attaching a 4-ball to J_1 *not* along one side of the minimal genus splitting of ∂B^4 (i.e. along $B^3 \subset S^3$), which clearly leaves J_1 unchanged, but rather along one side of the once-stabilized splitting of ∂B^4 . That is, B^4 is attached to J_1 along a solid torus, unknotted in ∂B^4 . But this is exactly a description of attaching a 2-handle to J_1 . So W can be viewed as J_1 with a single 2-handle attached. In the general situation, in which the product splitting is stabilized $\rho_0 - \rho_2$ times, W is homeomorphic to J_1 with $\rho_0 - \rho_2$ 2-handles attached. The result now follows from Lemma 3.2 and Proposition 2.4. \square

Remark. The link along which the 2-handles are attached has ρ_1 components and, viewed in S^3 , is part of a genus ρ_0 Heegaard splitting. So its tunnel number can be calculated: $\rho_1 - 1$ tunnels are needed to connect the link into a genus ρ_1 handlebody, and another $\rho_0 - \rho_1$ are needed to make it half of a Heegaard splitting. Hence the tunnel number is $\rho_0 - 1$. This fact may be useful, but anyway explains why [Sch1] could be done just knowing Property R for tunnel number one knots.

Corollary 3.4. *Suppose a Heegaard union $W = J^{\rho_1} \cup_{H^{\rho_0}} J^{\rho_2}$ is a homology ball and $\partial W \cong S^3$. If $\rho_0 \leq 3$ then $W = B^4$.*

Proof. By Lemma 3.2, $\rho_1 + \rho_2 \leq 3$, hence $\min\{\rho_1, \rho_2\} \leq 1$. The result then follows from Proposition 3.3 and Theorem 2.1. \square

4. Handlebody structure on 3-manifold complements

Suppose $M \subset S^4$ is a connected closed PL or smooth 3-submanifold. In this section we discuss the handlebody structure of each complementary component of M .

It is a classical result (cf. [KL]) that M can be isotoped so that it is in the form of a *rectified critical level embedding*. We briefly review what that means.

Informally, the embedding $M \subset S^4$ is in the form of a critical level embedding if it has a handle structure in which each handle is horizontal with respect to the natural height function on S^4 , and M intersects each region of S^4 between handle levels in a vertical collar of the boundary of the part of M that lies below (or, symmetrically, above). More formally, regard S^4 as the boundary of $D^4 \times [-1, 1]$, so S^4 consists of two 4-balls $D^4 \times \pm 1$ (called the poles) added to the ends of $S^3 \times [-1, 1]$. Let $p: S^3 \times [-1, 1] \rightarrow [-1, 1]$ be the natural projection. For $-1 < t < 1$ denote $p^{-1}(t)$ by S_t^3 . Then $M \subset S^3 \times [-1, 1] \subset S^4$ is a critical level embedding if there are a

collection $t_1 < t_2 < \cdots < t_n$ of values in $(-1, 1)$ and a collection of closed surfaces $F_1, \dots, F_n \subset S^3$ so that

- (1) $p(M) = [t_1, t_n] \subset (-1, 1)$.
- (2) For each $1 \leq i \leq n-1$, $M \cap (S^3 \times (t_i, t_{i+1})) = F_i \times (t_i, t_{i+1})$.
- (3) $M \cap S_{t_1}^3 = B^3$ with boundary F_1 .
- (4) For each $2 \leq i \leq n$, F_i is obtained from F_{i-1} by a j -surgery, some $0 \leq j \leq 3$. That is, there is a 3-ball $D^j \times D^{3-j} \subset S^3$ incident to F_{i-1} in $\partial D^j \times D^{3-j}$ and F_i is obtained from F_{i-1} by replacing $\partial D^j \times D^{3-j}$ with $D^j \times \partial D^{3-j}$.
- (5) For each $2 \leq i \leq n$, $M \cap S_{t_i}^3$ is the trace of the surgery above. That is, it is the union of F_{i-1} , F_i and $D^j \times D^{3-j}$.

Such an embedding gives rise to a handle structure on M with n handles added successively at levels t_1, \dots, t_n . j is the index of the handle $D^j \times D^{3-j}$. A critical level embedding is called *rectified* if, for $0 \leq j \leq 2$, each handle of index j occurs at a lower level than each handle of index $j+1$. Furthermore, all 0- and 1-handles lie below S_0^3 and all 2- and 3-handles lie above S_0^3 . See Figure 2.




Figure 2

Note that the surface $M \cap S_0^3$ is a Heegaard surface for M , since all 0- and 1-handles lie on one side (namely in $S^3 \times [-1, 0]$) and all 2- and 3-handles lie on the other ($S^3 \times [0, 1]$). In particular, $M \cap S_0^3$ is connected. It is easy to see, [Sch1, Lemma 1.4], though not completely obvious, that if the first 1-handle attached to the boundary of a 0-handle is incident to the 0-handle at only one end, then the handles cancel and there is a rectified embedding of M in which neither handle appears. So, minimizing the number of handles, we will henceforth assume that the first 1-handle incident to each 0-handle is incident to it in both ends. Equally important is the dual to this remark: the boundary of the core of any 2-handle is essential in the surface to which the 2-handle is attached. To summarize:

Lemma 4.1. *Any rectified critical level embedding of M may be isotoped rel $M \cap S_0^3$ to a rectified critical level embedding with no more (but perhaps fewer) handles of any index, such that*

- *the first 1-handle incident to each 0-handle is incident to it in both ends, and*
- *the core of any 2-handle attached in S_t^3 is a compressing disk for $M \cap S_{t-\varepsilon}^3$.*

We will henceforth consider only rectified critical level embeddings with these two properties.

Definition 4.2. The *genus* of the embedding of M in S^4 is the genus of the Heegaard surface $M \cap S_0^3$.

It will be important to understand how a rectified critical level embedding induces a handlebody structure on each of its closed complementary components X and Y . Let X denote the component of $S^4 - M$ that contains the south pole $D^4 \times \{-1\}$. For each generic $t \in (-1, 1)$ let Y_t^- , (resp. X_t^- , M_t^-) be the part of Y (resp. X , M) lying below level t or, more formally, the 4-manifold $Y \cap (S^3 \times [-1, t])$ (resp. $X \cap (S^3 \times [-1, t])$, 3-manifold $M \cap (S^3 \times [-1, t])$). Symmetrically, let Y_t^+ , (resp. X_t^+ , M_t^+) be the part of Y (resp. X , M) lying above level t , that is, the 4-manifold with boundary $Y \cap (S^3 \times [t, 1])$ (resp. $X \cap (S^3 \times [t, 1])$, 3-manifold with boundary $M \cap (S^3 \times [t, 1])$). Finally, let Y_t^* , (resp. X_t^* , M_t^*) be the part of Y (resp. X , M) lying at level t , that is the 3-manifold with boundary $Y \cap S_t^3$ (resp. $X \cap S_t^3$, closed surface $M \cap S_t^3$). Thus ∂Y_t^- is the union of M_t^- and Y_t^* . If $t_i < t < t_{i+1}$ then $\partial M_t^- = \partial M_t^+ = M_t^* = F_i \subset S_t^3$ and Y_t^* consists of a collection of closed complementary components of F_i in S_t^3 . Each component of F_i in S_t^3 is incident to Y_t^* on exactly one side and to X_t^* on the other.

Clearly as long as no t_i lies between the values $t < t'$, then $Y_t^\pm \cong Y_{t'}^\pm$, since the region between them is just a collar on part of the boundary. On the other hand, for each t_i , consider the relation between $Y_{t_i-\varepsilon}^-$ and $Y_{t_i+\varepsilon}^-$. We know that F_i is obtained from F_{i-1} by doing j -surgery along a j -disk in $S^3 - F_{i-1}$. If that j -handle lies on the Y side of F_{i-1} (in the sphere $S_{t_i-\varepsilon}^3$) then $Y_{t_i+\varepsilon}^*$ is homeomorphic to just $Y_{t_i-\varepsilon}^*$ with that j -handle removed. So $Y_{t_i+\varepsilon}^-$ is still just $Y_{t_i-\varepsilon}^-$ with a collar added to part of its boundary, but only to the complement of the j -handle in $Y_{t_i-\varepsilon}^*$. Hence it is still true that $Y_{t_i+\varepsilon}^- \cong Y_{t_i-\varepsilon}^-$. On the other hand, if the j -handle lies on the X side of F_{i-1} , then $Y_{t_i+\varepsilon}^-$ is homeomorphic to $Y_{t_i-\varepsilon}^-$ but with a (4-dimensional) j -handle added, namely the product of the interval $[t_i, t_i + \varepsilon]$ with the 3-dimensional j -handle added to $M_{t_i-\varepsilon}^-$ in $S_{t_i}^3$.

We have then the general rule, sometimes called the rising water rule (cf. Figure 3):

Lemma 4.3. (1) *If the j -surgery at level t_i has its core in Y , then $Y_{t_i+\varepsilon}^- \cong Y_{t_i-\varepsilon}^-$.*

(2) *If the j -surgery at level t_i has its core in X , then $Y_{t_i+\varepsilon}^- \cong Y_{t_i-\varepsilon}^-$ with a j -handle attached.*




Figure 3

Of course the symmetric statements hold for X . Note that since X contains the south pole, $X_{t_0-\varepsilon} \cong B^4$ whereas $Y_{t_0-\varepsilon} = \emptyset$. Just as M_0^* is a Heegaard surface for M , X_0^- and Y_0^- are connected 4-manifolds, constructed from just 0- and 1-handles. In other words, there are integers $n_x, n_y \geq 0$ so that $X_0 = \natural_{n_x}(S^1 \times D^3)$ and $Y_0 = \natural_{n_y}(S^1 \times D^3)$.

Each handle in M_0^- corresponds to a handle of the same index in exactly one of X_0^- or Y_0^- , so there is a connection between n_x, n_y and the genus g of M_0^* : The critical level embedding defines a handlebody structure on M_0^- with a 0-handles and b 1-handles, where

$$b - a + 1 = g.$$

If $a > g$ then there would be at least one 0-handle in the critical level embedding which is first incident to a 1-handle on a single one of its ends, violating the Handle Cancellation Lemma 4.1. So

$$a \leq g.$$

Let a_x, a_y (resp. b_x, b_y) denote the number of 0- (resp. 1-) handles in the critical level embedding whose cores lie in X and Y . We have from above that $a_x + a_y = a$, $b_x + b_y = b$, $n_x = b_y - a_y$ and $n_y = b_x - a_x + 1$. (The asymmetry is explained by noting that the south pole is a 0-handle for X .) It follows that

$$n_x + n_y = g.$$

Another way of counting n_x and n_y is this: Suppose a 1-handle at critical level t_i has its core lying in X , say. If the ends of the 1-handle lie in distinct components of F_{i-1} then the 1-handle adds a 1-handle to Y but nothing to its genus. In contrast, if the ends of the 1-handle lie on the same component of F_{i-1} then it adds 1 to the genus of Y . A count of the total number of the latter sort of 1-handles lying in X (resp. Y) gives n_x (resp. n_y).

For everything that has been said about X^- and Y^- there is a dual statement for X^+ and Y^+ , easily obtained by just inverting the height function. The result is that,

beyond the standard 4-dimensional duality of handle structures on X and on Y , there is a kind of 3-dimensional duality between handles in X and handles in Y , induced by the 3-dimensional duality of handles in M . See Figure 4.




Figure 4

To be concrete: X_0^+ is also a solid 4-dimensional handlebody. To determine its genus, consider the core of each 2-handle, say at critical height $t_i > 0$. If the core of the 2-handle lies on the X side of F_{i-1} then the cocore lies on the Y side of F_i so it corresponds to a 1-handle in X_0^+ . This 1-handle adds genus to X_0^+ if and only if the boundary of the 2-handle is non-separating in F_{i-1} .

To see how this occurs, consider the “dual rule” to Lemma 4.3. That is, suppose again that F_i is obtained from F_{i-1} by doing j -surgery along a j -disk in $S^3 - F_{i-1}$ and ask how $Y_{t_i-\varepsilon}^+$ and $Y_{t_i+\varepsilon}^+$ differ. If the j -surgery at level t_i has its core in Y , then, viewed from above instead of below, there is a corresponding $3-j$ surgery with its core in X . So, following the argument of Lemma 4.3, $Y_{t_i-\varepsilon}^+ \cong Y_{t_i+\varepsilon}^+$ with a $(3-j)$ -handle attached. On the other hand, if the core of the j -handle lies in X , Y^+ is unchanged. This might be called the descending hydrogen rule (cf. Figure 5).

To summarize all possibilities:

Lemma 4.4. *Suppose F_i is obtained from F_{i-1} by doing j -surgery along a j -disk in $S^3 - F_{i-1}$.*

- (1) *If the j -surgery at level t_i has its core in Y , then*

- $Y_{t_i+\varepsilon}^- \cong Y_{t_i-\varepsilon}^-$,
- $X_{t_i+\varepsilon}^- \cong X_{t_i-\varepsilon}^-$ with a j -handle attached,
- $Y_{t_i-\varepsilon}^+ \cong Y_{t_i+\varepsilon}^+$ with a $(3-j)$ -handle attached,
- $X_{t_i-\varepsilon}^+ \cong X_{t_i+\varepsilon}^+$.

(2) If the j -surgery at level t_i has its core in X , then

- $Y_{t_i+\varepsilon}^- \cong Y_{t_i-\varepsilon}^-$ with a j -handle attached,
- $X_{t_i+\varepsilon}^- \cong X_{t_i-\varepsilon}^-$,
- $Y_{t_i-\varepsilon}^+ \cong Y_{t_i+\varepsilon}^+$,
- $X_{t_i-\varepsilon}^+ \cong X_{t_i+\varepsilon}^+$ with a $(3-j)$ -handle attached.




Figure 5

Here is a simple example of how this 3-dimensional duality can be useful:

Proposition 4.5. *Suppose there is a rectified critical level embedding of $M = S^3$ in S^4 so that the 0- and 1-handles, as they are successively attached, all lie on the X -side. Then $X \cong B^4$.*

Proof. Following Lemma 4.4, X has no 0 or 1-handles, so it only has 2- and 3-handles. Dually (in the standard 4-dimensional handle duality of X), X can be constructed with only 1 and 2-handles. Neither of these statements, in itself, is enough to show that X is a 4-ball.

Consider, however, what the given information tells us about Y , following Lemma 4.4 applied to the construction of Y from above: The possible 2- and 3-handles in the construction of X from below correspond respectively to 1- and 0-handles in the construction of Y from above. Similarly, the lack of 0- and 1-handles (beyond the south pole) for X constructed from below corresponds to a lack of 3- and 2-handles for Y when constructed from above. Hence Y has only 0- and 1-handles, i.e. it is a 4-dimensional handlebody. On the other hand, because it is the complement of S^3 in S^4 it is a homotopy 4-ball, so the handlebody must be of genus 0, i.e. Y is a 4-ball. Then its complement X is also a 4-ball. \square

5. Two proofs of the genus 2 Schoenflies Conjecture

Informed by the ideas above, we present two proofs of the genus 2 Schoenflies Conjecture. The first is similar in spirit (though different in detail) to the original proof of [Sch1]. The second uses a different approach, one that aims to simplify the picture by reimbedding X or Y .

Here is a more general statement, relevant to the classical approach:

Proposition 5.1. *Suppose a 3-sphere M has a genus g rectified critical level embedding in S^4 with at most two 0-handles or at most two 3-handles. If the generalized Property R conjecture is true for links of $g - 1$ components then M divides S^4 into two PL 4-balls.*

Proof. Perhaps inverting the height function, assume without loss of generality that M has at most two 3-handles. The roles of X and Y can be interchanged by passing the lowest 0-handle over the south pole, so we can also assume without loss of generality that the first (that is, the lowest) 3-handle for M lies in Y and so represents the addition of a 3-handle to X . The second 3-handle (and so the last handle) of M either lies in X or in Y , but these options are isotopic by passing the handle over the north pole. So, via an isotopy of this handle, we can choose whether both 3-handles of M lie in Y (and so represent attaching of 3-handles to X and not Y) or one each lies in X and Y . See Figure 6.




Figure 6

Now consider the genera n_x and n_y of the 4-dimensional handlebodies X_0^- , Y_0^- , with $g = n_x + n_y$. If $n_x = 0$ then X_0^- is a 4-ball. X is obtained from this 4-ball by attaching some number of 2 and 3-handles, and also a 4-handle if the north pole of S^4 lies in X . There are as many, total, of 2- and 4-handles as there are 3-handles (since X is a homotopy 4-ball) and the argument of the previous paragraph ensures that we can arrange it so that X contains at most one 3-handle. Viewed dually, this means that X can be constructed from $\partial X = S^3$ with no 3-handles, and at most one each of 1- and 2-handles. The result then follows from Corollary 2.2.

If $n_x \geq 1$ then $n_y \leq g - 1$ and, first arranging as above so that Y has no 3-handles, the result again follows from the proof of Proposition 2.3. \square

Corollary 5.2 ([Sch1]). *Each complementary component of a genus 2 embedding of $M = S^3$ in S^4 is a 4-ball.*

Proof. As noted above, we can assume that the number a of 0-handles in the rectified embedding of M is no larger than $g = 2$. Proposition 5.1 then shows the result follows from Property R, via Corollary 2.2. \square

The reimbedding proof of the genus 2 Schoenflies Conjecture begins with a more general claim that follows from our results above for Heegaard unions:

Proposition 5.3. *Suppose $M \cong S^3$ has a rectified critical level embedding in S^4 so that Y_0^* (resp. X_0^*) is a handlebody of genus ρ_0 . If the generalized Property R conjecture is true for $\lfloor \rho_0/2 \rfloor$ components then $Y \cong B^4$ (resp. $X \cong B^4$).*

Proof. It was noted above that Y_0^- is a 4-dimensional handlebody and M_0^- is a 3-dimensional handlebody. The latter fact, and the hypothesis, imply that $M_0^- \cup Y_0^*$ is a Heegaard splitting of ∂Y_0^- . Viewing the critical level embedding from the top down we symmetrically see that Y_0^+ is a 4-dimensional handlebody and Y is a Heegaard union of Y_0^- and Y_0^+ along Y_0^* .

Let ρ_1, ρ_2 denote the genera of Y_0^- and Y_0^+ respectively. Since M is a 3-sphere, each complementary component of M is a homotopy 4-ball. In particular, following lemma 3.2, $\rho_1 + \rho_2 = \rho_0$. The result now follows from Proposition 3.3. \square

Proposition 5.3 suggests a clear strategy for a proof of the general Schoenflies Conjecture, assuming the generalized Property R Conjecture: Given a rectified critical level embedding of $M = S^3$ in S^4 , try to reimbed X (or Y), still a rectified critical level embedding, so that afterwards, either the 3-manifold X_0^* or its complement Y_0^* is a handlebody, or at least more closely resembles a handlebody. *For even if a series of reimbeddings, first of X , then of its new complement Y' , then of the new complement of Y' , etc, eventually leads to a handlebody cross-section at height 0, we are finished.* Indeed, once one of the complementary components of the multiply reimbedded M is a 4-ball, we have that both are, hence the previous complementary components, in succession, leading back to the original X and Y are all 4-balls. (This is more formally explained in the proof of Corollary 8.2.) What follows is a proof of the genus 2 Schoenflies Conjecture built on this strategy.

In order to be as flexible as possible in reimbedding X or Y we first prove a technical lemma which roughly shows that, at the expense of some vertical rearrangement of the 3-handles (or, dually, the 0-handles), the core of a 2-handle (resp, the cocore of a 1-handle) can be moved rel its boundary to another position without affecting the

isotopy class of M or even the embedding of M below the specified 2-handle (resp. above the specified 1-handle).

Suppose, as above, M is a rectified critical level embedded S^3 in S^4 .

Lemma 5.4 (Prairie-Dog Lemma). *Let $E \subset S_{t_i}^3 - F_{i-1}$ be the core of the 2-handle added to F_{i-1} at critical level $t_i > 0$ and let t be a generic height such that $t_{i-1} < t < t_i$. Let $E' \subset S_{t_i}^3 - F_{i-1}$ be another disk on the same side as E , with $\partial E'$ isotopic to ∂E in F_{i-1} . Then there is a proper isotopy of M_t^+ in $S^3 \times [t, 1]$ so that afterwards*

- *the new embedding M' of S^3 is still a rectified critical level embedding,*
- *the critical levels and their indices are the same for M and M' ,*
- *the core of the 2-handle at critical level t_i is E' , and*
- *for any generic level t below the level of the first 3-handle, $M_t^- \cong M_t'^-$.*

Proof. With no loss we take $\partial E'$ parallel (hence disjoint) from ∂E . Let k be the number of 2-handles above level t_i and $n = |E \cap E'|$. The proof is by induction on the pair (k, n) , lexicographically ordered.

Case 1. $k = n = 0$. In this case, the 2-handle attached at level t_i is the last 2-handle attached and E' is disjoint from E . Then the union of E and E' (and a collar between their boundaries) is a 2-sphere in $S^3 - F_{i-1}$; if it bounds a 3-ball in $S^3 - F_{i-1}$ then the disks are isotopic and there is nothing to prove. If it does not bound a 3-ball, let S be the parallel reducing sphere for $S^3 - F_i$ and $B \subset S^3$ be the ball it bounds on the side that does not contain the component to which E and E' are attached. Since t_i is the highest 2-handle, each component of $F_i \cap B$ is a sphere and each is eventually capped off above t_i by a 3-handle.

If all are capped off by 3-handles that lie within B , push all of $M \cap (B \times [t, 1])$ vertically down to a height just above t so that afterwards, E is isotopic to E' in $S_{t_i-\varepsilon}^3 - M$. Perform the isotopy, then push $M \cap (B \times [t, 1])$ back up, so that the 3-handles are attached above height t_i . The number of 3-handles attached (namely, the number of components of F_i) is the same, so, although perhaps rearranged in order, the critical heights at which the 3-handles are attached can be restored to the original set of critical heights. See Figure 7.

The picture is only a little changed if one of the components of F_i lying in B is eventually capped off by a 3-handle not in B . The proof is by induction on the number of such handles. Consider the highest such handle, say at level t_j , capping off a sphere component S of $F_j \cap B$ in $S_{t_j}^3$. Let B' be the 3-ball component of $S_{t_j}^3 - S$ that does lie in B , i.e. the complement of the 3-handle in $S_{t_j}^3$. If there are no components of F_j in the interior of B' then the 3-handle is isotopic to B' via passing over the north pole. This isotopy decreases by one the number of 3-handles not lying in B , completing the inductive step. If some components of F_j do lie in B' note that eventually they are




Figure 7

capped off by 3-handles lying in B' (by choice of t_j). Simply push $M \cap (B' \times [t_j, 1])$ down below level t_j and do the pole pass described above.

Case 2. $k = 0, n > 0$. Consider an innermost disk $E'_0 \subset E'$ cut off by E in E' . Then the union of E'_0 and the subdisk E_0 of E bounded by $\partial E'_0$ is a sphere bounding a ball B whose interior is disjoint from E . If no component of F_i lies in B , E_0 can be isotoped past E'_0 , reducing n by at least one and maybe more, thereby completing the inductive step. If some components of F_i lie in B , then follow the recipe given in Case 1. For example, if all components of $F_i \cap B$ eventually bound 3-handles that lie in B , push $M \cap (B \times [t, 1])$ vertically down to just above level t , do the isotopy, then raise $M \cap (B \times [t, 1])$ back up again.

Case 3. $k > 0$. Depending on whether $n = 0$ or $n \geq 1$, let S and B be the reducing sphere and 3-balls described in cases 1) and 2) above. The inductive hypothesis and a standard innermost disk argument tells us that any 2-handle attached above level t_i can, starting from highest to lowest, be replaced by a 2-handle disjoint from the sphere S . Suppose t_j is the level of the first 3-handle; after the replacement the entire product $S \times [t, t_j]$ is disjoint from M . In fact, following the argument of Case 2, we can isotope the 3-handles of M (possibly rearranging the ordering of the 3-handles) so that all of $S \times [t, 1]$ is disjoint from M . Then push $M \cap (B \times [t, 1])$ down to just above level t , do the isotopy across B as described in Case 1 or 2, then precisely restore the height of $M \cap (S^3 \times [t, 1])$. \square

Note that since all moves are by isotopy, X and Y do not change.

Lemma 5.5 (Torus Unknotting Lemma). *Suppose that Y_0^* (or, symmetrically, X_0^*) lies in a knotted solid torus $W \subset S_0^3$. Let $h_0: W \rightarrow S^1 \times D^2$ be an orientation-preserving homeomorphism to the unknotted solid torus $S^1 \times D^2 \subset S_0^3$. Then there is a reembedding $h: Y \rightarrow S^4$ to a rectified critical level embedding so that $h(Y) \cap S_0^3 = h_0(Y_0^*)$ and the number of handles of each index is unchanged. For t any generic height between the highest 0-handle and lowest 3-handle, both of M_t^\pm are unchanged.*

Proof. If ∂W compresses in X_0^* , there is nothing to prove: ∂ -reduce W to get a 3-ball, which can be isotoped into $S^1 \times D^2 \subset S_0^3$ and that same isotopy can be applied at every level of $S^3 \times I \subset S^4$.

So we assume that ∂W is incompressible in X_0^* . In each successively increasing critical level $t_i > 0$ ask whether the 2-handle attached at t_i can be replaced, as in the Prairie-Dog Lemma 5.4, by a 2-handle that lies in W . If it can be done, then do so. This may alter the critical level embedding of M , but only above level t_{i-1} . If success is possible at the critical levels of all 2-handles, the same can be accomplished for the 3-handles, as described in Case 2 of the proof of Lemma 5.4. Similarly, at each successively decreasing critical level $t_j < 0$ try to replace cocores of 1-handles by disks that lie in W . If this can be done for all 1-handles, then also replace cocores of 0-handles by 3-balls in W . If success is possible for all 1- and 2-handles, hence at all levels, then $M \cap (\partial W \times [-1, 1]) = \emptyset$ and so $Y \subset W \times [-1, 1]$. Then the function $h_0 \times [-1, 1]$ on $W \times [-1, 1]$, restricted to $Y \subset W \times [-1, 1]$, is the required reimbedding.

We are left with the case where successful replacement of the core of a 2-handle or cocore of a 1-handle is not always possible. Suppose, without loss of generality, that $t_i > 0$ is the lowest critical level for which the core of the associated 2-handle cannot be replaced by one that lies in $W \subset S_{t_i}^3$. Without loss, we assume that the replacements of lower 2-handles have been done, so $Y \cap (S^3 \times [0, t_i - \varepsilon]) \subset W \times [0, t_i - \varepsilon]$. In particular, the core of the 2-handle must lie in $X_{t_i - \varepsilon}^*$.

Choose a disk $D \subset X_{t_i - \varepsilon}^*$ so that its boundary is the same as that of the core of the 2-handle and, among all such disks, D intersects ∂W in as few components as possible. An innermost circle of $D \cap \partial W$ then cuts off a subdisk of D whose boundary is essential in ∂W (by choice of D) so, since W is knotted, the subdisk must be a meridian disk for the solid torus W . So at the generic level $t_i - \varepsilon$, ∂W compresses in $X_{t_i - \varepsilon}^* \cap W$. In particular, $Y_{t_i - \varepsilon}^*$ lies in a 3-ball $B \subset W$. It is a classical result that simple coning extends the homeomorphism $h_0|_B: B \rightarrow h_0(B) \subset S^1 \times D^2 \subset S^3$ to an orientation-preserving homeomorphism $H: S^3 \rightarrow S^3$. Define then the embedding of Y_0^+ into $S^3 \times [0, 1]$ so that $h|Y \cap (S^3 \times [0, t_i - \varepsilon]) = h_0 \times [0, t_i - \varepsilon]$ and, for $t \geq t_i - \varepsilon$, $h|Y_t^* = H|Y_t^*$.

The same argument applies symmetrically to construct $h|Y_0^-$. Either all co-cores of 1-handles can be replaced by disks in W , in which case we afterwards simply use h_0 at every level $t \in [-1, 0]$ or there is a highest critical level t_i for which the cocore of the associated handle on M cannot be replaced by a disk in W and we apply the symmetric version of the construction above. \square

Here then is an alternative proof of Corollary 5.2:

Proof. Like any surface in S^3 , the genus 2 surface M_0^* compresses in S^3 , and so it compresses into either X_0^* or Y_0^* , say the former. Maximally compress M_0^* in X_0^* .

If X_0^* is a handlebody then Proposition 5.3 says $X \cong B^4$, hence its complement $Y \cong B^4$.

If X_0^* is not a handlebody, then the surface F resulting from maximally compressing the surface M_0^* into X_0^* consists of one or two tori. Like any surface in S^3 , F compresses in S^3 . The torus component of F that compresses in the complement of F bounds a solid torus W on the side on which the compressing disk lies. That side cannot lie in X_0^* , since F is maximally compressed in that direction, so W must contain M_0^* and so indeed all of Y_0^* . The solid torus W is knotted, else F would still compress further into X_0^* . Now apply the Torus Unknotting Lemma 5.5 to reembed Y in S^4 in a level-preserving way so that afterwards W is unknotted; in particular, afterwards F does compress further into the (new) complement of Y_0^* . After perhaps a further iteration of the argument (when F originally consisted of two tori) we have a level-preserving re-embedding of Y in S^4 so that afterwards its complement is a handlebody. It follows from Proposition 5.3 then that after such a reembedding $S^4 - Y \cong B^4$, hence also $Y \cong B^4$. \square

6. Straightening connecting tubes between tori in S^3

Enlightened by Corollary 3.4, observe that there is no generalized Property R obstacle to applying Proposition 5.3 to the proof of the genus 3 Schoenflies Conjecture. All that is needed is a sufficiently powerful version of the Torus Unknotting Lemma 5.5 that would instruct us how to reembed some complementary component Y of a genus 3 S^3 in S^4 so that its new complement in S_0^3 more closely resembles a handlebody (e.g. it ∂ -reduces to a surface of lower genus than X_0^* or Y_0^* did originally).

Fuelling excitement in this direction is the classic theorem of Fox [Fo] that any compact connected 3-dimensional submanifold of S^3 can be reembedded as the complement of handlebodies. What seems difficult to find is a way to extend such a reembedding of Y_0^* to all of Y , as is done in Lemma 5.5. It is crucial in the proof of Lemma 5.5 that a solid torus has a unique meridian, whereas of course higher genus handlebodies have infinitely many meridians.

For genus 3 embeddings, there is indeed enough information to make such a reembedding strategy work. The key is a genus 2 analogue of the Torus Unknotting Lemma, called the Tube Straightening Lemma. We precede it with a preparatory lemma in 3-manifold topology, whose proof is reminiscent of that in [Ga3] or [Th]:

Lemma 6.1. *Suppose $F \subset S^3$ is a genus 2 surface and $k \subset F$ is a separating curve in F . Denote the complementary components of F by U and V and suppose k bounds a disk E in V so that $U \cup \eta(E)$ is reducible. Suppose $N \subset V$ is a compact 3-manifold. Then at least one of the following must hold:*

- any simple closed curve in ∂N that bounds a disk in $S^3 - N$ bounds a disk in $V - N$,
- N can be isotoped in V to be disjoint from E , or
- k bounds a disk in U .

Proof. Suppose some component of ∂N is a sphere S . Since V is irreducible, S bounds a ball B in V . Nothing is lost by adding B to N , if N is incident to S on the outside of B , or removing $B \cap N$ from N , if N is incident to S on the inside of B . So we may as well assume that ∂N has no sphere components. Essentially the same argument shows that we may take $V - N$ to be irreducible. For if S is a reducing sphere bounding a ball B in V , then any curve in $B \cap \partial N$ that bounds a disk in $S^3 - N$ bounds a disk in $B - N \subset V - N$ and any curve in $\partial N - B$ that bounds a disk in $(V - N) \cup B$ also bounds a disk in $V - N$, so without loss, we may delete $B \cap N$ from N . The proof then will be by induction on $-\chi(\partial N) \geq 0$, assuming now that $V - N$ is irreducible.

Case 1. ∂N does not compress in $V - N$. If ∂N does not compress in $S^3 - N$ either, then the first conclusion holds vacuously. Suppose then that $S^3 - N$ is boundary-reducible. A reducing sphere S for $U \cup \eta(E)$ must separate the two tori that are obtained from F by compressing along E , since otherwise one ball that S bounds in S^3 would lie entirely in $U \cup \eta(E)$. This implies that S intersects V in an odd number of copies of E or, put another way, it intersects U in a planar surface P with an odd number of boundary components lying on a regular neighborhood of k in F . Consider then the result of 0-framed surgery on k in the manifold $S^3 - N$: P can be capped off to give a sphere which is non-separating in the new manifold, since a meridian of $\eta(k)$ intersects P in an odd number of points. On the other hand, $S^3 - N$ itself is ∂ -reducible. No options a)-e) in [Sch2, Theorem 6.2] are consistent with this outcome (in particular the manifold called M' there having a non-separating sphere) so we conclude that $S^3 - (N \cup \eta(k))$ is either reducible or ∂ -reducible. In the latter case, consider how a ∂ -reducing disk D would intersect the surface $F - \eta(k)$. We know the framing of $F \cap \partial\eta(K)$ is a 0-framing (since each component of $F - \eta(k)$ is a Seifert surface for k) so $D \cap F$, if non-empty, consists entirely of simple closed curves. An innermost disk in D cut off by the intersection (perhaps all of D) lies either in U or $V - N$. But each component of $F - \eta(K)$ is a once-punctured torus, so if it compresses in U or $V - N$ so does its boundary, i.e. a copy of k . Hence we have that k bounds a disk in either U or $V - N$. Since V is irreducible, in the latter case the disk can be isotoped to E in V , thereby isotoping N in V off of E .

The same argument applies if $S^3 - (N \cup \eta(k))$ is reducible: since both U and $V - N$ are irreducible, such a reducing sphere must intersect $F - \eta(k)$ and an innermost disk cut off by the intersection leads to the same contradiction.

Case 2. ∂N compresses in $V - N$. The proof is by contradiction: Choose an essential curve $\ell \subset \partial N$ and a compressing disk D for ∂N in $V - N$ so that ℓ bounds a disk

D_ℓ in $S^3 - N$ but does not bound a disk in $V - N$ and, among all such choices of ℓ , D_ℓ , D , $|D \cap D_\ell|$ is minimal. If D and D_ℓ are disjoint, then let $N' = N \cup \eta(D)$. If there is then a sphere component of $\partial N'$, the ball it bounds in V can, without loss, be deleted from N' . In any case (perhaps after deleting the ball if a sphere appears in $\partial N'$), $-\chi(\partial N') < -\chi(\partial N)$ and the inductive hypothesis holds for N' . But the conclusion for N' implies the conclusion for N , which is contained in N' (e.g. if ℓ bounds a disk in $V - N'$ it bounds a disk in $V - N$), so this is impossible.

If D and D_ℓ are not disjoint, note that all curves of intersection must be arcs, else an innermost disk cut off in D could be used to surger D_ℓ and would lower $|D \cap D_\ell|$. Similarly, let D' denote the disk cut off from D by an outermost arc of $D \cap D_\ell$ in D and let $D_{\ell'}, D_{\ell''} \subset V - N$ denote the two disks obtained by the ∂ -compression of D_ℓ to ∂N along D' . Both of these disks intersect D in fewer components than D_ℓ did, so by choice of ℓ and D_ℓ the boundary of each new disk must bound a disk in $V - N$ (of course, if either curve is inessential in ∂N then it automatically bounds a disk in $\partial N \subset V$). A standard innermost disk argument shows that $D_{\ell'}$ and $D_{\ell''}$ can be taken to be disjoint; band them together via the band in ∂N which undoes the ∂ -compression by D' . To be more explicit, note that the arc $\partial D' \cap \partial N$ defined a band move on ∂D_ℓ that split ∂D_ℓ into $\partial D_{\ell'}$ and $\partial D_{\ell''}$. Undo that band move to recover the curve ∂D_ℓ , now bounding a disk (namely the band sum of $D_{\ell'}$ and $D_{\ell''}$) that lies in $V - N$. This contradicts our original choice of ℓ . \square

Lemma 6.2 (Tube Straightening Lemma). *Suppose that $Y \cap S_0^3$ (or, symmetrically, $X \cap S_0^3$) lies in $V \subset S_0^3$, with closed complement U , and $\partial U = \partial V$ is of genus 2. Suppose V contains a separating compressing disk E so that the manifold U_+ obtained from U by attaching a 2-handle along E is reducible. Then there is an embedding $h_0: V \rightarrow S^3$ so that $h_0(\partial E)$ bounds a disk in U' the complement of $h_0(V)$. There is also a reembedding $h: Y \rightarrow S^4$ to a rectified critical level embedding so that*

- $h(Y) \cap S_0^3 = h_0(Y_0^*)$,
- the number of handles of each index is unchanged, and
- for t any generic height between the highest 0-handle and lowest 3-handle, both of M_t^\pm are unchanged.

Proof. Let $h_0: V \rightarrow S^3$ be the reembedding (unique up to isotopy) that replaces the 1-handle in V that is dual to E with a handle intersecting the reducing sphere for U_+ in a single point. Then after the reembedding ∂E bounds a disk in the complement of $h_0(V)$, namely the complement of E in the reducing sphere.

In each successively increasing critical level $t_i > 0$ ask whether the 2-handle attached at t_i can be replaced, as in the Prairie-Dog Lemma 5.4, by a 2-handle that lies in V . If it can, then do so. This may alter the critical level embedding of M , but only above level t_{i-1} . If success is possible at the critical levels of all 2-handles,

the same can be accomplished for the 3-handles, as described in Case 2 of the proof of Lemma 5.4. Similarly, at each successively decreasing critical level $t_j < 0$ try to replace cocores of 1-handles by disks that lie in V . If this can be done for all 1-handles, then also replace cocores of 0-handles by 3-balls in V . If success is possible for all 1- and 2-handles, hence at all levels, then $M \cap (\partial V \times [-1, 1]) = \emptyset$ and so $Y \subset V \times [-1, 1]$. Then the function $h_0 \times [-1, 1]$ on $V \times [-1, 1]$, restricted to $Y \subset V \times [-1, 1]$, is the required reimbedding.

We are left with the case where successful replacement of the core of a 2-handle or cocore of a 1-handle is not always possible. Suppose, without loss of generality, that $t_i > 0$ is the lowest critical level for which the core of the associated 2-handle cannot be replaced by one that lies in $V \subset S^3_{t_i}$. Without loss, we assume that the replacements of lower 2-handles have been done, so $Y \cap (S^3 \times [0, t_i - \varepsilon]) \subset (V \times [0, t_i - \varepsilon])$. In particular, the core of the 2-handle must lie in $X^*_{t_i - \varepsilon}$.

Now apply Lemma 6.1 using $Y^*_{t_i - \varepsilon}$ for N . By assumption, the boundary of the core of the 2-handle bounds no disk in $V \cap X^*_{t_i - \varepsilon}$ so the first possible conclusion of Lemma 6.1 cannot hold. If the last holds, there was nothing to prove to begin with. (Take $h = \text{identity}$.) Hence we conclude that the second conclusion holds: $Y^*_{t_i - \varepsilon}$ can be isotoped to be disjoint from E . But once this is true, the reimbedding h_0 has no effect on $Y^*_{t_i - \varepsilon}$; that is, $Y^*_{t_i - \varepsilon}$ is isotopic to $h_0(Y^*_{t_i - \varepsilon})$. Hence we can define $h|(Y \cap (S^3 \times [0, 1]))$ to be $h_0 \times [0, t_i - \varepsilon]$ on $Y \cap (S^3 \times [0, t_i - \varepsilon])$, followed by a quick isotopy of $h_0(Y^*_{t_i - \varepsilon})$ to $Y^*_{t_i - \varepsilon/2}$ followed by the unaltered embedding above level $t_i - \varepsilon/2$. Note that this unaltered embedding is not necessarily the original embedding, because of changes made while ensuring that earlier 2-handles lie in V .

Finally, the argument can be applied symmetrically on $Y \cap (S^3 \times [-1, 0])$. \square

7. Weak Fox reimbedding via unknotting and straightening

In this section we show that, for a genus 3 surface in S^3 , the operations of torus unknotting and tube straightening described above suffice to give a weak version of Fox reimbedding. That is, for a genus 3 surface $F \subset S^3$ there is a sequence of such reimbeddings, not necessarily all operating on the same complementary component of F , so that eventually a complementary component is a handlebody. Although the context of this section appears to be 3-manifold theory, the notation is meant to be suggestive of the eventual application to the genus 3 Schoenflies Conjecture. In particular, the terms *torus unknotting* and *tube straightening* as used in this section refer to the 3-dimensional reimbedding h_0 given in, respectively, Lemma 5.5 and Lemma 6.2.

Suppose $F \subset S^3$ is a surface dividing S^3 into two components denoted X and Y . Suppose D_1 is a compressing disk for F in X giving rise to a new surface $F_1 \subset S^3$ with complementary components $X_1 = X - \eta(D_1)$ and $Y_1 = X \cup \eta(D_1)$. Suppose

D_2 is a compressing disk for F_1 in X_1 or Y_1 , giving rise to a new surface $F_2 \subset S^3$ with complementary components X_2 and Y_2 . Continue to make such a series of compressions via compressing disks $D_i, i = 1, \dots, n$ so that each D_i lies either in X_{i-1} or Y_{i-1} (the complementary components of $F_{i-1} \subset S^3$) until all components of F_n are spheres.

Definition 7.1. *F can be straightened* if there is a sequence of torus unknottings and tube straightenings of F , as described in Lemmas 5.5 and Lemma 6.2, so that afterwards either

- X or Y is a handlebody, or
- the order of the compressions D_1, \dots, D_n can be rearranged so that afterwards some ∂D_i is inessential in F_{i-1} and so can be eliminated from the sequence.

In the spirit of Lemma 5.4, any disk D_i can be replaced by a disk D'_i in the same complementary component, so long as ∂D_i and $\partial D'_i$ are isotopic in F_{i-1} .

The following series of lemmas assumes we are given such a sequence for $F \subset S^3$, that the sequence is as short as possible, and that the first compressing disk D_1 lies in X . There are, of course, symmetric statements if $D_1 \subset Y$.

Lemma 7.2. *If $D_2 \subset X_1$ then F can be straightened. More specifically, there is a sequence of torus unknottings which convert X into a handlebody*

Proof. Since $X_1 = X - \eta(D_1)$, the disks D_1 and D_2 can be thought of as disjoint disks in X and the compressions given by D_1, D_2 can be performed simultaneously. If any component of F_2 is a sphere, one of the disks is redundant and could be removed from the sequence of compressions. Since $\chi(F_2) = 0$, we can then assume that F_2 consists of one, two, or three tori, depending on how the disks D_1, D_2 are arranged.

The proof is by induction on the number of components in F_2 . Suppose F_2 is a single torus; since it is in S^3 it bounds a solid torus on at least one side. If a solid torus that it bounds lies in X_2 then the original X was a handlebody and we are done. If the solid torus that F_2 bounds lies in Y_2 then all of F also lies in that solid torus. After a torus unknotting, F_2 also bounds a solid torus in X_2 and X is a handlebody, as required.

Suppose D_1 is separating, so one of the components T of F_1 is a torus. If T bounds a solid torus in X_1 then we may as well have used the meridian of that torus for D_1 . This switch would lower the number of components of F_2 , so we can invoke the inductive hypothesis. If, on the other hand, T bounds a solid torus in Y_1 then all of F also lies in that solid torus. After perhaps a torus unknotting, T bounds a solid torus in X_1 as well, and again we could replace D_1 by a meridian of that torus and invoke the inductive hypothesis.

The same argument applies if D_2 is separating – just switch the order of D_1 and D_2 . So the only remaining possibility is that D_1 and D_2 are both non-separating but together they separate F , so F_2 consists of two tori. If either bounds a solid torus in X_2 then a meridian for that solid torus could be substituted for D_2 , thereby lowering the number of components of F_2 and completing the inductive step. If neither bounds a solid torus in X_2 then the solid torus W in S^3 that one of them does bound must contain the other torus, and so in fact W contains all of F . After a torus unknotting, ∂W will, on its other side, also bound a solid torus $W' \subset X_2$. After that reimbedding we can replace D_2 by a meridian of W' and invoke the inductive hypothesis. \square

Lemma 7.3. *If $D_1 \subset X$ and $D_2 \subset Y_1$ are both separating, and $D_2 \subset Y \subsetneq Y_1$ (i.e. D_2 does not pass through the 1-handle dual to D_1) then F can be straightened.*

Proof. Since the interiors of both D_1, D_2 are disjoint from F , their order can be rearranged, so there is symmetry between the two. Since D_1 is separating there is a torus component $T_1 \subset F_1$ and T_1 bounds a component W_1 of X_1 whose interior is disjoint from F_2 . After perhaps a torus unknotting of its complement we may as well assume that W_1 is a solid torus. Eventually some compression D_i will compress T_1 to a sphere; if $D_i \subset W_1$ then we could have done D_i before D_1 , making D_1 (coplanar to D_i) redundant, and thereby reduced the number of compressions. Thus we may as well assume $D_i \subset Y_{i-1}$, so W_1 is an unknotted solid torus. Similarly, the torus component $T_2 \subset F_2$ not incident to D_1 bounds a solid unknotted torus $W_2 \subset Y_2$. See Figure 8.

Assume, with no loss of generality, that $D_3 \subset Y_2$. We have already seen that D_3 cannot compress T_2 , else the number of compressions could be reduced. It follows that D_3 either compresses T_1 or it compresses the third torus T_3 created from F by the compressions D_1, D_2 . Either case implies that the component $U_+ = Y_2 - W_2$ of Y_2 is reducible. Apply tube straightening to $U = U_+ \cap Y$ so that the tube dual to D_1 passes through the reducing sphere of U_+ once. After the straightening, the disk D_i that eventually compresses T_1 into Y_i lies in Y , so it can be the first compression, making D_1 redundant. \square

Lemma 7.4. *If $D_1 \subset X$ and $D_2 \subset Y_1$ are both separating and if D_3 or both D_4 and D_5 are towards the X side, then F can be straightened.*

Proof. In this case there is not necessarily symmetry between D_1 and D_2 , but the argument of Lemma 7.3 still applies if the compression disk for T_3 lies on the X -side rather than the Y -side or if T_2 compresses before T_3 . So that is what we now verify: If D_3 lies on the X -side it compresses T_2 or T_3 into X_2 and we are done. On the other hand, if D_4 and D_5 both lie on the X -side then since T_1 compresses on the Y -side, one of D_4 or D_5 is the compression disk for T_3 . \square




Figure 8

Lemma 7.5. *If $D_1 \subset X$ is separating, and Y is ∂ -reducible, then either $D_2 \subset Y \subsetneq Y_1$ or F can be straightened.*

Proof. Consider the torus component T_1 of F_1 and the component $W_1 \subset X_1$ it bounds. As noted above, the interior of W_1 is disjoint from F_1 and so, perhaps after a torus unknotting, we can assume that W_1 is a solid torus. Moreover, the disk D_i that eventually compresses T_1 lies on the Y -side and not in W_1 . Hence W_1 is an unknotted solid torus.

Following Lemma 7.2 we may as well assume that $D_2 \subset Y_1$. By assumption Y is ∂ -reducible and, after attaching a 2-handle (a neighborhood of D_1) to get Y_1 the resulting manifold still is ∂ -reducible, via D_2 . It follows from the Jaco handle addition lemma [Ja] that there is a ∂ -reducing disk $J \subset Y$ for Y whose boundary is disjoint from ∂D_1 . Take J to be non-separating, if this is possible.

If ∂J lies on T_1 then it is parallel to the disk D_i that eventually compresses T_1 so we may as well do that compression before D_1 , making D_1 redundant and so reducing the number of compressions. So we henceforth suppose ∂J lies on the other, genus 2 component of $F - \partial D_1$, and so lies on $\partial F_1 - T_1$.

If ∂J is inessential in F_1 then it is parallel to ∂D_1 in F . Thus ∂D_1 also bounds the disk $J \subset Y$ and the union of the two gives a reducing sphere for Y_1 that intersects the 1-handle dual to D_1 in a single point. It follows that the longitude ∂D_i of T_1 bounds a disk in Y and we are done as before.

If ∂J is essential in F_1 compress Y along J to get Y_J and consider the component

$W_J \subset Y_J$ (in fact all of Y_J if J is non-separating) such that $\partial D_1 \subset \partial W_J$ and ∂W_J has genus 2. The manifold W_J^+ obtained by attaching $\eta(D_1)$ to W_J has boundary the union of two tori. W_J^+ can also be viewed as a component of $Y_1 - \eta(J)$.

We claim that either $D_2 \subset Y$ or W_J^+ is reducible. This is obvious if there is a disk component of $D_2 - J$ that cannot be removed by an isotopy or if $\partial D_2 \cap J = \emptyset$. The alternative then is that $D_2 \cap J$ is a non-empty collection of arcs. Consider an outermost disk D' cut off from D_2 by J . We may as well assume $D' \subset W_J^+$. For if it's not then J is separating, D' is a meridian of the other complementary component (a solid torus) and we may as well have used D' for J , thereby eliminating the case that J is separating. With then $D' \subset W_J^+$, compress the torus boundary component of ∂W_J^+ along D' to get a reducing sphere for W_J^+ .

Finally, if W_J^+ is reducible, apply tube straightening to the surface ∂W_J , replacing the handle dual to D_1 by a handle intersecting the reducing sphere once, allowing the same conclusion as before. \square

Corollary 7.6. *If $D_1 \subset X$ and $D_2 \subset Y_1$ are both separating, and Y is ∂ -reducible, then F can be straightened.*

Proof. Combine Lemmas 7.5 and 7.3. \square

Corollary 7.7. *If $D_1 \subset X$ is separating then either*

- *at least two of the three non-separating compressing disks are on the Y -side and, if D_2 is separating, Y is ∂ -irreducible, or*
- *F can be straightened.*

Proof. Following Lemmas 7.2, 7.4, and 7.5 we may as well assume that $D_2 \subset Y \subsetneq Y_1$ and D_2 is non-separating. As before, let T_1 be the torus component of F_1 bounding a component W_1 of X_1 whose interior is disjoint from F_1 .

If the compressing disk D_i that eventually compresses T_1 lies on the X -side, then $D_i \subset W_1$ (and W_1 is a solid torus). No earlier compressing disk can be incident to T_1 so in fact D_i could have been done before D_1 , making the latter redundant. This reduces the number of compressions.

If, on the other hand, D_i lies on the Y -side then both D_i and D_2 are non-separating disks lying on the Y -side, as required. \square

Lemma 7.8. *If $D_1 \subset X$ is non-separating, then either all succeeding disks D_2, D_3 are non-separating or F can be straightened.*

Proof. Since D_1 is non-separating, F_1 is a genus 2 surface. Its complementary component Y_1 contains all of Y . If D_2 is also non-separating then F_3 is a torus, for which any compressing disk is non-separating, giving the result.

So suppose D_2 is separating. Following Lemma 7.2 we may as well assume $D_2 \subset Y_1$, so ∂Y_2 consists of two tori. Hence D_3 is non-separating and compresses one of the tori. If $D_3 \subset Y_2$ then D_3 could have been done before D_2 , making D_2 redundant. If $D_3 \subset X_2$ then the result of the compression is a sphere which could have been viewed as a reducing sphere for Y_2 . Apply tube straightening, using Y_1 for N . After that reimbedding, $D_3 \subset X_1 \subsetneq X_2$ does not pass through D_2 so the compression along D_3 could be done before the compression along D_2 , making D_2 redundant. \square

We note in passing, though the fact will not be used, that if there are non-separating compressing disks on both sides, they may be taken to be disjoint:

Proposition 7.9. *If $D_1 \subset X$ is non-separating, and Y has a non-separating ∂ -reducing disk E , then either $\partial D_1 \cap \partial E = \emptyset$ or F can be straightened.*

Proof. Following Lemma 7.2 we may as well assume that $D_2 \subset Y_1$. By assumption Y is ∂ -reducible and, after attaching a 2-handle (a neighborhood of D_1) to get Y_1 the resulting manifold still ∂ -reducible, via D_2 . It follows from the Jaco handle addition lemma [Ja] that there is a ∂ -reducing disk $J \subset Y$ for Y whose boundary is disjoint from ∂D_1 . Take J to be non-separating, if this is possible.

Suppose first that ∂J is inessential in F_1 . Then the disk it bounds in F_1 contains both copies of D_1 resulting from the compression of F along D_1 . Put another way, J cuts off a component W_J from Y that has torus boundary and whose interior is disjoint from F . Following, perhaps, a torus unknotting, we may assume that W_J is a solid torus. A standard innermost disk, outermost arc argument between J and D_2 ensures that they can be taken to be disjoint, so D_2 compresses the other, genus 2 component of $Y - \eta(J)$. This compression, together with the compression via the meridian of W_J , ∂ -reduces Y to one or two components, each with a torus boundary. After perhaps some further torus unknottings, Y becomes then a handlebody, as required.

So suppose henceforth that ∂J is essential in F_1 . Compress Y along J to get Y_J and consider the component $W_J \subset Y_J$ (in fact all of Y_J if J is non-separating) such that $\partial D_1 \subset \partial W_J$ and ∂W_J has genus 2. The manifold W_J^+ obtained by attaching $\eta(D_1)$ to W_J has boundary a torus. W_J^+ can also be viewed as a component of $Y_1 - \eta(J)$.

Consider an outermost disk E' of E cut off by J , or let $E' = E$ if E is disjoint from J . We may as well assume that E' lies in W_J , since if J is separating and E' lies in the other component, we should have taken E' for J . If E' is inessential in W_J then $E' = E$ is parallel to J (since E is non-separating), so $\partial D_1 \cap \partial E = \emptyset$ and we are done. If E' is essential in W_J then each component of $Y - (\eta(J) \cup \eta(E'))$ has interior disjoint from F and is bounded by a torus. Following some torus unknottings we can take them to be solid tori. In that case Y is a handlebody, as required. \square

8. The genus 3 Schoenflies Conjecture

We now apply the results of the previous sections to complete the proof of the genus 3 Schoenflies Conjecture.

Theorem 8.1. *Suppose M is a genus 3 rectified critical level embedding of S^3 in S^4 . Then after a series of reimbeddings of M via other genus 3 rectified critical level embeddings, each of which changes at most one of the complementary components of M , one of those complementary components is B^4*

Proof. We assume that any possible genus 3 rectified critical level reimbedding of M that preserves at least one complementary component and simultaneously decreases the number of handles has been done. If any further such a sequence of reimbeddings via Lemmas 6.2 or 5.5 (reimbeddings that do not raise the number of handles) results in X_0^* (resp. Y_0^*) becoming a handlebody, then X (resp. Y) is B^4 , via Proposition 5.3. So we assume no such further sequence exists and use the results of the previous section to see if there are other options. With no loss of generality, assume the cocore D_1 of the highest 1-handle lies in X_0^* and let E_1 denote the core of the lowest 2-handle.

Following Lemma 7.2 we can assume that the cocore D_2 of the previous 1-handle lies on the Y -side and the cocore E_2 of the next 1-handle lies on the side opposite the one on which E_1 lies.

Claim. Perhaps after rearranging the ordering of the handles, at least one of D_1 and E_1 is non-separating.

Suppose D_1 and E_1 are both separating. If both D_1 and E_1 lie in X_0^* , then it follows from Corollary 7.7 that at least two of the non-separating cores of the 2-handles and at least two of the non-separating cocores of the 1-handles all lie on the Y -side of the surfaces to which they are attached. So both Y_0^+ and Y_0^- are 4-dimensional handlebodies of genus at least 2. But the Mayer–Vietoris sequence for Y_0^+ , Y_0^- glued along Y_0^* then contradicts $H_*(Y) \cong H_*(B^4)$, cf. the proof of Lemma 3.2.

On the other hand, if $D_1 \subset X_0^*$ and $E_1 \subset Y_0^*$ then, following Corollary 7.6 and Lemma 7.5, D_2 is non-separating and lies in Y_0^* . Then interchange D_1 and D_2 , using the 1-handle dual to D_2 as the highest 1-handle. The new arrangement establishes the claim.

Following the claim, we can, with no loss, assume that E_1 is non-separating. Then according to Lemma 7.8, all of the cores of 2-handles are non-separating, so each surface F_i at or above height $t = 0$ are connected. Hence there is at most one 3-handle in M and, passing this 3-handle over the north pole if necessary, this guarantees that each of X and Y have induced handle structures without 3-handles. Following the comments preceding Lemma 4.4, the sum of the genera of the 4-dimensional handlebodies X_0^- and Y_0^- is 3, so one of them, say X_0^- , has genus ≤ 1 . X is then obtained from the genus 0 or 1 handlebody X_0^- by attaching some 2-handles but no

3-handles. Moreover, ∂X is a sphere. If $(X_0^-) = 0$ (resp. $\text{genus}(X_0^-) = 1$) then no (resp. one) 2-handle must be attached, to ensure that $H_1(X) = H_2(X) = 0$. In the first case, since no 2-handles (hence no handles at all) are attached, $X \cong X_0^- \cong B^4$. In the second case, one 2-handle is attached to $X_0^- \cong S^1 \times D^3$ and so $X \cong B^4$, by Corollary 2.2. \square

Corollary 8.2. *Each complementary component of a genus 3 rectified critical level embedding of S^3 in S^4 is a 4-ball.*

Proof. Let M be a genus 3 rectified critical level embedding of S^3 . Following Theorem 8.1 there is a sequence of such embeddings

$$M = M_0, M_1, M_2, \dots, M_n$$

so that one of the complementary components of M_n is B^4 and, furthermore, for each $i = 0, \dots, n-1$, one of the complementary components of M_i is homeomorphic to a complementary component of M_{i+1} . The argument is by induction on n , exploiting the fact that the complement of B^4 in S^4 is B^4 .

Since one complementary component of M_n is B^4 , both complementary components are. If $n = 0$ we are done. For $n \geq 1$, note that since one complementary component of M_{n-1} is homeomorphic to a complementary component of M_n , that complementary component is B^4 . This completes the inductive step. \square

References

- [BC] J. Birman and R. Craggs, The μ -invariant of 3-manifolds and certain structural properties of the group of homeomorphisms of a closed oriented 2-manifold. *Trans. Amer. Math. Soc.* **237** (1978), 283–309. [Zbl 0383.57006](#) [MR 0482765](#)
- [Fo] R. Fox, On the imbedding of polyhedra in 3-space. *Ann. of Math.* **49** (1948), 462–470. [Zbl 0032.12502](#) [MR 0026326](#)
- [Ga1] D. Gabai, Foliations and the topology of 3-manifolds. II. *J. Differential Geom.* **26** (1987), 461–478. [Zbl 0627.57012](#) [MR 0910017](#)
- [Ga2] D. Gabai, Foliations and the topology of 3-manifolds. III. *J. Differential Geom.* **26** (1987), 479–536. [Zbl 0639.57008](#) [MR 0910018](#)
- [Ga3] D. Gabai, Genus is superadditive under band connected sum. *Topology* **26** (1987), 209–210. [Zbl 0621.57004](#) [MR 0895573](#)
- [Ja] W. Jaco, Adding a 2-handle to a 3-manifold: an application of property R . *Proc. Amer. Math. Soc.* **92** (1984), 288–292. [Zbl 0564.57009](#) [MR 0754723](#)
- [KL] C. Kearton and W. B. R. Lickorish, Piecewise linear critical levels and collapsing. *Trans. Amer. Math. Soc.* **170** (1972), 415–424. [Zbl 0248.57007](#) [MR 0310899](#)
- [Ki] R. Kirby, A calculus for framed links in S^3 . *Invent. Math.* **45** (1978), 35–56. [Zbl 0377.55001](#) [MR 0467753](#)

- [LP] F. Laudenbach, and V. Poénaru, A note on 4-dimensional handlebodies. *Bull. Soc. Math. France* **100** (1972), 337–344. [Zbl 0242.57015](#) [MR 0317343](#)
- [Sch1] M. Scharlemann, The four-dimensional Schoenflies Conjecture is true for genus two imbeddings. *Topology* **23** (1984), 211–217. [Zbl 0543.57011](#) [MR 0744851](#)
- [Sch2] M. Scharlemann, Producing reducible 3-manifolds by surgery on a knot. *Topology* **29** (1990), 481–500. [Zbl 0727.57015](#) [MR 1071370](#)
- [Th] A. Thompson, Property P for the band-connect sum of two knots. *Topology* **26** (1987), 205–207. [Zbl 0628.57005](#) [MR 0895572](#)
- [Wa] F. Waldhausen, Heegaard-Zerlegungen der 3-Sphäre. *Topology* **7** (1968), 195–203. [Zbl 0157.54501](#) [MR 0227992](#)

Received August 9, 2006

Martin Scharlemann, Mathematics Department, University of California, Santa Barbara, CA, U.S.A.

E-mail: mgscharl@math.ucsb.edu