

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 44 (1989)  
**Heft:** 5

**Artikel:** An extension of the isoperimetric inequality on the sphere  
**Autor:** Bollobás, Béla  
**DOI:** <https://doi.org/10.5169/seals-41619>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

monotonicity theorems. The use of these in the proof of l'Hôpital's rule was made by Lettenmeyer [4]. Since monotonicity theorems are known to hold for Dini derivatives, it is clear from our exposition that the right-hand derivatives can be replaced in Theorem 1–2 without affecting their validity by Dini derivatives. The following counterexample:

$$f(x) = x + \sin x \cos x, \quad g(x) = f(x) e^{\sin x}$$

$\lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)} = 0$  and no limit for  $\frac{f(x)}{g(x)}$  as  $x \rightarrow \infty$  was given already in 1879 by O. Stolz [6], who also showed that Theorem 3 (with ordinary rather than one-sided derivatives) can be deduced from Theorem 2. A simple proof based on the Newton-Leibniz formula was given by Boas [2] but one may conjecture that the method was already known to Huntington [3].

R. Vyborny and R. Nester  
University of Queensland, St. Lucia (Australia)

#### REFERENCES

- 1 Boas R. P.: Counter-examples to L'Hôpital's Rule. *American Math. Monthly* 93, 644–645 (1986).
- 2 Boas R. P.: L'Hôpital's rule without mean-value theorems. *American Math. Monthly* 76, 1051–1053 (1969).
- 3 Huntington E. U.: Simplified proof of L'Hôpital's theorem on indeterminate forms. *Bulletin of Amer. Math. Soc.* 29, 207 (1923).
- 4 Lettenmeyer F.: Über die sogenannte Hospitalsche Regel. *J. Reine Angew. Math* 174, 246–247 (1936).
- 5 Miller A. D. and Vyborny R.: Some Remarks on Functions with One-sided Derivatives. *American Math. Monthly* 93, 471–475 (1986).
- 6 Stolz O.: Über die Grenzwerte der Quotienten. *Math. Ann.* 15, 556–559 (1879).

## An extension of the isoperimetric inequality on the sphere

We shall consider the  $n$ -dimensional sphere  $S^n = \{x \in \mathbb{R}^{n+1} : |x| = 1\}$ , endowed with the spherical distance function  $d(x, y)$  and the (normalized) Lebesgue measure  $\mu$ . For  $x \in S^n$  and  $0 \leq \theta \leq \pi$ , the *spherical cap* of centre  $x$  and radius  $\theta$  is  $C(x, \theta) = \{y \in S^n : d(x, y) \leq \theta\}$ . It is well known that if  $A \subset S^n$  and  $\mu(A) = \mu(C)$  for some spherical cap  $C$ , then the diameter of  $A$  is at least as large as the diameter of  $C$ . This is usually considered to be a variant of the isoperimetric inequality on the sphere  $S^n$ ; it is, in fact, an immediate consequence of the isoperimetric inequality. Our aim is to extend this inequality and thereby answer a question raised by Paul Erdős [4].

For  $k \geq 2$ , define the  $k$ -diameter  $d_k(A)$  of a set  $A$  in a metric space by

$$d_k(A) = \sup \left\{ \min_{1 \leq i < j \leq k} d(x_i, x_j) : x_1, \dots, x_k \in A \right\}.$$

Thus  $d_k(A) \leq d$  if and only if  $A$  does not contain  $k$  points, any two of which are at distance greater than  $d$ ; in particular,  $d_2(A)$  is precisely the diameter of  $A$ . We shall show that if  $A \subset S^n$  and  $0 < \mu(A) = \mu(C)$  for some spherical cap  $C$  then  $d_k(A) \geq d_k(C)$  for every  $k \geq 2$ .

The proof we shall give makes use of compression operators and closely follows Benyamini's [2] proof of the classical isoperimetric inequality on the sphere. Benyamini's proof, in turn, was inspired by Baernstein and Taylor [1]. In spirit, the compression operators on the sphere are very close to the compression operators frequently used in the study of set systems in combinatorics (see [3; Chapters 16 and 17 and [4]).

Let  $A$  be a subset of  $S^n$  and  $z \in S^n$ . The *compression*  $\gamma_z(A)$  of  $A$  in the direction of  $z$  is defined as follows. For  $x \in S^n$ , let  $x^+ = x - \langle x, z \rangle z + |\langle x, z \rangle| z$  and  $x^- = x - \langle x, z \rangle z - |\langle x, z \rangle| z$ , where  $\langle \cdot, \cdot \rangle$  denotes the inner product in  $\mathbb{R}^{n+1}$ . Thus the line through  $x$ , in the direction of  $z$ , meets  $S^n$  precisely in  $x^+$  and  $x^-$ , where  $\langle x^+, z \rangle = -\langle x^-, z \rangle \geq 0$ ; furthermore,  $x^+ = x^-$  if and only if  $\langle x, z \rangle = 0$ .

The compression operator  $\gamma_z$  pushes the points of  $A \cap \{x^+, x^-\}$  towards  $x^+$ : if  $A \cap \{x^+, x^-\} = \{x^-\}$  then

$$\gamma_z(A) \cap \{x^+, x^-\} = \{x^+\}$$

and if  $A \cap \{x^+, x^-\} \neq \{x^-\}$  then

$$\gamma_z(A) \cap \{x^+, x^-\} = A \cap \{x^+, x^-\}.$$

It is trivial that if  $A$  is measurable then so is  $\gamma_z(A)$  and we have  $\mu(\gamma_z(A)) = \mu(A)$ ; furthermore, if  $A$  is closed, so is  $\gamma_z(A)$ . The compression operators map caps into caps:  $\gamma_z(C(x, \theta)) = C(x^+, \theta)$ ; furthermore, for any two measurable sets  $A$  and  $B$ ,

$$\mu(A \cap B) \leq \mu(\gamma_z(A) \cap \gamma_z(B)). \quad (1)$$

Thus  $\gamma_z$  not only compresses as much of a set  $A$  into the hemisphere  $\{x^+ : x \in S^n\} = \{x \in S^n : \langle x, z \rangle \geq 0\}$  as possible, but it also compresses sets closer to each other. In this note, the most important property of compression operators is that they do not increase the  $k$ -diameter.

**Lemma 1.** If  $A \subset S^n$ ,  $z \in S^n$  and  $k \geq 2$  then  $d_k(\gamma_z(A)) \leq d_k(A)$ .

*Proof.* It suffices to show that if  $d_k(\gamma_z(A)) > d$  then  $d_k(A) \geq d$ . Let then  $d_k(\gamma_z(A)) > d$ . Then there is a set  $X = \{x_1, \dots, x_k\} \subset \gamma_z(A)$  with  $d(x_i, x_j) \geq d$  for  $i \neq j$ . We claim that  $A$  contains a  $k$ -subset  $X'$  with minimal distance at least  $d$ , so  $d_k(A) \geq d$ .

In proving this claim we may assume that  $x_1, \dots, x_l$  are the points of  $X = \{x_1, \dots, x_k\}$  that do not belong to  $A$ . Then  $x_i = x_i^+$  for  $1 \leq i \leq l$  and  $X' = \{x_1^-, \dots, x_l^-, x_{l+1}, \dots, x_k\} \subset A$ .

Furthermore, the minimal distance in this  $k$ -subset  $X'$  of  $A$  is at least  $d$ . Indeed, for  $1 \leq i < j \leq l$ ,  $d(x_i^-, x_j^-) = d(x_i^+, x_j^+) \geq d$  since  $x_i^+, x_j^+ \in X$ , and for  $l+1 \leq i < j \leq k$  we have  $d(x_i, x_j) \geq d$  since  $x_i, x_j \in X$ . Let now  $1 \leq i \leq l$  and  $l+1 \leq j \leq k$ . If  $x_j = x_j^-$  then  $d(x_i^-, x_j^-) = d(x_i^-, x_j^-) = d(x_i^+, x_j^+) \geq d$  since  $x_i^+, x_j^+ \in X$ . Finally, if  $x_j = x_j^+$  then  $d(x_i^-, x_j) = d(x_i^-, x_j^+) \geq d(x_i^+, x_j^+) \geq d$  since  $x_i^+, x_j^+ \in X$ .  $\square$

Loosely speaking, our aim is to show that if  $A$  is a closed subset of  $S^n$  then  $A$  can gradually be transformed into a spherical cap of measure at least  $\mu(A)$  and  $k$ -diameter at most  $d_k(A)$ . Lemma 1 tells us that  $A$  can be transformed into  $\gamma_z(A)$  for every  $z \in S^n$ . The next lemma, which is essentially trivial, shows that we can take limits in the Hausdorff metric: the  $k$ -diameter is continuous in this metric and, in fact, every Borel measure on  $S^n$  is upper semi-continuous. Let  $H$  be the metric space of closed non-empty subsets of  $S^n$  with the Hausdorff metric  $d(A, B) = \sup \{d(a, B), d(b, A) : a \in A, b \in B\}$ . Since  $S^n$  is compact,  $H$  is also a compact metric space.

**Lemma 2.** Let  $\nu$  be a Borel measure on  $S^n$  and let  $A, A_1, A_2, \dots \in H, A_s \rightarrow A$ . Then

$$\nu(A) \geq \lim_{s \rightarrow \infty} \nu(A_s) \quad \text{and} \quad d_k(A) = \lim_{s \rightarrow \infty} d_k(A_s).$$

*Proof.* (i) Given  $\varepsilon > 0$ , let  $\delta > 0$  be such that  $\nu(A_\delta) < \nu(A) + \varepsilon$ , where  $A_\delta = \{x \in S^n : d(x, A) < \delta\}$ . If  $s$  is large enough then  $A_s \subset A_\delta$  so  $\nu(A_s) < \nu(A) + \varepsilon$ , showing that  $\nu$  is upper semi-continuous.

(ii) Suppose  $d(A, B) < \delta$  where  $A, B \in H$ , and  $x_1, \dots, x_k \in A$ . Then for each  $x_i$  there is a  $y_i \in B$  such that  $d(x_i, y_i) < \delta$ . Clearly  $d(y_i, y_j) > d(x_i, x_j) - 2\delta$  so  $d_k(B) \geq d_k(A) - 2\delta$ . Interchanging  $A$  and  $B$  we see that  $d_k(A) \geq d_k(B) - 2\delta$ . Hence, given  $\varepsilon > 0$ , if  $s$  is large enough to guarantee that  $d(A_s, A) < \varepsilon/2$  then we have  $|d_k(A_s) - d_k(A)| \leq \varepsilon$ .  $\square$

We are ready to prove the main result of this note. As usual, we shall write  $\mu^*$  for the outer measure defined by  $\mu$ .

**Theorem 3.** Let  $A$  be a non-empty subset of  $S^n$  and let  $C$  be a cap of measure  $\mu^*(A)$ . Then  $d_k(A) \geq d_k(C)$  for every  $k \geq 2$ .

*Proof.* The assertion is trivial if  $\mu^*(A) = 0$  or  $\mu^*(A) = \mu(S^n)$ . Furthermore, since  $d_k(A) = d_k(\bar{A})$ , we may assume that  $A$  is a closed set of measure  $m$ ,  $0 < m < \mu(S^n)$ .

Let  $K$  be the minimal closed subset of  $H$  containing  $A$  and closed under  $\gamma_z$  for every  $z \in S^n$ . By Lemmas 1 and 2, every set in  $K$  has measure at least  $m$  and  $k$ -diameter at most  $d_k(A)$ . For a Borel subset  $M$  of  $S^n$ , define  $\nu(M) = \mu(M \cap C)$ , where  $C$  is our spherical cap of measure  $m$ . Then  $\nu$  is a Borel measure on  $S^n$ ; by Lemma 2, this measure  $\nu$  is upper semi-continuous so its supremum on  $K$  is attained on some set  $M \in K$ . To complete the proof, we shall show that  $M$  contains the cap  $C$ .

Suppose that this is not the case. Then there is a cap  $= C(x, \theta)$ ,  $\theta > 0$ , such that  $D \subset C \setminus M$ . Since  $\mu(M) \geq \mu(C)$ , this implies that  $\mu(M \setminus C) > 0$  so there is a cap  $E = C(y, \mu)$ ,  $0 < \mu \leq \theta$ , such that  $E \cap C = \emptyset$  and  $\mu(M \cap E) > 0$ . By replacing  $\theta$  by  $\mu$ , we may assume that  $\mu = \theta$ .

Let  $z = (x - y)/\|x - y\|$ . Then  $\gamma_z(E) = D$ ,  $\gamma_z(C) = C$  and  $\gamma_z(C \setminus D) = C \setminus D$ . Hence, by (1),

$$\begin{aligned}\mu(\gamma_z(M) \cap C) &= \mu(\gamma_z(M) \cap (C \setminus D)) + \mu(\gamma_z(M) \cap D) \\ &\geq \mu(M \cap (C \setminus D)) + \mu(M \cap E) = \mu(M \cap C) + \mu(M \cap E) > \mu(M \cap C).\end{aligned}$$

Since  $\gamma_z(M) \in K$ , this contradicts the choice of  $M$ , so the proof is complete.  $\square$

Let us remark that a slight variant of the proof above gives the following assertion. Let  $K$  be a non-empty closed subset of  $H$  which is also closed under the operators  $\gamma_z$ , i.e. which is such that  $\gamma_z(A) \in K$  for all  $A \in K$  and  $z \in S^n$ . Then  $K$  contains all caps of measure  $m = \sup \{\mu(A) : A \in K\}$ .

Also, it is easily seen that the proof above implies various extensions of Theorem 3. For example, given finite sets  $X, Y \subset S^n$  with  $|X| = |Y|$ , let us write  $X \leq Y$  if for every  $d > 0$ , the number of pairs in  $X$  at distance at least  $d$  is not more than the number of pairs in  $Y$  at distance at least  $d$ . Furthermore, for sets  $A, B \subset S^n$ , let us write  $A \leq B$  for the assertion that for every finite set  $X \subset A$  there is a finite set  $Y \subset B$  with  $|Y| = |X|$  and  $X \leq Y$ . Then the following assertion holds. Let  $A$  be a non-empty closed subset of  $S^n$  and let  $C$  be a cap of measure  $\mu(A)$ . Then  $C \leq A$ .

Béla Bollobás

Department of Pure Mathematics and Mathematical Statistics  
University of Cambridge, England

#### REFERENCES

- 1 Baernstein A., Taylor B. A.: Spherical rearrangements, subharmonic functions and \*-functions in  $n$ -space, *Duke Math. J.* **43**, 245–268 (1976).
- 2 Benyamini Y.: Two point symmetrization, the isoperimetric inequality on the sphere and some applications, *Longhorn Notes*, The University of Texas, Texas Functional Analysis Seminar, pp. 53–76, 1983–1984.
- 3 Bollobás B.: *Combinatorics*, Cambridge University Press, xii + 177, Cambridge, England, 1986.
- 4 Frankl P.: The shifting technique in extremal set theory, in «*Surveys in Combinatorics 1987*» (Whitehead C., ed.), LMS Lecture Note Series 123, Cambridge University Press, pp. 81–110, Cambridge, 1987.

## Winch curves

A taut rope connects a point in the origin of a rectangular coordinate system with a point in  $R(a, 0)$ . If the latter starts moving along the line  $x = a$ , it will trail the point in the origin. For each point  $P$  of the curve that is created in this way we have  $PQ = a$ , where  $Q$  is the intersection of the tangent to the curve in  $P$  with the line  $x = a$ . This curve, known as the tractrix, is represented by an equation that can be found as follows.

In the rectangular triangle  $PSQ$  (see fig. 1) we have

$$PQ = a, \quad PS = a - x, \quad SQ = (a - x) dy/dx.$$