

Stein manifolds

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **14 (1968)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **22.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

usual coboundary map $\delta : C^l(\mathfrak{U}, S) \rightarrow C^{l+1}(\mathfrak{U}, S)$ which makes the system a complex. We put $Z^l(\mathfrak{U}, S) = \text{Ker } \delta \subset C^l(\mathfrak{U}, S)$ and $B^l(\mathfrak{U}, S) = \delta(C^{l-1}(\mathfrak{U}, S))$. The l -th cohomology group $H^l(\mathfrak{U}, S)$ with respect to the open covering \mathfrak{U} is $Z^l(\mathfrak{U}, S)/B^l(\mathfrak{U}, S)$. An open covering $\mathfrak{B} = \{V_\nu\}_{\nu \in N}$ is finer than an open covering $\mathfrak{U} = \{U_i\}_{i \in J}$ if there exists an index map $\tau : N \rightarrow J$ such that $V_\nu \subset U_{\tau(\nu)}$ for $\nu \in N$. It follows that an element of $\Gamma(U_{\tau(\nu_0)} \dots \tau(\nu_l), S)$ can be restricted to a continuous cross-section over $V_{\nu_0} \dots \nu_l$. In this way we get a map $\tau^* : C^l(\mathfrak{U}, S) \rightarrow C^l(\mathfrak{B}, S)$. The following diagram is commutative:

$$\begin{array}{ccc} & \tau^* & \\ C^l(\mathfrak{U}, S) & \rightarrow & C^l(\mathfrak{B}, S) \\ \delta \downarrow & & \delta \downarrow \\ & \tau^* & \\ C^{l+1}(\mathfrak{U}, S) & \rightarrow & C^{l+1}(\mathfrak{B}, S) \end{array}$$

It follows that we have a map $\tau^* : Z^l(\mathfrak{U}, S) \rightarrow Z^l(\mathfrak{B}, S)$. Let us put $Z^l(X, S) = \bigcup_{\mathfrak{U}} Z^l(\mathfrak{U}, S)$, where \mathfrak{U} runs over all open coverings of X . In

$Z^l(X, S)$ we can introduce an equivalence relation \approx as follows: Let $\xi_1 \in Z^l(\mathfrak{U}, S)$ and $\xi_2 \in Z^l(\mathfrak{U}_1, S)$. We put $\xi_1 \approx \xi_2$ iff there exists \mathfrak{U}_2 such that \mathfrak{U}_2 is finer than \mathfrak{U} and \mathfrak{U}_1 and $\xi_1|_{\mathfrak{U}_2} - \xi_2|_{\mathfrak{U}_2} \in B^l(\mathfrak{U}_2, S)$. Here we have put $\xi_\nu|_{\mathfrak{U}_2} = \tau_\nu^*(\xi_\nu)$ where τ_ν^* comes from an index map $\tau_\nu : \mathfrak{U}_2 \rightarrow \mathfrak{U}, \mathfrak{U}_1$. It is easy to check that the equivalence relation defined on $Z^l(X, S)$ is independent of the index maps. Now $H^l(X, S)$ is the set of equivalence classes in $Z^l(X, S)$. Because $C^l(\mathfrak{U}, S)$ is a module over the ring $I(X)$ of holomorphic functions on X it follows that $H^l(\mathfrak{U}, S)$ and $H^l(X, S)$ are modules over $I(X)$. We have a natural homomorphism $H^l(\mathfrak{U}, S) \rightarrow H^l(X, S)$. Let now $X' \subset X$ be an open subset. Then X' is a complex analytic manifold. We put $S' = S|_{X'}$ and $\mathfrak{U}' = \mathfrak{U} \cap X' = \{U_i \cap X'\}$ and obtain an open covering of X' . The restriction of cross-sections gives a homomorphism $\gamma : C^l(\mathfrak{U}, S) \rightarrow C^l(\mathfrak{U}', S')$ which commutes with δ and any index map τ . Thus we obtain restriction homomorphisms: $H^l(\mathfrak{U}, S) \rightarrow H^l(\mathfrak{U}', S')$ and $H^l(X, S) \rightarrow H^l(X', S')$.

STEIN MANIFOLDS

A complex analytic manifold X is a Stein manifold if: 1) X is holomorphically convex, i.e. if $D = (x_\nu)_1^\infty$ is an infinite discrete set, then there exists $f \in I(X)$ such that $|f(D)| = \sup_\nu |f(x_\nu)|$ is infinite. 2) X can be

spread holomorphically, i.e. for any $x \in X$ there exists $f_1 \dots f_N \in I(X)$ such that x is an isolated common zero of $f_1 \dots f_N$.

Let X be a complex analytic manifold. A Stein covering $\mathfrak{U} = \{U_i\}_{i \in J}$ of X is an open covering of X such that every U_i is Stein. We shall often use the following result:

Leray's Theorem: If \mathfrak{U} is a Stein covering of X then $H^l(\mathfrak{U}, S) \rightarrow H^l(X, S)$ is an isomorphism for every coherent analytic sheaf S .

The isomorphism between $H^l(\mathfrak{U}, S)$ and $H^l(X, S)$ means the following: If $\underline{\xi} \in H^l(X, S)$ there exists $\xi \in Z^l(\mathfrak{U}, S)$ such that ξ maps into $\underline{\xi}$ under the natural homomorphism $Z^l(\mathfrak{U}, S) \rightarrow H^l(X, S)$ and moreover if $\underline{\xi} \in Z^l(\mathfrak{U}, S)$ is mapped into zero in $H^l(X, S)$ there exist $\eta \in C^{l-1}(\mathfrak{U}, S)$ such that $\xi = \delta\eta$ in $C^l(\mathfrak{U}, S)$.

DIRECT IMAGES OF SHEAVES

Let X and Y be complex analytic manifolds. Let $\psi : X \rightarrow Y$ be a holomorphic map and let S be an analytic sheaf on X . Now X is fibered by the fibers $X(y) = \psi^{-1}(y)$ for $y \in Y$. Let U be an open neighborhood of a point $y \in Y$, then $V = \psi^{-1}(U)$ is an open set in X . Hence V is a complex analytic manifold and the restriction of S to V gives an analytic sheaf on V . We can now define $H^l(V, S)$. Let us put $H_y^l = \bigcup_U H^l(\psi^{-1}(U), S)$

where U runs over all open neighborhoods of y in Y . In H_y^l we introduce an equivalence relation as follows: $\xi_1 \in H^l(\psi^{-1}(U_1), S)$ and $\xi_2 \in H^l(\psi^{-1}(U_2), S)$ are equivalent iff there exists $U = U(y)$ in Y such that $U \subset U_1 \cap U_2$ and $\xi_1|_{\psi^{-1}(U)} = \xi_2|_{\psi^{-1}(U)}$ in $H^l(\psi^{-1}(U), S)$. We let $\psi_{(l)}(S)_{(y)}$ denote the set of equivalence classes in H_y^l . The equivalence class generated by $\xi \in H^l(\psi^{-1}(U), S)$ is denoted by ξ_y . The set $\psi_{(l)}(S)_{(y)}$ is called the set of germs of cohomology classes of dimension l along the fiber $X(y)$. Now $\psi_{(l)}(S)_{(y)}$ is an $\mathcal{O}_{y,Y}$ -module. For if $g_y \in \mathcal{O}_{y,Y}$ we have a representative $g \in I(U)$ for some open neighborhood U of y . Then $g \circ \psi \in I(\psi^{-1}(U))$. If $\xi_y \in \psi_{(l)}(S)_{(y)}$ and U is sufficiently small we can find a representative $\xi \in H^l(\psi^{-1}(U), S)$ for ξ_y . Then we put $g_y \cdot \xi_y = ((g \circ \psi)\xi)_y$. Now we form $\psi_{(l)}(S) = \bigcup_{y \in Y} \psi_{(l)}(S)_{(y)}$ where we introduce a sheaf topology.

A base of the open sets are $\{\xi_y : y \in U\}$ for $\xi \in H^l(\psi^{-1}(U), S)$. If $\xi \in H^l(X, S)$ then the map $y \rightarrow \xi_y$ is a cross-section in $\psi_{(l)}(S)$. We call it the direct image of ξ and denote it by $\psi_{(l)}(\xi)$. The sheaf $\psi_{(l)}(S)$ is the direct