

6. The exact sequence

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **23 (1977)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **19.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

and so $k^{q+1}s \simeq k^{q+1}t$ over X^n . Hence, by induction, we obtain (5.1). Of course the value of r can often be improved in particular cases.

COROLLARY (5.2). *Let E be a fibre bundle over X with locally compact fibre F , which admits a cross-section. Choose a cross-section and so regard E as an ex-space. Let $f: E \rightarrow E$ be an ex-map such that $g: F \rightarrow F$ is null-homotopic, where $g = f|F$. Then $f^{r+1} \simeq c$, the trivial ex-map, where $r = \text{reg}(X)$.*

To see this, take $M = M_X(E, E)$, in (5.1), and define $k: M \rightarrow M$ by post-composition with f . We take s, t to be the cross-sections $f', e': X \rightarrow M$ determined by f, e , and obtain (5.2).

Now let $\alpha, \beta \in \pi_X(E, E)$ be elements such that

- (i) $\alpha^2 = \beta^2$ and $\alpha\beta = \beta\alpha$,
- (ii) $\Phi_*\alpha = \Phi_*\beta$,

where $\Phi_*: \pi_X(E, E) \rightarrow \pi(F, F)$ is given by restriction. Suppose that $E = \Sigma E'$, for some ex-space E' , and that $\alpha = \Sigma_*\alpha', \beta = \Sigma_*\beta'$, for some $\alpha', \beta' \in \pi_X(E', E')$. Take f in (5.2) to be a representative of $\alpha - \beta$. Then f^{r+1} is a representative of $(\alpha - \beta)^{r+1} = 2^r(\alpha - \beta)\alpha^r$, and so (5.2) shows that

$$(5.3) \quad 2^r\alpha = 2^r\beta.$$

Applications will be given in §8 below.

6. THE EXACT SEQUENCE

Let X be a CW-complex with basepoint x_0 a 0-cell. Let $p: M \rightarrow X$ be a fibration with fibre $N = p^{-1}(x_0)$, and let Γ denote the function-space of cross-sections. By evaluating at x_0 we obtain a fibration $q: \Gamma \rightarrow N$. It may be noted that, under fairly general conditions, this fibration admits a cross-section if and only if the original fibration is trivial, in the sense of fibre homotopy type.

Now choose a basepoint $y_0 \in N$ so that $q^{-1}(y_0) = \Gamma_0$, the space of pointed cross-sections. Choose such a cross-section s as basepoint in $\Gamma_0 \subset \Gamma$, and consider the homotopy exact sequence of the fibration as follows:

$$\dots \rightarrow \pi_{r+1}(N) \xrightarrow{\Delta} \pi_r(\Gamma_0) \xrightarrow{u_*} \pi_r(\Gamma) \xrightarrow{q_*} \pi_r(N) \rightarrow \dots$$

Note that Γ_0 is a deformation retract of $\tilde{\Gamma}_0$, the space of pointed maps $t: X \rightarrow M$ such that $pt \simeq 1$.

In particular, suppose that $X = S^n$ ($n \geq 1$). Then s determines a homotopy equivalence $k: \Omega^n(N) \rightarrow \Gamma_0$ as follows. Consider the map $l: \Omega^n(N) \rightarrow \tilde{\Gamma}_0$ which transforms each pointed map $f: S^n \rightarrow N$ into the track sum $s + jf$, where $j: N \subset M$. Then l is a homotopy equivalence and k is obtained by composing l with a deformation retraction $\tilde{\Gamma}_0 \rightarrow \Gamma$. In this case, therefore, we can transform our exact sequence into

$$\dots \rightarrow \pi_{r+1}(N) \xrightarrow{D} \pi_{r+n}(N) \xrightarrow{v_*} \pi_r(\Gamma) \xrightarrow{q_*} \pi_r(N) \rightarrow \dots,$$

where $v_* = u_*k_*$ and $D = k_*^{-1} \Delta$. We refer to this as the *modified exact sequence* of the evaluation fibration. The operator D has been determined in §3 of [8]. Specifically, let $\sigma \in \pi_n(M)$ denote the class of s , and let $\alpha \in \pi_{r+1}(N)$. Then

$$(6.1) \quad j_* D\alpha = [j_* \alpha, \sigma],$$

the Whitehead product in $\pi_*(M)$.

We shall be particularly concerned with the tail end of this sequence, which reads

$$\pi_1(N) \xrightarrow{D} \pi_n(N) \xrightarrow{v_*} \pi(\Gamma) \xrightarrow{q_*} \pi(N).$$

Let $t: S^n \rightarrow M$ also be a cross-section. Suppose that we have a path λ in N from $s(x_0)$ to $t(x_0)$. We can regard λ as a vertical homotopy of s into t over $\{x_0\}$. The obstruction to extending this to a vertical homotopy over S^n is an element

$$\delta(s, t; \lambda) \in \pi_n(N).$$

If μ is another path in N from $s(x_0)$ to $t(x_0)$ the track difference $\lambda - \mu$ forms a loop in N and it is easy to check that the homotopy class $\alpha \in \pi_1(N)$ of this loop satisfies the relation

$$D\alpha = \delta(s, t; \lambda) - \delta(s, t; \mu).$$

Hence s and t are vertically homotopic if and only if the obstruction is contained in $D\pi_1(N)$.

Now let E_i ($i=1, 2$) be a sectioned bundle over X with locally compact fibre F_i . We can apply the above to the function-space bundle $M = M_X(E_1, E_2)$ with fibre $N = N(F_1, F_2)$, and obtain useful information about the ex-homotopy groups $\pi_X(\Sigma^r E_1, E_2)$ ($r=1, 2, \dots$). Details are given in [9] where the operator D is calculated, as follows, in case E_1 and E_2 are sphere-bundles over $X = S^n$.

Given a representation $\phi: SO(m) \rightarrow SO(q)$ write

$$J_\phi = J \circ \phi_*: \pi_r SO(m) \rightarrow \pi_{r+q}(S^q),$$

where J denotes the usual Hopf-Whitehead homomorphism. For example, if $q > m$ and ϕ is the inclusion then

$$(6.2) \quad J_\phi = (-1)^{m-q} \Sigma_*^{m-q} J,$$

by (3.2) of [5] (cf. [8]). If $q = 2m$ and $\phi = 1 \oplus 1$ it is easily seen that

$$(6.3) \quad J_\phi = 2(-1)^m \Sigma_*^m J.$$

Consider the function-space $N = N(S^p, S^q)$ of pointed maps $S^p \rightarrow S^q$. We identify $\pi_i(N)$ ($i=0, 1, \dots$) with $\pi_{i+p}(S^q)$ in the standard way (see [15]). Let G be a topological group and let

$$\phi: G \rightarrow SO(p), \quad \psi: G \rightarrow SO(q)$$

be representations of G . We regard S^p, S^q as pointed G -spaces using ϕ, ψ , respectively. Choose a principal G -bundle P over S^n with classifying element $\theta \in \pi_{n-1}(G)$, and take $E_1 = P_\#(S^p)$, $E_2 = P_\#(S^q)$. Then the operator D in our exact sequence is given

$$(6.4) \quad D\alpha = \alpha \circ \Sigma_*^{r+p-q+1} J_\psi \theta - J_\phi \theta \circ \Sigma_*^{n+p-q-1} \alpha,$$

where $\alpha \in \pi_{r+p+1}(S^q)$. The case $r = 1$ of this result will be needed in §8 below.

7. THE ADJOINT G -BUNDLE

Let X be any space and let P be a principal G -bundle over X . We regard P as a (right) G -space in the usual way. By a *principal automorphism* we mean an equivariant fibre-preserving map of P into itself. By the *adjoint G -bundle* we mean the sectioned bundle $Q = P_\#G$, where G acts on itself by conjugation. Note that Q is a group ex-space since G is a group G -space. We can construct Q from $G \times P$ by identifying

$$(7.1) \quad (gag^{-1}, b) \sim (a, bg) \quad (a \in G, b \in P)$$

for all $g \in G$. The group ex-structure is given by

$$\{a_1, b\} \cdot \{a_2, b\} = \{a_1 \cdot a_2, b\} \quad (a_1, a_2 \in G),$$

where $\{ \ , \ }$ denotes the equivalence class of $(\ , \)$. Every principal automorphism f of P determines a cross-section $f': X \rightarrow Q$ as follows.