

## 5. Construction of an algebraic model for the space OF SECTIONS OF A FIBER BUNDLE ([20], [18], [13]).

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **24 (1978)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.04.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

a morphism of  $A$  in  $C^*(L_M)$ . Then one proves directly that it induces an isomorphism in cohomology. The fact that  $A$  is also a model for  $\Gamma$  was proved in a similar way (cf. [14]).

When  $M$  has a finite dimensional model, one can construct a model for  $\Gamma$  which is finite dimensional in each degree, and with it one can make explicit calculations.

Note that the inclusion  $C_{\Delta}^*(L_M, \Omega_M) \rightarrow C^*(L_M, \Omega_M)$  is a model for the evaluation map  $\Gamma \times M \rightarrow E$  associating to a section  $s$  and a point  $x$  of  $M$  the element  $s(x)$  of  $E$ .

For computations along the lines of the spectral sequence of Gelfand-Fuks, see Cohen and Taylor [22].

The proof of theorem 1' is very similar to the proof of theorem 1. In the next paragraph, we explain the construction of an algebraic model for  $\Gamma_G$  suitable for computations. In § 6, we indicate briefly why this is a model for  $\Gamma_G$ .

## 5. CONSTRUCTION OF AN ALGEBRAIC MODEL FOR THE SPACE OF SECTIONS OF A FIBER BUNDLE ([20], [18], [13]).

As a guide, consider first the geometric situation. Let  $p: E \rightarrow M$  be a fiber bundle with base space  $M$ , fiber  $F$  and let  $\Gamma$  be the space of continuous sections of  $E$ .

We have the commutative diagramm

$$\begin{array}{ccc}
 & & e \\
 & & \longrightarrow \\
 M \times \Gamma & \xrightarrow{\quad} & E \\
 \downarrow & \searrow & \swarrow p \\
 & & M \\
 \downarrow & & \downarrow \\
 \Gamma & & * \\
 & \searrow & \\
 & & *
 \end{array}$$

1)

where  $e$  is the evaluation map associating to the point  $x$  of  $M$  and the section  $s$  the point  $s(x)$  of  $E$ . The other maps are natural projections (\* is a point).

Suppose that a topological group  $G$  acts on  $M$  and also on  $E$  in a way compatible with  $p$ . Then  $G$  acts also on  $\Gamma$ , and all the maps in the diagramm are equivariant.

For a space  $X$  on which  $G$  acts, let us denote by  $X_G$  the bundle with fiber  $X$  associated to the principal universal  $G$ -bundle  $P$  with base space  $BG (= *_G)$ .

From 1) we get the corresponding commutative diagramm

$$\begin{array}{ccc}
 (M \times \Gamma)_G & \longrightarrow & E_G \\
 \downarrow & \searrow & \swarrow \\
 \Gamma_G & & M_G \\
 & \searrow & \downarrow \\
 & & BG
 \end{array}$$

2)

We try now to construct an algebraic analogue of this diagramm. *We assume that the connectivity of the fiber  $F$  of  $E$  is bigger than the dimension  $n$  of  $M$ .*

Choose a  $DG$ -algebra  $B$  which is a model of  $BG$  and assume that we can represent the bundle  $M_G$  by a  $DG$ -algebra  $A$ , the projection being represented by a morphism  $B \rightarrow A$ , and such that  $A$ , as a module over  $B$ , is free and finite dimensional with a basis  $s_1, \dots, s_k$ , where the degree of  $s_i$  is not bigger than  $n$  (see examples below).

Then we construct the Postnikov decomposition (or minimal model) of the bundle  $E_G \rightarrow M_G$ . Algebraically, this means that we take a model for  $E_G$  which is a tensor product  $A \otimes \Lambda(x_\alpha)$ , where  $\Lambda(x_\alpha)$  is a free graded algebra on an ordered set of generators  $x_\alpha$ , the differential of each  $x_\alpha$ , being in the subalgebra generated by  $A$  and the preceding  $x_\beta$ . Of course the natural inclusion of  $A$  in  $A \otimes \Lambda(x_\alpha)$  has to be a model for the projection  $E_G \rightarrow M_G$ . Such a model, with a finite number of generators  $x_\alpha$  in each degree, always exists if  $F$  is 1-connected and with finite dimensional cohomology, and if  $G$  is a connected Lie group (cf. [13], [18]).

A model for  $\Gamma_G$  will be the algebra  $B \otimes \Lambda(x^i_\alpha)$ , where  $\Lambda(x^i_\alpha)$  is the free algebra on generators  $x^i_\alpha$ ,  $i = 1, \dots, k$ , and  $\deg x^i_\alpha = \deg x_\alpha - \deg s^i$ . By our assumptions,  $\deg x^i_\alpha > 0$ .

A model for the map  $e$  will be the morphism

$$\varepsilon: A \otimes \Lambda(x_\alpha) \rightarrow B \otimes \Lambda(x^i_\alpha)$$

of  $A$ -algebras defined by

$$\varepsilon(1 \otimes x_\alpha) = \sum_i s^i \otimes x_\alpha^i.$$

The differential on  $B \otimes \Lambda(x^i)$  is then uniquely defined by the conditions that  $B \otimes \Lambda(x^i)$  should be a  $DG$ -algebra over  $B$  and that  $\varepsilon$  should commute with the differential given by the isomorphism with  $A \otimes_B (B \otimes \Lambda(x^i_\alpha))$ .

The algebraic analogue of diagramm 2) is the commutative diagramm of  $DG$ -algebras

$$\begin{array}{ccccc}
 & & A \otimes_B (B \otimes \Lambda(x^i_\alpha)) & \longleftarrow & A \otimes \Lambda(x_\alpha) \\
 & & \uparrow & \swarrow & \nearrow \\
 2) & & B \otimes \Lambda(x^i_\alpha) & & A \\
 & & \swarrow & \uparrow & \\
 & & & B & 
 \end{array}$$

*Examples.*

1. For  $M$ , take the 2-sphere  $S^2$  and for  $E$  the trivial bundle  $S^2 \times S^4$ , so that  $\Gamma$  is the space of continuous maps of  $S^2$  in  $S^4$ . The group  $G$  will be the rotation group  $SO_3$  acting on  $S^2$  as usual and trivially on  $S^4$ .

As model  $B$  for  $BG$  we take the polynomial algebra  $R[p_1]$  in a generator  $p_1$  of degree 4. A model for  $M_G$  is the algebra  $A$  quotient of the polynomial algebra  $\Lambda(s, p_1)$ , where  $\deg s = 2$ , by the ideal generated by  $s^2 - p_1$ . The differential is zero. The elements 1 and  $s$  form a basis for the  $B$ -module  $A$ .

A minimal model for the bundle  $E_G$  is  $A \otimes \Lambda(x, y)$ , where  $\Lambda(x, y)$  is the free algebra with generators  $x$  of degree 4, and  $y$  of degree 7, and  $dy = x^2$ .

According to the preceding recipe, a model for  $\Gamma_G$  is the algebra  $R[p_1] \otimes \Lambda(x, y, \bar{x}, \bar{y})$  with  $\deg \bar{x} = 2$ ,  $\deg \bar{y} = 5$ , the image of  $x$  by  $\varepsilon$  being  $1 \otimes x + s \otimes \bar{x}$ , similarly for  $y$ . The differential is given by  $dx = d\bar{x} = 0$ ,  $dy = x^2 + p_1 \bar{x}^2$ ,  $d\bar{y} = 2x\bar{x}$ .

2. Take  $M$  as the circle,  $E$  as the product  $S^1 \times F$ , where  $F$  is a simply connected space, so that  $\Gamma$  is just the space of continuous maps of  $S^1$  in  $F$  (case studied by Sullivan [19]). For  $G$  we take the group of rotations of the circle, acting trivially on  $F$ .

Represent  $F$  by its minimal model  $\Lambda(x_\alpha)$ . A model  $B$  for  $BG$  is the polynomial algebra  $R[e]$  in a generator  $e$  of degree 2 and a model  $A$  for  $M_G$

is the free commutative algebra  $\Lambda(s, e)$ , where  $\deg s = 1$  and  $ds = e$ . As a  $B$ -module, it is free with basis 1 and  $s$ . A model for  $E_G$  is just  $\Lambda \otimes \Lambda(x_\alpha)$ .

As model for  $\Gamma_G$ , we take  $R[e] \otimes \Lambda(x_\alpha, \bar{x}_\alpha)$ , where  $\deg \bar{x}_\alpha = \deg x_\alpha - 1$ , the image of  $x_\alpha$  by  $\epsilon$  being  $1 \otimes x_\alpha + s \otimes \bar{x}_\alpha$ . The differential  $d$  is described as follows (compare with Sullivan [18] or [19]). Let  $h$  be the derivation of degree  $-1$  of  $\Lambda(x_\alpha, \bar{x}_\alpha)$  given by  $hx_\alpha = \bar{x}_\alpha$  and  $h\bar{x}_\alpha = 0$ . Then if  $d_0$  denotes the differential in  $\Lambda(x_\alpha)$  identified to a subalgebra of  $\Lambda(x_\alpha, \bar{x}_\alpha)$ , we have

$$de = 0, dx_\alpha = d_0x_\alpha - e\bar{x}_\alpha, d\bar{x}_\alpha = -hd_0x_\alpha$$

*Remark.* In the case where  $E$  is the bundle described in § 4, its minimal model  $\Lambda \otimes \Lambda(x_\alpha)$  over  $M_G$  is complicated, because there is an infinite number of generators  $x_\alpha$  (except for  $n=1$ ) labelled by a basis of the rational homotopy of a wedge of spheres, so by a basis of the free graded Lie algebra  $L(n)$  generated by the spheres of this wedge (cf. [13]).

## 6. SKETCH OF THE PROOF OF THE MAIN THEOREM AND APPLICATIONS

We represent the universal principal  $G$ -bundle as a limit of finite dimensional bundles  $P_k$  and we denote by  $\Omega_P$  the inverse limit of algebras of forms  $\Omega_{P_k}$ .

First note that we can replace  $C^*(L_M; G)$  by the  $DG$ -algebra  $C^*(L_M, \Omega_P)_G$  of  $G$ -basic elements in  $C^*(L_M, \Omega_P)$  (compare with Cartan [5], exposé 20).

A model for  $E_G$  will be the algebra  $C_\Delta^*(L_M, \Omega_{M \times P})_G = [C_\Delta^*(L_M, \Omega_M \hat{\otimes} \Omega_P)_G]$  and a model for the evaluation map will be the inclusion of this  $DG$ -algebra in  $C^*(L_M, \Omega_{M \times P})_G$ .

In the construction of § 5, we choose  $B = \Omega_{BG}$  as model for  $BG$  and, instead of taking for  $A$  a finite dimensional module over  $B$ , we take the  $DG$ -algebra  $\Omega_{M_G} \approx [\Omega_{M \times P}]_G$  as model for  $M_G$ . We have to build the model for  $\Gamma_G$  along the same lines as in § 5, but in more intrinsic terms like in [13]. The minimal model (or Postnikov decomposition of  $E_G$ ) will be of the form  $A \otimes S^*(V)$ , where  $S^*(V)$  denotes the algebra of symmetric multilinear forms on a graded vector space  $V$  (cf. [13]).

As an algebra, the model for  $\Gamma_G$  will be the algebra  $S_B^*(A \otimes V, B)$  of continuous symmetric  $B$ -multilinear forms on the graded  $B$ -module  $A \otimes V$ . One can construct a map of this model in  $C^*(L_M, \Omega_{M \times P})_G$  and prove that it induces an isomorphism in cohomology.