

§5. Proofs of Theorem 1.21 and related results

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **28 (1982)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **20.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

THEOREM 4.13. *Suppose that there exists a real number $\gamma(E) > 0$ such that (1.7) holds. Let $\varepsilon > 0$, and suppose that $x \geq c_{34}(E, \varepsilon)$ and $(\log_2 x)^2 (\log x)^{-1} \leq \alpha \leq 1 + \{1 + \log \gamma(E) - \varepsilon\} (\log_2 x)^{-1}$.*

Then

$$\begin{aligned} x^{1-\alpha} \exp \left\{ -c_{35}(E) \frac{\log x}{\log_2 x} \right\} &\leq S(x, \alpha (\log x) (\log_2 x)^{-1}; E, \omega) \\ &\leq x^{1-\alpha} \exp \left\{ \frac{2\alpha (\log x) \log_3 x}{\log_2 x} + c_{36}(E) \frac{\log x}{\log_2 x} \right\}. \end{aligned}$$

This can be obtained from Theorems 1.11 and 1.14 (take

$$y = \alpha (\log x) (\log_2 x)^{-1}$$

and use the inequalities

$$\log_2 y \leq \log_3 x, y \geq \log_2 x \geq \gamma(E) \log_2 x.$$

Theorem 4.13 should be compared with Theorem 1.6.

§5. PROOFS OF THEOREM 1.21 AND RELATED RESULTS

In estimating $S(x, y; E, \Omega)$ (defined by (1.1)), we do not need any assumption such as (1.7). Hence we emphasize that throughout the remainder of this paper, E is merely assumed to be any nonempty set of primes. (We shall sometimes assume explicitly that E has at least two members.) The smallest member of E will always be denoted by p_1 (and the smallest member of $E - \{p_1\}$, if it exists, by p_2). When x and v are positive real numbers, the function $\Lambda = \Lambda(x, v; E)$ is always defined by (1.22).

The subsequent work depends heavily on the following elementary lemma [13, p. 690]:

LEMMA 5.1. *If $x > 0$ and $1 \leq z < p_1$, then*

$$\sum_{n \leq x} z^{\Omega(n; E)} < p_1 (p_1 - z)^{-1} x e^{(z-1)E(x) + 4z}.$$

For the special case $E = P$, there is a recent paper of DeKoninck and Hensley [1] giving various estimates for $\sum_{n \leq x}^* z^{\Omega(n)}$, where z is complex and * indicates that the prime factors of n are restricted to lie in a certain range. DeKoninck and Hensley get sharp results, but their work is rather complicated and does not seem applicable to the problems discussed here.

If y is real and $z \geq 1$, then

$$\begin{aligned} \sum_{n \leq x} z^{\Omega(n; E)} &\geq \sum_{n \leq x, \Omega(n; E) \geq y} z^{\Omega(n; E)} \\ &\geq z^y \text{card} \{n \leq x : \Omega(n; E) \geq y\}. \end{aligned}$$

Hence Lemma 5.1 immediately yields

LEMMA 5.2. *If $x > 0$, y is real, and $1 \leq z < p_1$, then*

$$\begin{aligned} &\text{card} \{n \leq x : \Omega(n; E) \geq y\} \\ &< p_1 (p_1 - z)^{-1} x \exp \{(z-1) E(x) - y \log z + 4z\}. \end{aligned}$$

LEMMA 5.3. *Let $x > 0$, $0 < v \leq y < p_1 v$. Then*

$$\begin{aligned} &\text{card} \{n \leq x : \Omega(n; E) \geq y\} \\ &< c_{37} (p_1) (p_1 - y/v)^{-1} x \exp \{y - v - y \log (y/v) + p_1 \Lambda\}. \end{aligned}$$

Proof: In Lemma 5.2, use the inequality $E(x) \leq v + \Lambda$ and take $z = y/v$ to get an approximate minimum. Q.E.D.

We observe in passing that Lemma 5.2 can also be used when $y \geq p_1 v$. In order to get a reasonably good result in this case by the same method, one needs to minimize the function

$$g(z) = (z-1)v - y \log z - \log(p_1 - z)$$

on the interval $1 \leq z < p_1$. Assuming that y is rather large, one can see with some computation that $g(z)$ is approximately minimized when

$$z = p_1 (1 - (2y)^{-1}),$$

and this z satisfies $1 \leq z < p_1$ whenever $y \geq 1$. With this value of z , Lemma 5.2 yields

$$\text{card} \{n \leq x : \Omega(n; E) \geq y\} \leq c_{38} (p_1) y p_1^{-y} x e^{(p_1-1)v + p_1 \Lambda} \quad (5.4)$$

for $x > 0$, $y \geq 1$. When E is the set of all primes and $x \geq 3$, we can take $v = \log_2 x$, $\Lambda = O(1)$. Thus (5.4) is already sharper and more general

than (1.20) (which is due to Erdős and Sárközy [3]). However, Theorem 1.18 shows that it may be of interest to take y as large as $(\log x)(\log p_1)^{-1}$, and we shall now prove that when y is relatively large, the factor y on the right-hand side of (5.4) can be replaced by a much smaller quantity.

LEMMA 5.5. Write $F = E - \{p_1\}$ (if F is empty, we define $\Omega(n; F) = 0$ for all n). Let $x > 0, y \geq 0$, and let $k = [y] + 1$. For integers a with $0 \leq a \leq k$, define

$$C_a = \{m \leq xp_1^{-a} : p_1 \nmid m \text{ and } \Omega(m; F) \geq k - a\}.$$

Then

$$S(x, y; E, \Omega) = [xp_1^{-k}] + \sum_{a=0}^{k-1} \text{card } C_a.$$

Proof: For $0 \leq a \leq k$, define

$$B_a = \{n \leq x : p_1^a \parallel n \text{ and } \Omega(np_1^{-a}; F) \geq k - a\}$$

(recall that $p_1^a \parallel n$ means $p_1^a \mid n$ and $p_1^{a+1} \nmid n$). It is easy to see that

$$\{n \leq x : \Omega(n; E) > y\} = \{n \leq x : p_1^k \mid n\} \cup \bigcup_{a=0}^{k-1} B_a.$$

Since the sets $\{n \leq x : p_1^k \mid n\}, B_0, B_1, \dots, B_{k-1}$ are disjoint, we have

$$S(x, y; E, \Omega) = \text{card } \{n \leq x : p_1^k \mid n\} + \sum_{a=0}^{k-1} \text{card } B_a.$$

But the mapping $n \mapsto np_1^{-a}$ establishes a one-to-one correspondence between B_a and C_a , so the result follows. Q.E.D.

Proof of Theorem 1.21: If $E = \{p_1\}$, then by Lemma 5.5,

$$S(x, y; E, \Omega) \leq xp_1^{-y},$$

and (1.23) follows. Thus we may assume that $F = E - \{p_1\}$ is not empty. Let p_2 be the smallest member of F , and let $k = [y] + 1$. By Lemma 5.5,

$$S(x, y; E, \Omega) = [xp_1^{-k}] + \sum_{a=1}^k \text{card } C_{k-a}. \quad (5.6)$$

To estimate

$$\text{card } C_{k-a} = \text{card } \{m \leq xp_1^{a-k} : p_1 \nmid m \quad \text{and} \quad \Omega(m; F) \geq a\}$$

from above, we apply Lemma 5.2 (with E replaced by F and p_1 by p_2). Since

$$F(xp_1^{a-k}) \leq F(x) \leq E(x) \leq v + \Lambda,$$

we obtain

$$\begin{aligned} \text{card } C_{k-a} &< p_2 (p_2 - z)^{-1} xp_1^{a-k} \exp \{(z-1)(v+\Lambda) - a \log z + 4z\} \\ &= H(a, z), \end{aligned} \tag{5.7}$$

say, and this holds for each integer a ($1 \leq a \leq k$) and each real z with $1 \leq z < p_2$. In applying (5.7), we are free to choose z to depend on a . Write $Q = \max \{k, p_1 v\}$, and for each a ($1 \leq a \leq Q$), let z_a be any real number satisfying $1 \leq z_a < p_2$. Then by (5.6) and (5.7),

$$\begin{aligned} S(x, y; E, \Omega) &\leq xp_1^{-k} + \sum_{a=1}^k H(a, z_a) \\ &\leq xp_1^{-k} + \sum_{1 \leq a \leq v} H(a, z_a) + \sum_{v < a \leq p_1 v} H(a, z_a) \\ &\quad + \sum_{p_1 v < a \leq Q} H(a, z_a). \end{aligned} \tag{5.8}$$

For $1 \leq a \leq v$, take $z_a = 1$. With this choice, we have

$$\begin{aligned} \sum_{1 \leq a \leq v} H(a, z_a) &\ll xp_1^{-k} \sum_{1 \leq a \leq v} p_1^a \ll xp_1^{-y+v} \\ &\ll xp_1^{-y} e^{(p_1-1)v}. \end{aligned} \tag{5.9}$$

For $v < a \leq p_1 v$, the quantity $(z-1)v - a \log z$ in (5.7) is minimized by taking $z = a/v = z_a$. With this choice of z_a , we have $1 < z_a \leq p_1$ and

$$p_2 (p_2 - z_a)^{-1} \leq p_2 (p_2 - p_1)^{-1} \leq 1 + p_1,$$

so

$$H(a, z_a) \leq c_{39} (p_1) xp_1^{a-k} e^{(p_1-1)\Lambda} (v^a e^{-v} / a^a e^{-a}).$$

By Stirling's formula, $a^a e^{-a} \gg a! a^{-1/2}$, so we get

$$\begin{aligned} \sum_{v < a \leq p_1 v} H(a, z_a) &\leq c_{40} (p_1) xp_1^{-y} v^{1/2} e^{-v+p_1\Lambda} \sum_{v < a \leq p_1 v} \frac{(p_1 v)^a}{a!} \\ &\leq c_{40} (p_1) xp_1^{-y} v^{1/2} e^{(p_1-1)v+p_1\Lambda}. \end{aligned} \tag{5.10}$$

For $p_1 v < a \leq Q$, we let all the numbers z_a have the same value $p_1 (1 + \theta)$, where θ is a real number about which we assume only that $0 < \theta < p_2 p_1^{-1} - 1$ (the last inequality being needed in order to have $z_a < p_2$). With this choice of z_a , (5.7) yields

$$\begin{aligned} & \sum_{p_1 v < a \leq Q} H(a, z_a) \\ & \leq p_2 \{p_2 - p_1 (1 + \theta)\}^{-1} x p_1^{-k} \exp \{(p_1 - 1 + p_1 \theta) (v + \Lambda) + 4p_1 (1 + \theta)\} \\ & \quad \times \sum_{p_1 v < a \leq Q} (1 + \theta)^{-a}. \end{aligned} \quad (5.11)$$

The last sum on the right does not exceed

$$\sum_{a > p_1 v} (1 + \theta)^{-a} < (1 + \theta) \theta^{-1} (1 + \theta)^{-p_1 v}. \quad (5.12)$$

After combining this estimate with (5.11), we would like to minimize the contribution of the essential terms $e^{p_1 \theta v} \theta^{-1} (1 + \theta)^{-p_1 v}$. Since

$$\log(1 + \theta) \geq \theta - \theta^2/2 \quad \text{for } \theta \geq 0, \quad (5.13)$$

we have

$$p_1 \theta v - \log \theta - p_1 v \log(1 + \theta) \leq -\log \theta + p_1 v \theta^2/2,$$

and here the right-hand side would be minimized by taking θ to be $(p_1 v)^{-1/2}$. However, we must also choose $\theta < p_2 p_1^{-1} - 1$ (so that $z_a < p_2$). If we take

$$\theta = (2p_1 v^{1/2})^{-1}, \quad (5.14)$$

then because of our assumption that $v \geq 1$, we have

$$\theta \leq (2p_1)^{-1} < p_2 p_1^{-1} - 1.$$

Combining (5.11), (5.12), (5.13), and (5.14), and observing that

$$\begin{aligned} p_2 \{p_2 - p_1 (1 + \theta)\}^{-1} & \leq p_2 (p_2 - p_1 - 1/2)^{-1} \\ & = 1 + (p_1 + 1/2) (p_2 - p_1 - 1/2)^{-1} \leq c_{41} (p_1), \end{aligned}$$

we obtain finally

$$\sum_{p_1 v < a \leq Q} H(a, z_a) \leq c_{42} (p_1) x p_1^{-y} v^{1/2} e^{(p_1 - 1)v + p_1 \Lambda}. \quad (5.15)$$

The theorem now follows from (5.8), (5.9), (5.10), and (5.15). Q.E.D.

Since

$$E(x) \leq \sum_{p \leq x} p^{-1} = \log_2 x + O(1) \quad \text{for } x \geq 2,$$

one would always want to choose $v \leq \log_2 x$. Thus (1.23) is superior to (5.4) whenever $y \geq (\log_2 x)^{1/2}$. Furthermore, consideration of derivatives shows that

$$y - v - y \log(y/v) \leq (p_1 - 1)v - y \log p_1 \quad \text{for } 0 < v \leq y \leq p_1 v,$$

and hence Lemma 5.3 is superior to Theorem 1.21 whenever

$$1 \leq v \leq y \leq p_1 v - v^{1/2}.$$

REFERENCES

- [1] DEKONINCK, J.-M. and D. HENSLEY. Sums taken over $n \leq x$ with prime factors $\leq y$ of $z^{\Omega(n)}$, and their derivatives with respect to z . *J. Indian Math. Soc. (N.S.)* 42 (1978), 353-365.
- [2] ERDÖS, P. et J.-L. NICOLAS. Sur la fonction : nombre de facteurs premiers de N . *Ens. Math.* 27 (1981), 3-30.
- [3] ERDÖS, P. and A. SÁRKÖZY. On the number of prime factors of integers. *Acta Sci. Math. (Szeged)* 42 (1980), 237-246.
- [4] HALÁSZ, G. Remarks to my paper "On the distribution of additive and the mean values of multiplicative arithmetic functions". *Acta Math. Acad. Sci. Hungar.* 23 (1972), 425-432.
- [5] HARDY, G. H. and S. RAMANUJAN. The normal number of prime factors of a number n . *Quart. J. Math.* 48 (1917), 76-92.
- [6] HARDY, G. H. and E. M. WRIGHT. *An introduction to the theory of numbers*. 3rd ed., Oxford Univ. Press, Oxford, 1954.
- [7] KOLESNIK, G. and E. G. STRAUS. On the distribution of integers with a given number of prime factors. *Acta Arith.* 37 (1980), 181-199.
- [8] KUBILIUS, J. P. *Probabilistic methods in the theory of numbers*. Amer. Math. Soc. Translations of Mathematical Monographs, vol. 11, Providence, R.I., 1964.
- [9] ——— On large deviations of additive arithmetic functions. *Trudy Mat. Inst. Steklov.* 128 (1972), 163-171 (= *Proc. Steklov Inst. Math.* 128 (1972), 191-201).
- [10] LAURINČIKAS, A. On large deviations of arithmetic functions. *Litovsk. Mat. Sb.* 16 (1976), No. 1, 159-171 (= *Lithuanian Math. Trans.* 16 (1976), 97-104).
- [11] NEWMAN, M. Isometric circles of congruence groups. *Amer. J. Math.* 91 (1969), 648-656.
- [12] NORTON, K. K. Numbers with small prime factors, and the least k -th power non-residue. *Mem. Amer. Math. Soc.* 106 (1971).
- [13] ——— On the number of restricted prime factors of an integer. I. *Illinois J. Math.* 20 (1976), 681-705.