

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 33 (1987)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INVOLUTIONS IN SURFACE MAPPING CLASS GROUPS
Autor: McCarthy, John / Papadopoulos, Athanase
Kapitel: 5. Remarks and examples
DOI: <https://doi.org/10.5169/seals-87897>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Let C and C_1 be two simple closed curves on F representing respectively the classes $\mathbf{C}$ and $\mathbf{C}_1$, in such a way that C and C_1 are in a position of minimum-intersection number.

Consider a neighborhood of the union of C and C_1 obtained by taking the union of a thin tubular neighborhood of each of these curves, and let C_2 denote the collection of those boundary curves of this neighborhood which are not null-homotopic.

Suppose first of all that C_2 is not empty. Then we have $s_1(C_2) = C_2$ and $s_2(C_2) = C_2$. (To see this, one can represent s_1 (respectively s_2) by an isometry of some hyperbolic metric, and then consider the geodesics g and g_1 in the classes of C and C_1 . The isometry preserves the geodesics union $g \cup g_1$ and therefore it preserves an imbedded ε -neighborhood of that subset, and the boundary of the neighborhood). In this case, s_1 and s_2 have a common fixed point in $\mathbf{PMF}$.

Suppose now that C_2 is empty. We have $s_1 \circ s_2(\mathbf{C}) = \mathbf{C}$ and $s_1 \circ s_2(\mathbf{C}_1) = \mathbf{C}_1$, and $\mathbf{C}$ and $\mathbf{C}_1$ have the property that for any element $\mathbf{F}$ in $\mathbf{MF}$, we have either $i(\mathbf{F}, \mathbf{C}) \neq 0$ or $i(\mathbf{F}, \mathbf{C}_1) \neq 0$.

By assumption, $s_1 \circ s_2$ is reducible. Let n be an integer s.t. the map $(s_1 \circ s_2)^n$ preserves each component of the surface F cut along the reducing curve.

The mapping class $(s_1 \circ s_2)^n$ cannot have any pseudo-Anosov component, since if it had one, and if $\mathbf{F}^u$ denotes the class of the unstable foliation of that component, we have either $i(\mathbf{F}^u, \mathbf{C}) \neq 0$ or $i(\mathbf{F}^u, \mathbf{C}_1) \neq 0$. By the dynamics of a pseudo-Anosov (component) map on measured foliations space, the two classes of curves cannot be fixed by $s_1 \circ s_2$. Therefore, $s_1 \circ s_2$ cannot have pseudo-Anosov components.

So $(s_1 \circ s_2)^n$ has only finite order components.

By the same argument, $(s_1 \circ s_2)^n$ cannot have a non-trivial Dehn twist along a component of its reducing curve.

Therefore, $s_1 \circ s_2$ has only periodic components with no non-trivial Dehn twists along the reducing curve, so it is globally periodic, i.e. of finite order, a contradiction.

We conclude that $s_1 \circ s_2$ is pseudo-Anosov. This proves theorem 2.

5. REMARKS AND EXAMPLES

1. We can easily classify now the structure of the group generated by two involutions:

Given the two involutions s_1 and s_2 of $M(F)$, the subgroup G they generate is an order-2 extension of the cyclic subgroup generated by $s_1 \circ s_2$. The elements of G that are not in that subgroup are all conjugate to s_1 or s_2 . If s_1 and s_2 have a common fixed point in $\mathbf{T}$, the subgroup that they generate is finite. Otherwise, it is isomorphic to the infinite dihedral group $Z_2 * Z_2$.

2. In closing, we wish to point out that all three cases of Theorem 2 do in fact occur in every genus: To see that $s_1 \circ s_2$ can be of finite order we can take s_1 to be an horizontal rotation as in figure 2 and s_2 to be a vertical rotation as in figure 3. Since these rotations commute, $s_1 \circ s_2$ is an involution. (This example obviously generalizes to genus greater than two.)

To see that $s_1 \circ s_2$ can be reducible of infinite order, we can take s_1 to be a vertical rotation as in figure 3 and let $s_2 = s_1 \circ t_1 \circ t_b^{-1}$. Now s_2 is an involution by equation (1):

$$(17) \quad (s_2)^2 = s_1 \circ t_1 \circ t_b^{-1} \circ s_1 \circ t_1 \circ t_b^{-1} = t_b \circ t_1^{-1} \circ t_1 \circ t_b^{-1} = 1.$$

Moreover, $s_1 \circ s_2 = t_1 \circ t_b^{-1} - 1$ which is a reducible map of infinite order. (Again, this example obviously generalizes to higher genera.)

To see that $s_1 \circ s_2$ can be pseudo-Anosov we can make a similar construction. Let s_1 be an involution. Suppose that A is a family of disjoint nontrivial simple closed curves. Let $B = s_1(A)$. Now suppose that A and B fill up F . Let t_A be the product of the Dehn twists about the components of A and t_B be the corresponding product associated to B . Let $s_2 = s_1 \circ t_A \circ t_B^{-1}$. As in the reducible case just described, s_2 is an involution. Furthermore, $s_1 \circ s_2 = t_A \circ t_B^{-1}$, which is a pseudo-Anosov map by an algorithm of Long's [6] generalizing Thurston's algorithm described in [4]. An example of this construction of case (iii) of Theorem 2 is depicted in figure 11, where s_1 is again the vertical rotation. (Again, this example easily generalizes.)

Alternatively, one can give a nonconstructive argument as follows. Let s_1 be a vertical rotation as in figure 3. Since $s_1(a_1) = b$, we know that s_1




FIGURE 11.

is not in $\text{Fix}(s_1)$. On the other hand, $\text{Fix}(s_1)$ is clearly a closed set. Hence, we may find an open neighborhood of a_1 in $\mathbf{T} \cup \mathbf{PMF}$, U , such that U avoids $\text{Fix}(s_1)$. Now, we may find a pseudo-Anosov, f , both of whose fixed points lie in U . (For example, this can be achieved by conjugating any given pseudo-Anosov by a sufficiently high power of t_1 .) Since $\text{Fix}(s_1)$ is a compact set which avoids the repelling fixed point of f , it follows from the well known behavior of pseudo-Anosov maps on $\mathbf{T} \cup \mathbf{PMF}$ that $f^n(\text{Fix}(s_1))$ is contained in U for sufficiently large n . Choose n subject to this condition and let $s_2 = f^n \circ s_1 \circ f^{-n}$. Finally, since $\text{Fix}(s_2)$ is equal to $f^n(\text{Fix}(s_1))$, it follows from Theorem 2 that $s_1 \circ s_2$ is pseudo-Anosov.