L'Enseignement Mathématique
Commission Internationale de l'Enseignement Mathématique
37 (1991)
1-2: L'ENSEIGNEMENT MATHÉMATIQUE
COHOMOLOGY OF FINITELY GENERATED ABELIAN GROUPS
Huebschmann, Johannes
1. Introduction
https://doi.org/10.5169/seals-58730

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 06.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# COHOMOLOGY OF FINITELY GENERATED ABELIAN GROUPS

# par Johannes HUEBSCHMANN

# 1. INTRODUCTION

Let Q be a finitely generated abelian group. In this note we compute its integral cohomology explicitly. Our approach is to construct a suitable small model  $\mathscr{A}(Q)$  for the algebra of cochains on Q. Our model is very simple and relies on some differential homological algebra that has been created more than three decades ago [2-4], [8], [9], [28], [29]. Here is the model:

THEOREM A. With coefficients in an arbitrary commutative R, the cohomology ring of a finitely generated abelian group Q, given by a presentation

$$Q = \langle f_1, ..., f_m, t_1, ..., t_n ; t_j^{l_j} = 1 \rangle$$
,

is as an R-algebra isomorphic to the homology algebra of the following primitively generated cocommutative differential graded Hopf algebra  $\mathscr{A}(Q)$ :

Generators:  $z_1, ..., z_m, |z_j| = 1, x_1, ..., x_n, |x_i| = 1, c_1, ..., c_n, |c_i| = 2.$ 

Relations: The generators commute in the graded sense, save that the squares of the  $x_i$ 's are non-zero and given by

(1.1) 
$$x_i^2 = \frac{l_i(l_i-1)}{2} c_i, 1 \le i \le n;$$

in particular the squares of the  $z_j$ 's are zero.

Differential:  $d(z_j) = 0, 1 \leq j \leq m, \quad d(x_i) = l_i c_i, \quad d(c_i) = 0, 1 \leq i \leq n.$ 

For appropriate coefficients so that the diagonal map of this Hopf algebra induces a diagonal map on its homology, e.g. if R is a field or if the characteristic of R divides each  $l_i, 1 \le i \le n$ , the isomorphism is one of Hopf algebras.

Supported by the Deutsche Forschungsgemeinschaft under a Heisenberg grant.

### J. HUEBSCHMANN

Here as usual |w| denotes the degree of a homogeneous element w. A proof of this Theorem has been given in Section 3 of our paper [13]. To make the present paper self contained we reproduce the proof in Section 2 below.

The model philosophy has already succesfully been exploited by us in [13-20] for theoretical purposes and to do calculations in group cohomology and algebraic topology. Models of different kinds have been introduced and exploited in the literature for decades, see for example [1-3], [9-11], [21], and the literature there; more about e.g. rational homotopy theory and minimal models may be found in Halperin's book [11].

Our main aim is to deduce an explicit description of the integral cohomology ring of Q from Theorem A. To explain our result, we denote an *infinite* cyclic group by C and a *finite* cyclic group of order l by  $C_l$ . We then write our abelian group Q in the form  $Q = C_{l_1} \times \cdots \times C_{l_n} \times C^m$ . It is clear that, as a graded commutative algebra, the integral cohomology ring  $H^*(Q, \mathbb{Z})$  decomposes as

(1.2) 
$$H^*(Q, \mathbb{Z}) = H^*(C_{l_1} \times \cdots \times C_{l_n}, \mathbb{Z}) \otimes H^*(C^m, \mathbb{Z}) .$$

Moreover, cf. Eckmann [7],

(1.3) 
$$H^*(C^m, \mathbb{Z}) = \Lambda[z_1, \cdots, z_m]$$

where the notation  $z_j$  is abused somewhat. We therefore assume henceforth that Q is finite. Moreover, we suppose that things have been arranged in such a way that

(1.4) 
$$l_1 \mid l_2 \mid ... \mid l_n$$
,

where as usual u | v means that the number u divides the number v. It is classical that this is always possible. From Theorem A we deduce at once that every monomial of the kind

(1.5) 
$$x_i x_j \dots x_k, \quad 1 \leq i < j < \dots < k \leq n,$$

is a cycle in our algebra  $\mathscr{A}(Q)$  when taken over the ground ring  $R = \mathbb{Z}/l_i$  and hence determines a cohomology class  $x_i x_j \dots x_k \in H^*(-, \mathbb{Z}/l_i)$ , where a slight abuse of notation comes into play. For each  $l_i$ , let

(1.6) 
$$\beta_{l_i}: H^*(-, \mathbb{Z}/l_i) \to H^{*+1}(-, \mathbb{Z})$$

be the indicated Bockstein operation and, for each monomial of the kind (1.5), let

(1.7) 
$$\zeta_{x_i x_j \dots x_k} = \beta_{l_i}(x_i x_j \dots x_k) \in H^{*+1}(Q, \mathbb{Z}) ,$$

It is clear that, in our model  $\mathscr{A}(Q)$ , the class  $\zeta_{x_i x_j \dots x_k}$  is represented by the cocycle

(1.8) 
$$\frac{1}{l_i} d(x_i x_j \dots x_k) = c_i x_j \dots x_k - \frac{l_j}{l_i} c_j x_i \dots x_k + \dots \pm \frac{l_k}{l_i} c_k x_i x_j \dots x_{k-1}$$
.

Moreover, inspection of the Bockstein exact sequence

(1.9) 
$$H^*(Q, \mathbf{Z}) \xrightarrow{l_i} H^*(Q, \mathbf{Z}) \to H^*(Q, \mathbf{Z}/l_i) \xrightarrow{\beta_{l_i}} H^{*+1}(Q, \mathbf{Z})$$

shows that the class  $\zeta_{x_i x_j \dots x_k} \in H^{*+1}(Q, \mathbb{Z})$  has exact order  $l_i$ .

We now consider the graded algebra A(Q) that arises from  $\mathscr{A}(Q)$  by introducing the additional relations  $l_i c_i = 0$ . It is clear that, when we write  $\zeta_i$ for the obvious image of  $c_i \in \mathscr{A}(Q)$  in A(Q), as a graded algebra, A(Q) is generated by

(1.10.1) 
$$x_1, ..., x_n, |x_i| = 1, \zeta_1, ..., \zeta_n, |\zeta_i| = 2,$$

subject to the relations that

(1.10.2) the generators commute in the graded sense, save that the squares of the  $x_i$ 's are possibly non-zero and given by

$$x_i^2 = \frac{l_i(l_i-1)}{2} \zeta_i, 1 \leq i \leq n ;$$

and

$$(1.10.3) l_i \zeta_i = 0, 1 \leq i \leq n .$$

We note that when  $l_i$  is odd we have in fact  $x_i^2 = 0$ . By construction, the association

$$c_i \mapsto \zeta_i$$
,  $x_j \mapsto x_j$ 

yields a surjective morphism

$$(1.11) \qquad \qquad \mathscr{A}(Q) \to A(Q)$$

of differential graded algebras where the algebra A(Q) is understood to have zero differential. Since (1.11) is an isomorphism in degree 1 we do not distinguish in notation between  $x_j \in \mathscr{A}(Q)$  and its image in A(Q). For each monomial  $x_i x_j \dots x_k$  of the kind (1.5), we write

$$\tilde{\zeta}_{x_ix_j\ldots x_k}\in A(Q)$$

for the image of  $\zeta_{x_i x_j \dots x_k} \in \mathscr{A}(Q)$  under (1.11), so that

#### J. HUEBSCHMANN

(1.12) 
$$\widetilde{\zeta}_{x_i x_j \dots x_k} = \zeta_i x_j \dots x_k - \frac{l_j}{l_i} \zeta_j x_i \dots x_k + \dots \pm \frac{l_k}{l_i} \zeta_k x_i x_j \dots x_{k-1} \in A(Q)$$
.

Here is our main result.

THEOREM B. For a finite abelian group Q, the association

(1.13) 
$$\zeta_{x_i x_j \dots x_k} \mapsto \tilde{\zeta}_{x_i x_j \dots x_k} \in A(Q)$$

identifies  $H^*(Q, \mathbb{Z})$  with the graded subalgebra of A(Q) generated (as an algebra) by the  $\tilde{\zeta}_{x_i x_j \dots x_k} \in A(Q)$ .

This description of the cohomology ring should be compared with that in Chapman [5]. Our description of the cohomology ring has its advantages and disadvantages. An advantage is that it arises from a "small" model; indeed, modulo a prime p dividing each torsion coefficient  $l_i$ , our model boils down to the cohomology ring itself. To have a model as small as possible is important for explicit computations. A disadvantage of our description is that it is natural in the *presentation only*, and not in the group itself. Invariant descriptions of the *homology* of a finitely generated abelian group have been given by Hamsher [12] and Decker [6] in their Chicago ph. d. theses supervised by S. Mac Lane. We do not known whether invariant descriptions of the *cohomology* of a finitely generated abelian group have ever been worked out.

I am indebted to S. Mac Lane for discussions and for a number of comments about an earlier version of the paper.

# 2. The proof of Theorem A

To make the paper self contained we reproduce the following material from our paper [13]. The contents of the present Section are of course classical but apparently not as well known as they deserve.

Let R be a commutative ring with 1. By a Hopf algebra  $(H, \mu, \Delta, \eta, \varepsilon)$ over R we mean as usual a module H together with the structures  $(\mu, \eta)$  and  $(\Delta, \varepsilon)$  of an algebra and a coalgebra that are compatible, that is,  $\mu$  is a morphism of coalgebras or, equivalently,  $\Delta$  is a morphism of algebras; see e.g. VI.9 in Mac Lane [23]. We mention in passing that some authors call this a bialgebra and require a Hopf algebra to have the additional structure of what is called an *antipode*. For us this is actually of no account since this additional structure will always be present, but there is no need to spell it out. Examples of Hopf algebras are group rings, exterior Hopf algebras, divided