

§3. NILMANIFOLDS

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **38 (1992)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

In order to show the reverse inequality $s \leq k$, we must show that, for $p: \Lambda X \rightarrow \Lambda X / \Lambda^{>k} X$, p^* is injective. Plainly, by Poincaré duality, p^* is injective if and only if $p^*(\tau) \neq 0$. Hence, we prove this.

Suppose $p^*(\tau) = 0$. Let τ denote the representing cocycle in $\Lambda^{\geq k} X$ of the fundamental class τ . Let $p(\tau) = \bar{\tau} \in \Lambda X / \Lambda^{>k} X$ and consider $\bar{\tau}$ as an element in $\Lambda^{\leq k} X$. Now, $p^*(\tau) = 0$, so there exists $\bar{\alpha} \in \Lambda X / \Lambda^{>k} X$ with $d\bar{\alpha} = \bar{\tau}$. Consider $\bar{\alpha} \in \Lambda^{\leq k} X$ as well and note that $p(d\bar{\alpha}) = d\bar{\alpha} = \bar{\tau}$. Therefore, in ΛX

$$d\bar{\alpha} = \bar{\tau} + \Phi, \quad \text{where } \Phi \in \Lambda^{>k} X.$$

Similarly, of course, $\tau = \bar{\tau} + \Omega$ for $\Omega \in \Lambda^{>k} X$ and we obtain,

$$\tau = \bar{\tau} + \Omega = d\bar{\alpha} - \Phi + \Omega$$

with $\Omega - \Phi \in \Lambda^{>k} X$. But this means τ is cohomologous to $\Omega - \Phi \in \Lambda^{>k} X$, contradicting the definition of k . $\square$

§3. NILMANIFOLDS

A *nilmanifold* M is the quotient of a nilpotent Lie group N by a discrete cocompact subgroup π . The description below follows [7].

It is well known that N is diffeomorphic to some $\mathbf{R}^n$ and, therefore, M is a $K(\pi, 1)$. Furthermore, this entails the fact that π is a finitely generated torsionfree nilpotent group.

On the algebraic side, there is a refinement of the upper central series of π ,

$$\pi \supseteq \pi_2 \supseteq \pi_3 \supseteq \cdots \supseteq \pi_n \supseteq 1$$

with each $\pi_i / \pi_{i+1} \cong \mathbf{Z}$ whose length is invariant and is called the *rank* of π . So, for π above, $\text{rank}(\pi) = n$.

This description implies that any $u \in \pi$ has a decomposition $u = u_1^{x_1} \cdots u_n^{x_n}$, where $\langle u_n \rangle = \pi_n, \cdots \langle u_i \rangle = \pi_i / \pi_{i+1}$. The set $\{u_1, \cdots u_n\}$ is called a Malcev basis for π . Using this basis the multiplication in π takes the form

$$u_1^{x_1} \cdots u_n^{x_n} u_1^{y_1} \cdots u_n^{y_n} = u_1^{\rho_1(x, y)} \cdots u_n^{\rho_n(x, y)}$$

where

$$\rho_i(x, y) = x_i + y_i + \tau_i(x_1, \cdots x_{i-1}, y_1, \cdots y_{i-1}).$$

Example. $N = U_n(\mathbf{R})$, the group of upper diagonal matrices with 1's on the diagonal; $\pi = U_n(\mathbf{Z})$. A Malcev basis is given by $\{u_{ij} \mid 1 \leq i < j \leq n\}$ where $u_{ij} = I + e_{ij}$ and

$$\rho_{ij}(x, y) = x_{ij} + y_{ij} + \sum_{i < k < j} x_{ik} y_{kj}.$$

Consider the central extension $\pi_n \rightarrow \pi \rightarrow \bar{\pi}$. The cocycle for the extension is $\tau_n: \bar{\pi} \times \bar{\pi} \rightarrow \mathbf{Z}$. Of course $\bar{\pi}$ is also finitely generated torsionfree with refined upper central series,

$$\bar{\pi} = \frac{\pi}{\pi_n} \supseteq \frac{\pi_2}{\pi_n} \supseteq \cdots \supseteq \frac{\pi_{n-1}}{\pi_n} \supseteq \frac{\pi_n}{\pi_n} = 1.$$

Hence, $\text{rank}(\bar{\pi}) = n - 1$ and

$$\bar{\rho}_i(x, y) = \rho_i((x, 0), (y, 0)) = x_i + y_i + \tau_i(x_1, \cdots x_{i-1}, y_1, \cdots y_{i-1})$$

for $i < n$. Clearly, then, we may iterate this process and decompose π as n central extensions of the form

$$\mathbf{Z} \rightarrow G \rightarrow \bar{G}$$

with cocycles $\tau_i \in H^2(\bar{G}; \mathbf{Z})$ (with untwisted coefficients since the extension is central).

This description allows a geometric formulation:

$$\tau_n \in H^2(\bar{\pi}; \mathbf{Z}) \cong H^2(K(\bar{\pi}, 1); \mathbf{Z}) \cong [K(\bar{\pi}, 1), K(\mathbf{Z}, 2)]$$

by the usual identification of cohomology groups with sets of homotopy classes into $K(\mathbf{Z}, m)$'s. Now, $K(\mathbf{Z}, 2) = \mathbf{CP}(\infty)$, the classifying space for principal S^1 -bundles, so τ_n induces a bundle over $K(\bar{\pi}, 1)$,

$$\begin{array}{ccc} S^1 & \rightarrow & K(\pi, 1) \\ & & \downarrow \\ & & K(\bar{\pi}, 1) \xrightarrow{\tau_n} \mathbf{CP}(\infty). \end{array}$$

The total space of the bundle is clearly $K(\pi, 1)$ since the ensuing short exact sequence of fundamental groups is classified by τ_n .

Now, because we can iterate the algebraic decomposition of π , we obtain an iterated sequence of principal S^1 -bundles classified by the τ_i :

$$\begin{array}{ccccc}
S^1 & \rightarrow & M & = & K(\pi, 1) \\
& & \downarrow & & \\
S^1 & \rightarrow & M_{n-1} & \xrightarrow{\tau_n} & CP(\infty) \\
& & \downarrow & & \\
& & \vdots & & \\
& & \downarrow & & \\
S^1 & \rightarrow & M_1 & \xrightarrow{\tau_2} & CP(\infty) \\
& & \downarrow & & \\
& & * & \xrightarrow{\tau_1} & CP(\infty) .
\end{array}$$

We can assume (by finite dimensionality) that each τ_i has image in a finite $CP(n)$, so thus may be approximated by a smooth map. Hence, each M_j is a compact manifold with

$$\dim(M_j) = \dim(M_{j-1}) + 1 .$$

Thus, $\dim(M) = \text{rank}(\pi) = n$.

§4. CATEGORY OF NILMANIFOLDS

The decomposition of $M = K(\pi, 1)$ into a tower of principal S^1 -bundles is, in fact, the Postnikov decomposition of M with k -invariants the τ_i . By the fundamental theorem of rational homotopy theory, the minimal model has the form,

$$\Lambda(M) = (\Lambda(x_1, \dots, x_n), d) , \quad \deg(x_i) = 1$$

with $dx_i = \tau_i$, where τ_i is a cocycle representing the class $\tau_i \in H^2(M_{i-1}; \mathbf{Z})$. Note that $\Lambda(M)$ is an exterior algebra because all generators are in degree 1. Therefore, since $\dim M = n$, the only possibility for a cocycle representing the fundamental class is $x_1 \cdots x_n$. Hence, $e_0(M) = n$ and this immediately implies,

Proof of Theorem 1. $n = e_0(M) \leq \text{cat}_0(M) \leq \text{cat}(M) \leq \dim M = n$. $\square$

Example. Consider the 3-dimensional Heisenberg group $U_3(\mathbf{R})$ and mod out by $U_3(\mathbf{Z})$. The resulting M is a 3-manifold obtained as a principal bundle,

$$S^1 \rightarrow M \rightarrow T^2$$