Introduction

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **19.03.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.
Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.
Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*
ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

http://www.e-periodica.ch

L'Enseignement Mathématique, t. 39 (1993), p. 233-247

CARROUSEL MONODROMY AND LEFSCHETZ NUMBER OF SINGULARITIES

by Mihai TIBĂR

INTRODUCTION

Let $f: (\mathbf{X}, x) \to (\mathbf{C}, 0)$ be a holomorphic function on an analytic germ $(\mathbf{X}, x)$. Let h_f denote the monodromy of the germ $\Psi_f^\bullet(\mathbf{C}_\mathbf{X}^\bullet)_x$ of neighbouring cycles. One defines its *Lefschetz number*

$$\Lambda(h_f) := \sum_{i \geq 0} (-1)^i \text{ trace } [h_f; \Psi_f^i(\mathbf{C}_\mathbf{X}^\bullet)_x] ,$$

and its *zeta-function*

$$\zeta_{h_f}(t) := \prod_{i \geq 0} \det [I - t \cdot h_f ; \Psi_f^i(\mathbf{C}_\mathbf{X}^\bullet)_x]^{(-1)^{i+1}} .$$

We alternatively denote them by $\Lambda(f)$, respectively $\zeta_f(t)$.

A theorem of Eisenbud and Neumann [EN, Theorem 4.3] asserts that the zeta-function of a *fibred multilink* L is the product of the zeta-functions over all *splice components* of L. If the multilink is defined by some Cerf diagram $\Delta(l, f)$, then $\zeta_f(t)$ becomes the zeta-function of the multilink L, this time with coefficients in a local system. This observation of Némethi [Ne] enables him to prove an inductive formula for $\zeta_f(t)$, in terms of invariants of the so called EN-diagram (splice diagram); compare to the one of Eisenbud and Neumann [EN, p. 96]. Some quite strong results in the 3-dimensional link theory are involved in the proofs.

Our approach is based on Lê's carrousel construction and is therefore more geometric and selfcontained. It yields inductive formulae for $\Lambda(f)$ and $\zeta_f(t)$ directly from the Puiseux parametrization of $\Delta(l, f)$. Moreover, it clarifies the contribution, however essential in general, of the "nonessential" terms in this parametrization — which may be not clear from the definition of the splice diagram of an algebraic link given in [EN, p. 53], simply because such terms are completely omitted. One can therefore compare to our definitions 1.5 ÷ 7.

The formula for $\zeta_f(t)$ will be not the same, but quite similar to the ones before. The ingredients are zeta-functions of fibres over certain periodic points in the carrousel disc. We show in Sections 2 and 3 how to define these points from the Puiseux expansion of $\Delta(l,f)$. We end by some applications.

Aknowledgement. This work is based on a piece of the author's disertation [Ti]. He much benefited from talks with Dirk Siersma, whose paper [Si] incited him to do this research (see 3.8).

1. The carrousel revisited

1.1. We first briefly recall the carrousel construction, following closely [Lê-1] and [Lê-3], then give the necessary definitions for our study. One regards $(\mathbf{X}, x)$ as being embedded in $(\mathbf{C}^N, 0)$, for some sufficiently large $N \in \mathbf{N}$. We assume that, unless otherwise stated, all the irreducible components of $(\mathbf{X}, 0)$ have dimensions greater than 1.

Let $\mathscr{X}$ be a small enough representative of $(\mathbf{X}, 0)$. Let $\Gamma(l,f)$ be the *polar curve* of f with respect to a linear function $l: (\mathbf{X}, 0) \to (\mathbf{C}, 0)$, relatively to a fixed *Whitney stratification* $\mathscr{S}$ on $\mathscr{X}$ which satisfies *Thom condition* (a_f).

The polar curve $\Gamma(l,f)$ exists for a Zariski open subset $\hat{\Omega}_f$ in the space of linear germs $l: (\mathbf{C}^N, 0) \to (\mathbf{C}, 0)$. If one does not impose $\Gamma(l,f)$ to be reduced then one gets a larger set $\Omega_f \supset \hat{\Omega}_f$ which is sometimes useful to deal with (see e.g. Example 2.2). (We only mention that one can enlarge even Ω_f: modify its definition by allowing also nonlinear functions.)

1.2. Let $l \in \Omega_f$ and let $\Phi := (l,f): (\mathbf{X}, 0) \to (\mathbf{C}^2, 0)$. We denote by (u, λ) the pair of coordinates on $\mathbf{C}^2$.

The curve germ (with reduced structure) $\Delta(l,f) := \Phi(\Gamma(l,f))$ is called the *Cerf diagram* (of f with respect to l, relative to $\mathscr{S}$). We shall use the same notation $\Gamma(l,f)$, respectively $\Delta(l,f)$ for suitable representatives of these germs.

There is a fundamental system of "privileged" open polydiscs in $\mathbf{C}^N$, centred at 0, of the form $(D_\alpha \times P_\alpha)_{\alpha \in A}$ and a corresponding fundamental system $(D_\alpha \times D'_\alpha)_{\alpha \in A}$ of 2-discs at 0 in $\mathbf{C}^2$, such that Φ induces, for any $\alpha \in A$, a topological fibration

$$\Phi_\alpha: \mathscr{X} \cap (D_\alpha \times P_\alpha) \cap \Phi^{-1}(D_\alpha \times D'_\alpha \setminus (\Delta(l,f) \cup \{\lambda = 0\})) \to D_\alpha \times D'_\alpha \setminus (\Delta(l,f) \cup \{\lambda = 0\}) .$$