

REMARKS AND PROBLEMS ON FINITE AND PERIODIC CONTINUED FRACTIONS

Autor(en): **Mendès France, Michel**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-60426>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARKS AND PROBLEMS
ON FINITE AND PERIODIC CONTINUED FRACTIONS

by Michel MENDÈS FRANCE

SUMMARY. We present eight problems related to the length of continued fractions of rational numbers and to the length of the period of quadratic surds.

§1. A FRUSTRATING QUESTION

Let a and b be two coprime integers, $1 < b < a$. *Is it true that the sequence $(a/b)^n$, $n = 0, 1, 2, \dots$ is dense (mod 1)?* This very old problem of Pisot and Vijayaraghavan is still unanswered. Pisot, Vijayaraghavan and André Weil did however show that there exist infinitely many cluster points.

Are any one of these cluster points irrational? Even this seems unanswered. We address a simpler question, but before we must define the depth of a rational number x : it is simply the length $\delta(x)$ of the continued fraction of x

$$x = [c_0, c_1, c_2, \dots, c_\delta]$$

where we choose δ to be even ($c_\delta \geq 1$). For example

$$\delta(k) = 0, k \in \mathbf{Z}; \quad \delta\left(\frac{1}{2}\right) = 2; \quad \delta\left(\frac{3}{5}\right) = 4 .$$

Quite obviously $\delta(a/b) = O(\ln(b))$, $1 \leq b < a$ (see [8]).

Suppose that the sequence $(a/b)^n$ has an irrational cluster point ζ (mod 1). Then some subsequence $(a/b)^{n_j}$ (mod 1) tends to ζ hence

$$\delta((a/b)^{n_j}) \rightarrow \infty .$$

This observation led me to ask in 1971 [10] whether it was indeed true that

$$\sup_n \delta((a/b)^n) = \infty$$

whithout any other assumptions than $1 < b < a$, $(a, b) = 1$. The problem was solved by Y. Pouchet in an unpublished letter he sent me [14] and by G. Choquet in a series of Comptes Rendus à l'Académie des Sciences [2].

THEOREM 1. *If a and b are two coprime integers $1 < b < a$ then*

$$\lim_{n \rightarrow \infty} \delta((a/b)^n) = \infty .$$

Choquet's proof involves dynamical systems. He could only show that the "sup" is infinite. Pouchet's proof is number theoretical and uses the Mahler-Ridout theorem which strengthens Roth's famous result on the rational approximations of algebraic numbers.

It is a pity that Pouchet never published his result. Fortunately A. van der Poorten gave some details of the proof in [16].

§2. A QUESTION CONCERNING PISOT NUMBERS

Let $x > 1$ be a real number. Define the set

$$E(x) = \{x^n \pmod{1} \mid n \in \mathbf{N}\} \subset [0, 1]$$

Let $E'(x)$ be the derived set i.e. the set of cluster points of $E(x)$. Define $E^{(n)}(x)$ recursively to be the derived set of $E^{(n-1)}(x)$, $n \geq 1$. In [12] Pisot establishes that if x is a real algebraic number larger than 1 such that $E''(x) = \emptyset$ then x is a Pisot number. I ask the following question.

PROBLEM 1. *Is it true that if $x > 1$ is algebraic and if for some $k \in \mathbf{N}$ ($k \geq 2$) $E^{(k)}(x) = \emptyset$ then x is a Pisot number?*

A positive answer to this problem implies the weak form of Theorem 1, namely that the sup is infinite. Indeed, define $A_0 = \{0\}$ and for $k \geq 1$

$$A_{2k} = \{\zeta \in (0, 1) \mid \delta(\zeta) \leq 2k\} .$$

Let A'_{2k}, A''_{2k}, \dots be the derived sets of A_{2k} . Clearly $A'_{2k} = A_{2k-2}$, therefore

$$A_{2k}^{(k+1)} = \emptyset .$$

Now let $x > 1$ be a rational number which is not an integer. Suppose

$$\sup_n \delta(x^n) < \infty$$

Then for some k

$$E(x) \subset A_{2k}$$

hence

$$E^{(k+1)}(x) = \emptyset .$$

Assuming a positive answer to Problem 1, we conclude that x is a Pisot number, i.e. a rational integer. This contradicts the assumption hence

$$\sup_n \delta(x^n) = \infty . \quad QED$$

§3. MORE QUESTIONS ON $\delta(x^n)$

H. Heilbronn [7], T. Tonkov [15] and finally J.W. Porter [13] improving on one another established that as a tends to infinity

$$\frac{1}{\varphi(a)} \sum_{\substack{b < a \\ (a,b)=1}} \delta\left(\frac{a}{b}\right) = \frac{12}{\pi^2} \ln 2 \ln a + O(1) .$$

Independently, J.D. Dixon [6] showed that for all $\varepsilon > 0$ and for all $a, b, 1 < b < a < x$ with the exception of at most $o(x^2)$ couples, one has

$$\left| \delta\left(\frac{a}{b}\right) - \frac{12}{\pi^2} \ln 2 \ln a \right| \leq (\ln a)^{\frac{1}{2} + \varepsilon} .$$

See H. Daudé's work for a dual result [5]. These results suggest the second problem.

PROBLEM 2. *Is it true that for all coprime a and $b, 1 < b < a$*

$$(1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \delta\left(\left(\frac{a}{b}\right)^n\right) = \frac{12}{\pi^2} \ln 2 \ln b ?$$

The limit should indeed be what is stated above and not

$$\frac{12}{\pi^2} \ln 2 \ln a .$$

since it is b , the smallest of the two integers a and b that seems to control the behavior of $(a/b)^n$. It is also to be noticed that in Dixon's inequality, the $\ln a$ that appears on the left hand side could well be replaced by $\ln b$ since for "almost all" couples a, b , $\ln a \approx \ln b$.

Numerical evidence supports equality (1). Based on computer computation, Chr. Batut and M. Olivier showed that

$$\left| \frac{1}{n} \delta \left(\left(\frac{a}{b} \right)^n \right) - \frac{12}{\pi^2} \ln 2 \ln b \right|$$

is less than .02 for n in the range (4000, 5000) and for $a = 3, b = 2$ on the one hand and $a = 5, b = 2$ on the other hand.

§4. RELATED PROBLEMS

My initial (unsuccessful) attempts to prove Theorem 1 were based on the comparison of $\delta(ax/b)$ to $\delta(x)$. I was hoping that a relationship between both depths would give by induction some results on $\delta(xa^n/b^n)$. This turned out nonconclusive, yet I did obtain some results which I believe are interesting in themselves [10].

Let a, b, c, d be coprime integers and let $\Delta = |ad - bc|$. Consider the Möbius map

$$x \mapsto Tx = \frac{ax + b}{cx + d}.$$

THEOREM 2.

$$\limsup_{\delta(x) \rightarrow \infty} \frac{\delta(Tx)}{\delta(x)} = \Theta(\Delta)$$

where Θ takes odd integral values. As n increases to infinity $\Theta(n)$ behaves like $\ln n$. More precisely let

$$\alpha = \left(2 \ln \frac{1 + \sqrt{5}}{2} \right)^{-1}.$$

Then for all integer $n \geq 1$

$$1 + \alpha \ln n \leq \Theta(n) \leq 2(1 + \alpha \ln n).$$

Θ is linked to the depth by the formula

$$\Theta(n) = \max_{1 \leq b \leq n} \delta \left(\frac{b}{n} \right) + 1.$$

Finally

$$\liminf_{\delta(x) \rightarrow \infty} \frac{\delta(Tx)}{\delta(x)} = \frac{1}{\Theta(\Delta)}.$$

The following table gives the first values of Θ

n	1	2	3	4	5	6	7	8	9	10	...
$\Theta(n)$	1	3	3	3	5	3	5	5	5	5	...

Actually Theorem 1 can be improved. There exist two constants $C_1 = C_1(T)$ and $C_2 = C_2(T)$ such that for all rational x

$$\frac{1}{\Theta(\Delta)} \delta(x) - C_1 \leq \delta(Tx) \leq \Theta(\Delta) \delta(x) + C_2.$$

Both inequalities are sharp apart from the exact values of C_1 and C_2 .

5. MORE QUESTIONS

To every Möbius map T we associate the interval $I(T) = [\Theta^{-1}(\Delta), \Theta(\Delta)]$.

PROBLEM 3. *Is it true that for all $\zeta \in I(T)$ there exists a sequence of rational numbers x_n such that $\lim_{n \rightarrow \infty} \delta(x_n) = \infty$ and*

$$\lim_{n \rightarrow \infty} \frac{\delta(Tx_n)}{\delta(x_n)} = \zeta?$$

PROBLEM 4. *Let T_1, T_2, \dots, T_k be Möbius maps with pairwise coprime determinants $\Delta_1, \Delta_2, \dots, \Delta_k$.*

Is it true that for all

$$(\zeta_1, \zeta_2, \dots, \zeta_k) \in \prod_{i=1}^k I(T_i)$$

there exists a sequence of rational x_n with strictly increasing depths such that for all $i = 1, 2, \dots, k$

$$\lim_{n \rightarrow \infty} \frac{\delta(T_i x_n)}{\delta(x_n)} = \zeta_i?$$

Can k be infinite?

The following result should be mentioned at this point.

THEOREM 3. *There exists a sequence of rational numbers x_n with strictly increasing depths such that for all Möbius maps T*

$$\lim_{n \rightarrow \infty} \frac{\delta(Tx_n)}{\delta(x_n)} = 1 .$$

The proof is quite simple. To each irrational

$$x = [c_0, c_1, c_2, \dots]$$

we associate the sequence of best approximations

$$x_n = \frac{p_n}{q_n} = [c_0, c_1, c_2, \dots, c_n] .$$

Paul Lévy [9] showed that for almost all x

$$\ln q_n \sim \frac{\pi^2}{12 \ln 2} n$$

as n goes to infinity (see for example [1] p. 45). In other words, for almost all x

$$\delta(x_n) \sim \frac{12 \ln 2}{\pi^2} \ln q_n .$$

Therefore, for almost all x

$$\delta\left(\frac{ax_n + b}{cx_n + d}\right) \sim \frac{12 \ln 2}{\pi^2} \ln(cp_n + dq_n) .$$

Now $p_n \sim xq_n$ so that

$$cp_n + dq_n \sim (cx + d)q_n$$

$$\ln(cp_n + dq_n) \sim \ln q_n .$$

Hence for almost all x

$$\delta\left(\frac{ax_n + b}{cx_n + d}\right) \sim \delta(x_n) .$$

By countable intersection, we conclude that for almost all x and for all Möbius map T

$$\delta(Tx_n) \sim \delta(x_n) . \quad \text{QED}$$

PROBLEM 5. Let T be a given Möbius map and let $I(T)$ be the associated interval. Let $\zeta \in I(T)$. To compute the Hausdorff dimension of those x for which

$$\lim_{n \rightarrow \infty} \frac{\delta(Tx_n)}{n} = \zeta.$$

Extend this problem to higher dimensions in the spirit of problem 4.

§6. QUADRATIC SURDS

Let x be a real quadratic number. Its continued fraction expansion is ultimately periodic. Let $\pi(x)$ be its period. H. Cohen [3], followed by J. Cusick [4] and Paysant-Leroux [11] studied the action of a Möbius map on the period. They established that

$$\limsup_{\pi(x) \rightarrow \infty} \frac{\pi(Tx)}{\pi(x)} = R(\Delta)$$

where $R(\Delta)$ is an integer. Furthermore

$$A n \ln n \leq R(n) \leq B n \ln n + 1$$

for some constants $A > 0$, $B > 0$. A simple argument then shows that

$$\liminf_{\pi(x) \rightarrow \infty} \frac{\pi(Tx)}{\pi(x)} = \frac{1}{R(\Delta)}.$$

PROBLEM 6. Is it true that for all real quadratic irrational x

$$\sup_n \pi(x^n) = \infty ?$$

Define the interval

$$J(\Delta) = \left[\frac{1}{R(\Delta)}, R(\Delta) \right].$$

PROBLEM 7. Let $\zeta \in J(\Delta)$. Prove the existence of a sequence of real quadratic numbers x_n with strictly increasing period such that

$$\lim_{n \rightarrow \infty} \frac{\pi(Tx_n)}{\pi(x_n)} = \zeta.$$

Extend this result to higher dimensions as in Problem 4.

PROBLEM 8. *Does there exist a sequence x_n of quadratic numbers with strictly increasing period such that for all Möbius map T*

$$\lim_{n \rightarrow \infty} \frac{\pi(Tx_n)}{\pi(x_n)} = 1 ?$$

We believe some of our problems are relatively easy to solve. But quite obviously Problem 1, 2 and maybe 6 are deep.

REFERENCES

- [1] BILLINGSLEY, P. *Ergodic Theory and Information*. Wiley 1965.
- [2] CHOQUET, G. Répartition des nombres $k(3/2)^n$; et ensembles associés, (English summary) *C.R. Acad. Sci. Paris Sér. A-B* 290 (1980), no. 13, A575-A580; Algorithmes adaptés aux suites $(k\theta^n)$ et aux chaînes associées, (English summary) *C.R. Acad. Sci. Paris Sér. A-B* 290 (1980), no. 16, A719-A724; θ -jeux récurrents et application aux suites $(k\theta^n)$; solénoïdes de T^z , (English summary) *C.R. Acad. Sci. Paris Sér. A-B* 290 (1980), no. 19, A863-A868; Construction effective de suites $(k(3/2)^n)$. Etude des mesures $(3/2)$ -stables, (English summary) *C.R. Acad. Sci. Paris Sér. A-B* 291 (1980), no. 2, A69-A74; Les fermés $(3/2)$ -stables de T ; structure des fermés dénombrables; applications arithmétiques, (English summary) *C.R. Acad. Sci. Paris Sér. A-B* 291 (1980), no. 4, A-239-A244; θ -fermés; θ -chaînes et θ -cycles (pour $\theta = 3/2$), (English summary) *C.R. Acad. Sci. Paris Sér. I Math.* 292 (1981), no. 1, 5-10; θ -fermés et dimension de Hausdorff. Conjectures de travail. Arithmétique des θ -cycles (où $\theta = 3/2$), (English summary) *C.R. Acad. Sci. Paris Sér. I Math.* 292 (1981), no. 6, 339-344.
- [3] COHEN, H. Multiplication par un entier d'une fraction continue périodique. *Acta Arith.* 26 (1974), 129-148.
- [4] CUSICK, T. Integer multiples of periodic continued fractions. *Pacific J. Math.* 78 (1978), 47-60.
- [5] DAUDÉ, H. Thèse, Caen 20 Janv. 1993.
- [6] DIXON, J.D. The number of steps in the Euclidean algorithm. *J. of Number Theory* 2 (1970), 414-422. A simple estimate for the number of steps in the Euclidean algorithm. *Amer. Math. Monthly* 78 (1971), 374-376.
- [7] HEILBRONN, H. On the average length of a class of finite continued fractions. In *Abhandlungen aus Zahlentheorie und Analysis*, Berlin 1968, edited by P. Turàn, 87-96.
- [8] LAMÉ, G. Note sur la limite... *Comptes Rendus Acad. Sci. Paris* 19 (1844), 867-870.
- [9] LÉVY, P. Sur le développement en fraction continue d'un nombre choisi au hasard. *Compositio Mathem.* 3 (1936), 286-303.
- [10] MENDÈS FRANCE, M. Quelques problèmes relatifs à la théorie des fractions continues limitées. *Séminaire de Théorie des Nombres Bordeaux 1971-72*, 4 bis 01-09; Sur les fractions continues limitées, *Acta Arith.* 23 (1973), 207-215; The depth of a rational number, in *Topics in Number Theory, Debrecen 1974, Coll. Mathem. Soc. János Bolyai* 13, 183-194.

- [11] PAYSANT-LEROUX, R. Transformation par une fonction homographique... in *Journées Arith. Caen. SMF Astérisque 41-42*, 1977, 251-253.
- [12] PISOT, C. Répartition (mod 1) des puissances successives des nombres réels. *Commentari. Math. Helv.* 19 (1946-47), 153-160.
- [13] PORTER, J.W. On a theorem of Heilbronn. *Mathematika* 22 (1975), 20-28.
- [14] POURCHET, Y. Private letter to Mendès France, 1972.
- [15] TONKOV, T. On the average length of finite continued fraction. *Acta Arith.* 26 (1974), 47-57.
- [16] VAN DER POORTEN, A. Some problems of recurrent interest. In *Topics in Classical Number Theory, Budapest 1981, Coll. Math. Soc. János Bolyai 34*, edited by G. Halász, vol. 2, 1265-1294.

(Reçu le 13 avril 1993)

Michel Mendès France

Département de Mathématiques
Université de Bordeaux I
F-33405 Talence, France

ADDED IN PROOF

1. In a delightful article to appear "Origins of the Analysis of Algorithms", J. Shallit discusses the early history of $\delta(x)$, $x \in \mathbf{Q}$.

2. Recently (summer 1993), G. Grisel (University of Caen, France) managed to show that for a large class of quadratic irrationals x , $\pi(x^n)$ is indeed unbounded so that Problem 6 is partially solved.

Vide-leer-empty