

## 3.2 The GIT quotient $S^3 \times \mathbb{C}^n // \mathrm{SL}(n, \mathbb{C})$

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.09.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARK 6. It is not difficult to show that  $F^t$  is not injective if and only if there exists a proper quotient  $\bar{H}_C$  of  $H_C$ , and a form  $\bar{F} \in S^3 \bar{H}_C^\vee$  whose pull-back to  $H_C$  is the given form  $F$ . This means that the Hessians of cubic polynomials  $f \in \mathbf{C}[H_C]_3$  which ‘do not depend on all variables’ are automatically zero.

The converse holds for forms in  $b \leq 4$  variables, but not in general [G/N].

### 3.2 THE GIT QUOTIENT $S^3 H_C^\vee //_{SL(H_C)}$

Let  $V := S^3 H_C^\vee$  be the vector space of complex cubic forms. The reductive group  $G := SL(H_C)$  acts rationally on  $V$ , and therefore has a finitely generated ring  $\mathbf{C}[V]^G$  of invariants [H]. The inclusion  $\mathbf{C}[V]^G \subset \mathbf{C}[V]$  induces a regular map  $\pi: V \rightarrow V//_G$  onto the affine variety  $V//_G$  with coordinate ring  $\mathbf{C}[V]^G$ . It is well known that  $\pi$  is a categorical quotient, which is  $G$ -closed and  $G$ -separating, so that  $V//_G$  parametrizes precisely the closed  $G$ -orbits in  $V$ . Recall that a point  $v \in V$  is semi-stable if  $0 \notin \bar{G} \cdot v$ , and that  $v$  is stable if  $G \cdot v$  is closed in  $V$  and the isotropy group  $G_v$  is finite [M/F]. Denote the  $G$ -invariant, open subsets of semistable (stable) points in  $V$  by  $V^{ss}$  ( $V^s$ ).

The complement  $V \setminus V^{ss} = \pi^{-1}(\pi(0))$  consists of ‘Nullformen’, i.e. forms for which all polynomial invariants vanish. The open subset of stable points, which includes in particular all non-singular forms, has a geometric quotient, given by the restricted map  $\pi|_{V^s}: V^s \rightarrow \pi(V^s)$ .

REMARK 7. Let  $A_\circ \in GL(H)$  be a fixed automorphism of determinant  $\det A_\circ = -1$ , e.g.  $A_\circ = -id_H$  if  $b$  is odd.  $A_\circ$  induces a  $\mathbf{Z}/2$ -action on  $S^3 H^\vee /_{SL(H)}$  and on  $S^3 H_C^\vee /_{SL(H_C)}$ , for which the map  $c$  is equivariant.

Let  $\hat{G} \subset GL(H_C)$  be the semi-direct product of  $SL(H_C)$  and  $\mathbf{Z}/2$  generated by  $A_\circ$  and  $SL(H_C)$ . The invariant ring  $\mathbf{C}[V]^{\hat{G}}$  has an important topological interpretation: it consists of all polynomial invariants of complex cohomology rings of 1-connected, closed, oriented 6-dimensional manifolds with torsion-free homology.

#### EXAMPLE 5. Binary cubics ( $b = 2$ )

Choose linear coordinates  $X, Y$  on  $H_C$ , and write a cubic polynomial  $f \in \mathbf{C}[X, Y]_3$  in the form  $f = a_0 X^3 + 3a_1 X^2 Y + 3a_2 X Y^2 + a_3 Y^3$ .

We use  $a_0, a_1, a_2, a_3$  as coordinates on  $S^3 H_C^\vee$ , so that  $\mathbf{C}[S^3 H_C^\vee] = \mathbf{C}[a_0, a_1, a_2, a_3]$ . The discriminant  $\Delta(f)$  of  $f$  is a homogeneous

polynomial of degree 4 in the coefficients  $a_0, a_1, a_2, a_3$ , explicitly given by  $\Delta(f) = a_0^2 a_3^2 - 3a_1^2 a_2^2 - 6a_0 a_1 a_2 a_3 + 4a_0 a_2^3 + 4a_1^3 a_3$ .

The discriminant generates the ring of  $SL(H_C)$ -invariants,

$$\mathbf{C}[S^3 H_C^\vee]^{SL(H_C)} = \mathbf{C}[\Delta],$$

and it is easy to see that  $\Delta$  is also  $\mathbf{Z}/2$ -invariant. A cubic form  $f$  is stable if and only if it is semistable, if and only if it is non-singular [N]. The cone of nullforms  $\pi^{-1}(\pi(0))$  is the affine hypersurface  $(\Delta)_\circ \subset S^3 H_C^\vee$ ; it has a nice geometric interpretation in terms of the Hessian. The Hessian of the cubic  $f$  is the quadratic form

$$H_f = 6^2 [(a_0 a_2 - a_1^2) X^2 + (a_0 a_3 - a_1 a_2) XY + (a_1 a_3 - a_2^2) Y^2].$$

The set of forms  $f$  with vanishing Hessians  $H_f$  form the affine cone over the rational normal curve in  $\mathbf{P}(S^3 H_C^\vee)$ ; the hypersurface of nullforms is the cone over the tangential scroll of this curve. There are 4 different types of  $SL(H_C)$ -orbits in  $S^3 H_C^\vee$ , represented by the normal forms  $XY(X + \lambda Y)$ ,  $X^2 Y$ ,  $X^3$ ,  $0$ . The first type is stable, the others are nullforms, the orbits of  $X^3$  and  $0$  have vanishing Hessians.

#### EXAMPLE 6. Ternary cubics ( $b = 3$ )

The ring of  $SL(H_C)$ -invariants of ternary cubics is a weighted polynomial ring in 2 variables,  $\mathbf{C}[S^3 H_C^\vee]^{SL(H_C)} = \mathbf{C}[S, T]$  whose generators  $S, T$  have been found by S. Aronhold [A].  $S$  is a homogeneous polynomial of degree 4 in the coefficients of a cubic  $f$ ,  $T$  is homogeneous of degree 6, both polynomials are  $\mathbf{Z}/2$ -invariant. For a cubic of the form  $f = aX^3 + bY^3 + cZ^3 + 6dXYZ$ ,  $S$  and  $T$  are given by  $S = 4d(d^3 - abc)$  and  $T = 8d^6 + 20abc(d^3 - abc)$  respectively [P]. The general formulae, which take two pages to write down, can be found in the book of Sturmfels [St]. The discriminant of a form  $f$  is homogeneous of degree 12 in the coefficients of  $f$ ; in terms of Aronhold's invariants  $S, T$  it is simply given by  $\Delta = S^3 - T^2$ . We obtain the following overall picture: The GIT quotient for ternary cubics is an affine plane  $\mathbf{A}^2$  with coordinates  $S, T$ . The complement  $\mathbf{A}^2 \setminus (\Delta)_\circ$  of the discriminant curve is the geometric quotient of stable cubics. The  $\pi$ -fibers over a point  $(S, T) \neq (0, 0)$  on the discriminant curve  $(\Delta)_\circ$  consist of 3 types of  $SL(H_C)$ -orbits: nodal cubics with normal form  $X^3 + Y^3 + 6\alpha XYZ$ , reducible cubics formed by a smooth conic and a transversal line (normal form:  $X^3 + 6\alpha XYZ$ ), and cubics consisting of three lines in general position (normal form:  $6\alpha XYZ$ ); these cubics are properly

semi-stable for  $\alpha \neq 0$  with Aronhold invariants  $S = 4\alpha^4$ ,  $T = 8\alpha^6$ . The fiber of  $\pi$  over 0 contains 6 orbits with normal forms

$$Y^2Z - X^3, Y(X^2 - YZ), XY(X + Y), X^2Y, X^3,$$

and 0, of which the last 4 types have vanishing Hessians. For more details we refer to H. Kraft's book [Kr].

REMARK 8. The natural  $\mathbf{C}^*$ -action  $f \rightarrow \lambda \cdot f$  on cubic forms induces a weighted action on the GIT quotient  $S^3H_{\mathbf{C}}^{\vee}/SL(H_{\mathbf{C}})$ ,  $\lambda \cdot (S, T) = (\lambda^4 S, \lambda^6 T)$ . The associated weighted projective space  $\mathbf{P}^1(4, 6)$  with homogeneous coordinates  $\langle S, T \rangle$  is the good quotient for semi-stable plane cubic curves. Its affine part  $\mathbf{P}^1 \setminus (\Delta)_{\circ}$  is the moduli space of genus-1 curves. The  $PGL(H_{\mathbf{C}})$ -invariant  $J := \frac{S^3}{\Delta}$  gives the  $J$ -invariant of the corresponding curve.

### 3.3 ARITHMETICAL ASPECTS

Let  $c: S^3H^{\vee}/SL(H) \rightarrow S^3H_{\mathbf{C}}^{\vee}/SL(H)$  be the map which associates to the  $SL(H)$ -orbit  $\langle F \rangle$  of a symmetric trilinear form  $F \in S^3H^{\vee}$  the  $SL(H_{\mathbf{C}})$ -orbit  $\langle F \rangle_{\mathbf{C}}$  of its complexification. The  $c$ -fiber over  $\langle F \rangle_{\mathbf{C}}$  can be identified with the subset  $(SL(H_{\mathbf{C}}) \cdot F \cap S^3H^{\vee})/SL(H)$  of  $S^3H^{\vee}/SL(H)$ . C. Jordan has shown that these subsets are finite provided the cubic form  $f \in \mathbf{C}[H_{\mathbf{C}}]_3$  associated to  $F$  has a non-vanishing discriminant [J1]. Jordan's original proof, which is only two pages long, is somewhat hard to follow. The following theorem of A. Borel and Harish-Chandra provides, however, a vast generalization of Jordan's finiteness result:

THEOREM 3 (Borel/Harish-Chandra). *Let  $G$  be a reductive  $\mathbf{Q}$ -group,  $\Gamma \subset G$  an arithmetic subgroup,  $\xi: G \rightarrow GL(V)$  a  $\mathbf{Q}$ -morphism, and  $L \subset V$  a  $\Gamma$ -invariant sublattice of  $V_{\mathbf{Q}}$ . If  $v \in V$  has a closed  $G$ -orbit in  $V$ , then  $G_v \cap L/\Gamma$  is a finite set.*

*Proof.* [B].

COROLLARY 4. *Let  $F \in S^3H^{\vee}$  be a symmetric trilinear form on  $H$ . If the  $SL(H_{\mathbf{C}})$ -orbit of  $F$  in  $S^3H_{\mathbf{C}}^{\vee}$  is closed, then the fiber  $c^{-1}(\langle F \rangle_{\mathbf{C}})$  over  $\langle F \rangle_{\mathbf{C}}$  is finite.*

To check whether a  $SL(H_{\mathbf{C}})$ -orbit  $SL(H_{\mathbf{C}}) \cdot F$  is closed in  $S^3H_{\mathbf{C}}^{\vee}$ , one has a generalization of the Hilbert-criterion [Kr]:  $SL(H_{\mathbf{C}}) \cdot F$  is closed in  $S^3H_{\mathbf{C}}^{\vee}$  if and only if for every 1-parameter subgroup  $\lambda: \mathbf{C}^* \rightarrow SL(H_{\mathbf{C}})$ , for