

2. The index of an elliptic complex

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **19.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. THE INDEX OF AN ELLIPTIC COMPLEX

An *elliptic complex* (E, d) over a closed, oriented, n dimensional Riemannian manifold M consists of:

- a) a finite collection of finite dimensional complex vector bundles

$$E_0, E_1, \dots, E_k$$

- b) a collection of smooth differential operators

$$d_i : C^\infty(E_i) \rightarrow C^\infty(E_{i+1})$$

- c) The operators d_i are required to satisfy

$$d_{i+1} \cdot d_i = 0$$

and an additional technical condition called ellipticity. This condition makes possible the construction a virtual bundle, i.e. the formal difference of two vector bundles, over TM which carries a great deal of information about (E, d) . This virtual bundle $\sigma(E, d)$ is called the symbol of (E, d) and it defines a class $[\sigma(E, d)]$, also called the symbol, in the K theory with compact supports of TM .

EXAMPLES

1. The de Rham complex, where

$T_{\mathbb{C}}^*M$ = complexified cotangent bundle of M

$E_i = \Lambda^i T_{\mathbb{C}}^*M$ the i th exterior power of $T_{\mathbb{C}}^*M$

$C^\infty(E_i)$ = smooth complex i forms on M

d_i = the usual exterior derivative

2. The Dolbeault complex
 3. The Signature complex (see [AS])
 4. The twisted Spin complex.

SOME FACTS ABOUT ELLIPTIC COMPLEXES

Set $H^i(E, d) = \ker d_i / \text{image } d_{i-1}$. If M is compact, then $\dim H^i(E, d) < \infty$, and we may define

$$\text{Index}(E, d) = \sum_{i=0}^k (-1)^i \dim H^i(E, d).$$

This is a very important invariant. Special cases of (E, d) yield the

1. Euler class $\chi(M)$ of M (de Rham complex)
2. Signature of M (Signature complex)
3. Euler class $\chi(M, V)$ (Dolbeault complex)
4. $\hat{A}$ genus of M (Spin complex).

The Atiyah-Singer Index Theorem tells how to compute this invariant from topological information about M and (E, d) . In particular, it says

THEOREM 2.1 ([AS]).

$$\text{Index}(E, d) = \int_M Td(TM \otimes_{\mathbf{R}} \mathbf{C}) \cdot \text{ch}(\sigma(E, d)).$$

The theorems quoted above are all special cases of this theorem. We now give an idea of how to prove this deep and important theorem.

On each E_i choose an Hermitian inner product denoted $(\ , \)_i$. This induces an inner product $\langle \ , \ \rangle_i$ on $C^\infty(E_i)$ by the formula

$$\langle s_1, s_2 \rangle_i = \int_M (s_1(x), s_2(x))_i dx.$$

Using $\langle \ , \ \rangle_i$ we define the adjoints

$$d_i^* : C^\infty(E_i) \rightarrow C^\infty(E_{i-1})$$

by

$$\langle s_1, d_i^* s_2 \rangle_{i-1} = \langle d_{i-1} s_1, s_2 \rangle_i$$

where

$$s_1 \in C^\infty(E_{i-1}), \quad s_2 \in C^\infty(E_i).$$

The Laplacian $\Delta_i : C^\infty(E_i) \rightarrow C^\infty(E_i)$ is defined by

$$\Delta_i = d_{i-1} d_i^* + d_{i+1}^* d_i,$$

and it extends to a densely defined operator of $L^2(E_i)$, the space of L^2 sections of E_i , as follows. Δ_i is a diagonalizable operator, and any eigenvalue λ of Δ_i must be real and nonnegative. If M is compact there is a sequence of real numbers

$$0 = \lambda_0 < \lambda_1 < \lambda_2 < \cdots, \quad \lim_{j \rightarrow \infty} \lambda_j = \infty$$

such that for each E_i there is a sequence of finite dimensional subspaces of $C^\infty(E_i)$, denoted

$$E_i(\lambda_0), E_i(\lambda_1), E_i(\lambda_2), \dots$$

so that for any $s \in E_i(\lambda_j)$

$$\Delta_i s = \lambda_j s.$$

In addition

$$L^2(E_i) = \bigoplus_{j=0}^{\infty} E_i(\lambda_j).$$

Thus each element in $L^2(E_i)$ can be written as a (possibly infinite) sum of eigen functions and we may think of Δ_i as the infinite diagonal matrix

$$\begin{bmatrix} 0 & & & & & & \\ & \ddots & & & & & \\ & & 0 & & & & \\ & & & \lambda_1 & & & \\ & & & & \ddots & & \\ & & & & & \lambda_1 & \\ & & & & & & \lambda_2 \\ & & & & & & & \ddots \\ & & & & & & & & \lambda_2 \\ & & & & & & & & & \ddots \end{bmatrix}.$$

OTHER PROPERTIES OF Δ_i

- 1) $E_i(\lambda_0) = \ker \Delta_i \subset \ker d_i$ and the inclusion of $E_i(\lambda_0)$ in $\ker d_i$ induces an isomorphism

$$E_i(\lambda_0) \simeq H^i(E, d),$$

so

$$\text{Index}(E, d) = \sum_{i=0}^k (-1)^i \dim E_i(\lambda_0).$$

2) For each $\lambda_j > 0$, the sequence

$$0 \rightarrow E_0(\lambda_j) \xrightarrow{d_0} E_1(\lambda_j) \xrightarrow{d_1} \cdots \xrightarrow{d_{k-1}} E_k(\lambda_j) \rightarrow 0$$

is exact.

As a corollary, we have immediately

$$\sum_{i=0}^k (-1)^i \dim E_i(\lambda_j) = 0$$

for all $\lambda_j > 0$. These results rely on the fact that M is compact. For a general reference for the above facts, see [Wa].

The fact that Δ_i is diagonal implies that for any function $f : \mathbf{R} \rightarrow \mathbf{R}$, we may define

$$f(\Delta_i) : L^2(E_i) \rightarrow L^2(E_i)$$

by : for each $s \in E_i(\lambda_j)$ set $f(\Delta_i)s = f(\lambda_j)s$. i.e. the “matrix” of $f(\Delta_i)$ is

$$\begin{bmatrix} f(0) & & & & & \\ & \ddots & & & & \\ & & f(0) & & & \\ & & & f(\lambda_1) & & \\ & & & & \ddots & \\ & & & & & f(\lambda_1) \\ & & & & & & \ddots \end{bmatrix}$$

Note also that if $f(x)$ goes to zero rapidly enough as $x \rightarrow \infty$, then the trace of $f(\Delta_i)$, thought of as the usual trace applied to the infinite matrix above, will be a finite number. In this case, we say $f(\Delta_i)$ is of trace class. See [RS].

We are interested in the family of functions $f_t(x) = e^{-tx}$, $t > 0$. In fact, even if M is not compact, $e^{-t\Delta}$ makes sense and we have

THEOREM 2.2 (Seeley, [S]). *For $t > 0$, $e^{-t\Delta_i}$ is a smoothing operator on $L^2(E_i)$ and so if M is compact it is of trace class.*

Let $\pi_j : M \times M \rightarrow M$ be projection on the j th factor, $j = 1, 2$. To say an operator A on $L^2(E_i)$ is a smoothing operator means that there is a smooth section $k(x, y)$ of the bundle $\text{Hom}(\pi_2^* E_i, \pi_1^* E_i)$ over $M \times M$, so that for all $s \in L^2(E_i)$.

$$(As)(x) = \int_M k(x, y)s(y)dy.$$

Note that $k(x, y)$ is a linear map from $E_{i,y}$, the fiber over y , to $E_{i,x}$, the fiber over x , so $k(x, x) : E_{i,x} \rightarrow E_{i,x}$ has a well defined trace. The section $k(x, y)$ is called the Schwartz kernel of A . Any smoothing operator on a compact manifold is of trace class and its trace is given by $\text{tr}(A) = \int_M \text{tr } k(x, x)dx$.

To see this for $e^{-t\Delta_i}$, note that its Schwartz kernel $k_t^i(x, y)$ must be given as follows: For each λ_j choose an orthonormal basis ϕ_j^v , $v = 1, \dots, \dim E_i(\lambda_j)$ of $E_i(\lambda_j)$. Then

$$k_t^i(x, y) = \sum_{j=0}^{\infty} e^{-t\lambda_j} \left[\sum_v \phi_j^v(x) \phi_j^v(y) \right].$$

Here $k_t^i(x, y) : E_{i,y} \rightarrow E_{i,x}$ acts on $w \in E_{i,y}$ by

$$k_t^i(x, y)w = \sum_{j=0}^{\infty} e^{-t\lambda_j} \left[\sum_v (\phi_j^v(y), w)_i \cdot \phi_j^v(x) \right].$$

The trace of $k_t^i(x, x)$ is then given by

$$\sum_{j=0}^{\infty} e^{-t\lambda_j} \left[\sum_v (\phi_j^v(x), \phi_j^v(x))_i \right]$$

and the result follows by integrating over M .

Now, since $e^{-t\lambda_0} = 1$ for all t , we have $e^{-t\lambda_0} \sum_{i=0}^k (-1)^i \dim E_i(\lambda_0) =$

$\text{Index}(E, d)$, for all t . In addition $e^{-t\lambda_j} \sum_{i=0}^k (-1)^i \dim E_i(\lambda_j) = 0$ for $j > 0$, and for all t . Thus we have

THEOREM 2.3. *If M is compact, then for all $t > 0$,*

$$\begin{aligned} \text{Index}(E, d) &= \sum_{j=0}^{\infty} \left[\sum_{i=0}^k (-1)^i e^{-t\lambda_j} \dim E_i(\lambda_j) \right] \\ &= \sum_{i=0}^k \left[\sum_{j=0}^{\infty} (-1)^i e^{-t\lambda_j} \dim E_i(\lambda_j) \right] \\ &= \sum_{i=0}^k (-1)^i \text{tr } e^{-t\Delta_i}. \end{aligned}$$

The Index Theorem now follows from two other results.

- 1) Set $k_t(x) = \sum_{i=0}^k (-1)^i \text{tr } k_t^i(x, x)$. Then for t near 0, $k_t(x)$ has an asymptotic expansion of the form

$$k_t(x) = \sum_{j \geq -n} t^{j/2} a_j(x).$$

As $\int_M k_t(x) dx = \sum_{i=0}^k (-1)^i \text{tr } e^{-t\Delta_i} = \text{Index}(E, d)$ is independent of t , we have

$$\text{Index}(E, d) = \int_M a_0(x) dx.$$

Now, for any twisted Dirac operator D_F^+ , one can prove that the differential n form $a_0(x)dx$ is the degree n part of the form constructed from the connections on TM and F which represents $\hat{A}(TM) \cdot ch(F)$. Thus, we have

$$\text{Index}(D_F^+) = \int_M \hat{A}(TM) \cdot ch(F).$$

- 2) $\text{Index}(E, d)$ depends only on the K theory class $[\sigma(E, d)]$. Given this and the formula above for $\text{Index}(D_F^+)$, one may use well known arguments in K theory to extend the result in 1. to all elliptic complexes. The essential fact is that the symbols of twisted Dirac operators generate the K theory with compact supports of TM as an algebra over the K theory of M .

The difference between the formula in 1. and that in the Atiyah-Singer Index Theorem is accounted for by the fact that for the twisted Spin complex $(E^\pm \otimes F, D_F^+)$,

$$ch(\sigma(E^\pm \otimes F, D_F^+)) = ch(E^\pm, D^+) \cdot ch(F)$$

and

$$Td(TM \otimes_{\mathbf{R}} \mathbf{C}) \cdot \mathbf{ch}(\sigma(E^\pm, D^+)) = \hat{A}(TM).$$

For more on this see [ABP], [B], [G], [Gi], and [P].