

5.1 Big and Small Groups

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **22.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

In what follows, Γ will denote a finitely-presented group. Given a presentation of Γ , there is an associated 2-dimensional CW-complex K which we call the *presentation complex*. To form it, make a bouquet of circles indexed by the generators of Γ . Attach 2-cells based on the relations of Γ . (We allow trivial or repeated relations in the presentation.) This is the presentation complex.

DEFINITION 7. Put $b_0^{(2)}(\Gamma) = b_0^{(2)}(K)$, $b_1^{(2)}(\Gamma) = b_1^{(2)}(K)$, $\alpha_1(\Gamma) = \alpha_1(K)$ and $\alpha_2(\Gamma) = \alpha_2(K)$.

By Property 4 above, Definition 7 makes sense in that the choice of presentation of Γ does not matter.

As the invariants $b_0^{(2)}(\Gamma)$, $b_1^{(2)}(\Gamma)$, $\alpha_1(\Gamma)$ and $\alpha_2(\Gamma)$ will play an important role, let us state explicitly what they measure. First, from Property 5, $b_0^{(2)}(\Gamma)$ tells us whether or not Γ is infinite. In general, $b_0^{(2)}(\Gamma) = \frac{1}{|\Gamma|}$. Next, from Property 1, $b_1^{(2)}(\Gamma)$ tells us whether or not M has square-integrable harmonic 1-forms (or \tilde{K} has square-integrable harmonic 1-cochains). From Property 2, $\alpha_1(\Gamma)$ tells us whether or not the Laplacian Δ_0 , acting on functions on M , has a gap in its spectrum away from zero. In fact, Property 6 is just a restatement of Corollary 3. Finally, from Property 2, $\alpha_2(\Gamma)$ tells us whether or not the spectrum of the Laplacian on $\Lambda^1(M)/\text{Ker}(d)$ goes down to zero.

5.1 BIG AND SMALL GROUPS

Let us first introduce a convenient terminology for the purposes of the present paper.

DEFINITION 8. *The group Γ is big if it is nonamenable, $b_1^{(2)}(\Gamma) = 0$ and $\alpha_2(\Gamma) = \infty^+$. Otherwise, Γ is small.*

We recall that Δ_p denotes the Laplace-Beltrami operator on the universal cover M .

PROPOSITION 11. *Let X and M be as above. The group $\pi_1(X)$ is small if and only if $0 \in \sigma(\Delta_0)$ or $0 \in \sigma(\Delta_1)$.*

Proof. This follows immediately from Properties 1, 2, 4, 5 and 6 above. \square

The question arises as to which groups are big and which are small. Clearly any amenable group is small.

PROPOSITION 12. *Fundamental groups of compact surfaces are small.*

Proof. Suppose that Σ is a compact surface and $\Gamma = \pi_1(\Sigma)$. If Σ has boundary then Γ is a free group F_j on some number j of generators. If $j = 0$ or $j = 1$ then Γ is amenable. If $j > 1$ then $b_1^{(2)}(\Gamma) = j - 1 > 0$.

Suppose now that Σ is closed. If $\chi(\Sigma) \geq 0$ then Γ is amenable. If $\chi(\Sigma) < 0$ then $b_1^{(2)}(\Gamma) = -\chi(\Sigma) > 0$. \square

We now extend Proposition 12 to 3-manifold groups. We use some facts about compact connected 3-manifolds Y , possibly with boundary. (See, for example, [21, Section 6]). Again, all of our manifolds are assumed to be oriented. First, Y has a decomposition as a connected sum $Y = Y_1 \# Y_2 \# \dots \# Y_r$ of *prime* 3-manifolds. A prime 3-manifold is *exceptional* if it is closed and no finite cover of it is homotopy-equivalent to a Seifert, Haken or hyperbolic 3-manifold. No exceptional prime 3-manifolds are known and it is likely that there are none.

PROPOSITION 13 (Lott-Lück). *Suppose that Y is a compact connected oriented 3-manifold, possibly with boundary, none of whose prime factors are exceptional. Then $\pi_1(Y)$ is small.*

Proof. We argue by contradiction. Suppose that $\pi_1(Y)$ is big. First, $\pi_1(Y)$ must be infinite. If ∂Y has any connected components which are 2-spheres then we can cap them off with 3-balls without changing $\pi_1(Y)$. So we can assume that ∂Y does not have any 2-sphere components. In particular, $\chi(Y) = \frac{1}{2}\chi(\partial Y) \leq 0$. From [21, Theorem 0.1.1],

$$(5.3) \quad b_1^{(2)}(Y) = (r - 1) - \sum_{i=1}^r \frac{1}{|\pi_1(Y_i)|} - \chi(Y).$$

As this must vanish, we have $\chi(Y) = 0$ and either

1. $\{|\pi_1(Y_i)|\}_{i=1}^r = \{2, 2, 1, \dots, 1\}$ or
2. $\{|\pi_1(Y_i)|\}_{i=1}^r = \{\infty, 1, \dots, 1\}$.

It follows that ∂Y is empty or a disjoint union of 2-tori. As there are no 2-spheres in ∂Y , if $|\pi_1(Y_i)| = 1$ then Y_i is a homotopy 3-sphere. Thus Y is homotopy-equivalent to either

1. $\mathbf{RP}^3 \# \mathbf{RP}^3$ or
2. A prime 3-manifold Y' with infinite fundamental group whose boundary is empty or a disjoint union of 2-tori.

If Y is homotopy-equivalent to $\mathbf{R}P^3\#\mathbf{R}P^3$ then $\pi_1(Y)$ is amenable, which is a contradiction. So we must be in the second case. Using Property 3, we may assume that $Y = Y'$. Then as Y is prime, it follows from [24, Chapter 1] that either $Y = S^1 \times D^2$ or Y has incompressible (or empty) boundary. If $Y = S^1 \times D^2$ then $\pi_1(Y)$ is amenable. If Y has incompressible (or empty) boundary then from [21, Theorem 0.1.5], $\alpha_2(Y) \leq 2$ unless Y is a closed 3-manifold with an \mathbf{R}^3 , $\mathbf{R} \times S^2$ or Sol geometric structure. In the latter cases, Γ is amenable. Thus in any case, we get a contradiction. \square

The next proposition gives examples of big groups.

PROPOSITION 14.

1. *A product of two nonamenable groups is big.*
2. *If Y is a closed nonpositively-curved locally symmetric space of dimension greater than three, with no Euclidean factors in \tilde{Y} , then $\pi_1(Y)$ is big.*

Proof. 1. Suppose that $\Gamma = \Gamma_1 \times \Gamma_2$ with Γ_1 and Γ_2 nonamenable. Then Γ is nonamenable. Let K_1 and K_2 be presentation complexes with fundamental groups Γ_1 and Γ_2 , respectively. Put $K = K_1 \times K_2$. Then $\Gamma = \pi_1(K)$. Let $\Delta_p(\tilde{K})$, $\Delta_p(\tilde{K}_1)$ and $\Delta_p(\tilde{K}_2)$ denote the Laplace-Beltrami operator on p -cochains on \tilde{K} , \tilde{K}_1 and \tilde{K}_2 , respectively, as defined in Subsection 5.2 below. Then

$$(5.4) \quad \inf(\sigma(\Delta_1(\tilde{K}))) = \min(\inf(\sigma(\Delta_1(\tilde{K}_1))) + \inf(\sigma(\Delta_0(\tilde{K}_2))), \inf(\sigma(\Delta_0(\tilde{K}_1))) + \inf(\sigma(\Delta_1(\tilde{K}_2)))) > 0.$$

Using Proposition 11, the first part of the proposition follows.

2. If \tilde{Y} is irreducible then part 2. of the proposition follows from the second remark after Proposition 7. If \tilde{Y} is reducible then we can use an argument similar to (5.4). \square

REMARK. Let Γ be an infinite finitely-presented discrete group with Kazhdan's property T. From [6, p. 47], $H^1(\Gamma; \ell^2(\Gamma)) = 0$. This implies that Γ is nonamenable and $b_1^{(2)}(\Gamma) = 0$. We do not know if it is necessarily true that $\alpha_2(\Gamma) = \infty^+$.

5.2 TWO AND THREE DIMENSIONS

In this subsection we relate the zero-in-the-spectrum question to a question in combinatorial group theory. Let K be a finite connected 2-dimensional