

## 5.4 More Dimensions

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.09.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

- a.  $b_0^{(2)} \neq 0 : S^4, S^2 \times S^2, CP^2.$
- b.  $0 \in \sigma(\Delta_0 \text{ on } \Lambda^0 / \text{Ker}(d)) : \mathbf{R}^4, S^3 \times \mathbf{R}, S^2 \times \mathbf{R}^2, Nil^3 \times \mathbf{R}, Nil^4, Sol_0^4, Sol_1^4, Sol_{m,n}^4.$
- c.  $b_1^{(2)} \neq 0 : S^2 \times H^2.$
- d.  $0 \in \sigma(\Delta_1 \text{ on } \Lambda^1 / \text{Ker}(d)) : H^3 \times \mathbf{R}, \widetilde{SL}_2 \times \mathbf{R}, H^2 \times \mathbf{R}^2.$
- e.  $\chi > 0 : H^4, H^2 \times H^2, CH^2.$

Part 2. of the proposition follows.

- 3. Suppose that zero is not in the spectrum of  $\widetilde{X}$ . From Properties 7 and 9,  $\chi(X) = \tau(X) = 0$ . From the classification of complex surfaces,  $X$  has a geometric structure [32, p. 148–149]. This contradicts part 2. of the proposition.  $\square$

#### 5.4 MORE DIMENSIONS

In this subsection we give some partial positive results about the zero-in-the-spectrum question for covers of compact manifolds of arbitrary dimension. Let us first recall some facts about index theory [18]. Let  $X$  be a closed Riemannian manifold. If  $\dim(X)$  is even, consider the operator  $d + d^*$  on  $\Lambda^*(X)$ . Give  $\Lambda^*(X)$  the  $\mathbf{Z}_2$ -grading coming from (3.12). Then the signature  $\tau(X)$  equals the index of  $d + d^*$ . To say this in a more complicated way, the operator  $d + d^*$  defines a element  $[d + d^*]$  of the K-homology group  $K_0(X)$ . Let  $\eta : X \rightarrow \text{pt.}$  be the (only) map from  $X$  to a point. Then  $\eta_*([d + d^*]) \in K_0(\text{pt.})$ . There is a map  $A : K_0(\text{pt.}) \rightarrow K_0(\mathbf{C})$  which is the identity, as both sides are  $\mathbf{Z}$ . So we can say that  $\tau(X) = A(\eta_*([d + d^*])) \in K_0(\mathbf{C})$ .

We now extend the preceding remarks to the case of a group action. Let  $M$  be a normal cover of  $X$  with covering group  $\Gamma$ . The fiber bundle  $M \rightarrow X$  is classified by a map  $\nu : X \rightarrow B\Gamma$ , defined up to homotopy. Let  $\widetilde{d}$  be exterior differentiation on  $M$ . Consider the operator  $\widetilde{d} + \widetilde{d}^*$ . Taking into account the action of  $\Gamma$  on  $M$ , one can define a refined index  $\text{ind}(\widetilde{d} + \widetilde{d}^*) \in K_0(C_r^*\Gamma)$ , where  $C_r^*\Gamma$  is the reduced group  $C^*$ -algebra of  $\Gamma$ .

We recall the statement of the Strong Novikov Conjecture (SNC) [18, 29]. This is a conjecture about a countable discrete group  $\Gamma$ , namely that the assembly map  $A : K_*(B\Gamma) \rightarrow K_*(C_r^*\Gamma)$  is rationally injective. Many groups of a geometric origin, such as discrete subgroups of connected Lie groups or Gromov-hyperbolic groups, are known to satisfy SNC. There are no known groups which do not satisfy SNC.

PROPOSITION 19. *Let  $X$  be a closed Riemannian manifold with a surjective homomorphism  $\pi_1(X) \rightarrow \Gamma$ . Let  $M$  be the induced normal  $\Gamma$ -cover of  $X$ . Suppose that  $\Gamma$  satisfies SNC. Let  $L(X) \in H^*(X; \mathbf{C})$  be the Hirzebruch  $L$ -class of  $X$  and let  $*L(X) \in H_*(X; \mathbf{C})$  be its Poincaré dual. Then if  $\nu_*( *L(X)) \neq 0$  in  $H_*(B\Gamma; \mathbf{C})$ , zero lies in the spectrum of  $M$ . In fact,  $0 \in \sigma \left( \Delta_{\frac{\dim(X)}{2}} \right)$  if  $\dim(X)$  is even and  $0 \in \sigma \left( \Delta_{\frac{\dim(X) \pm 1}{2}} \right)$  if  $\dim(X)$  is odd.*

*Proof.* Suppose first that  $\dim(X)$  is even. Suppose that zero does not lie in the spectrum of  $M$ . Then the operator  $\tilde{d} + \tilde{d}^*$  is invertible. (More precisely, it is invertible as an operator on a Hilbert  $C_r^*\Gamma$ -module of differential forms on  $M$ .) This implies that  $\text{ind}(\tilde{d} + \tilde{d}^*)$  vanishes in  $K_0(C_r^*\Gamma)$ .

The higher index theorem says that

$$(5.10) \quad \text{ind}(\tilde{d} + \tilde{d}^*) = A(\nu_*([d + d^*])).$$

Let  $A_{\mathbf{C}} : K_0(B\Gamma) \otimes \mathbf{C} \rightarrow K_0(C_r^*\Gamma) \otimes \mathbf{C}$  be the complexified assembly map. Using the isomorphism  $K_0(B\Gamma) \otimes \mathbf{C} \cong H_{\text{even}}(B\Gamma; \mathbf{C})$ , the higher index theorem implies that in  $K_0(C_r^*\Gamma) \otimes \mathbf{C}$ ,

$$(5.11) \quad \text{ind}(\tilde{d} + \tilde{d}^*)_{\mathbf{C}} = A_{\mathbf{C}}(\nu_*( *L(X))).$$

By assumption,  $A_{\mathbf{C}}$  is injective. This gives a contradiction.

Let  $T$  be the operator obtained by restricting  $\tilde{d} + \tilde{d}^*$  to

$$\Lambda^{\frac{\dim(X)}{2}}(M) \oplus \overline{\tilde{d}\Lambda^{\frac{\dim(X)}{2}}(M)} \oplus \overline{*d\Lambda^{\frac{\dim(X)}{2}}(M)}.$$

One can show that the other differential forms on  $M$  cancel out when computing the rational index of  $\tilde{d} + \tilde{d}^*$ , so  $T$  will have the same index as  $\tilde{d} + \tilde{d}^*$ . Then the same arguments apply to  $T$  to give  $0 \in \sigma \left( \Delta_{\frac{\dim(X)}{2}} \right)$ .

If  $\dim(X)$  is odd, consider the even-dimensional manifold  $X' = X \times S^1$  and the group  $\Gamma' = \Gamma \times \mathbf{Z}$ . As the proposition holds for  $X'$ , it must also hold for  $X$ .  $\square$

COROLLARY 4. *Let  $X$  be a closed Riemannian manifold. Let  $[X] \in H_{\dim(X)}(X; \mathbf{C})$  be its fundamental class. Suppose that there is a surjective homomorphism  $\pi_1(X) \rightarrow \Gamma$  such that  $\Gamma$  satisfies SNC and the composite map  $X \rightarrow B\pi_1(X) \rightarrow B\Gamma$  sends  $[X]$  to a nonzero element of  $H_{\dim(X)}(B\Gamma; \mathbf{C})$ . Let  $M$  be the induced normal  $\Gamma$ -cover of  $X$ . Then on  $M$ ,  $0 \in \sigma \left( \Delta_{\frac{\dim(X)}{2}} \right)$  if  $\dim(X)$  is even and  $0 \in \sigma \left( \Delta_{\frac{\dim(X) \pm 1}{2}} \right)$  if  $\dim(X)$  is odd.*

*Proof.* As the Hirzebruch  $L$ -class starts out as  $L(X) = 1 + \dots$ , its Poincaré dual is of the form  $*L(X) = \dots + [X]$ . The corollary follows from Proposition 19.  $\square$

**COROLLARY 5.** *Let  $X$  be a closed aspherical Riemannian manifold whose fundamental group satisfies SNC. Then on  $\tilde{X}$ ,  $0 \in \sigma\left(\Delta_{\frac{\dim(X)}{2}}\right)$  if  $\dim(X)$  is even and  $0 \in \sigma\left(\Delta_{\frac{\dim(X)\pm 1}{2}}\right)$  if  $\dim(X)$  is odd.*

*Proof.* This follows from Corollary 4.  $\square$

#### EXAMPLES.

1. If  $X = T^n$  then Corollary 5 is consistent with Example 2 of Section 2.
2. If  $X$  is a compact quotient of  $H^{2n}$  then Corollary 5 is consistent with Example 3 of Section 2.
3. If  $X$  is a compact quotient of  $H^{2n+1}$  then Corollary 5 is consistent with Example 4 of Section 2.
4. If  $X$  is a closed nonpositively-curved locally symmetric space then Corollary 5 is consistent with the second remark after Proposition 7.

If  $X$  is a closed aspherical manifold, it is known that SNC implies that the rational Pontryagin classes of  $X$  are homotopy-invariants [18] and that  $X$  does not admit a Riemannian metric of positive scalar curvature [29]. Thus we see that these three questions about aspherical manifolds, namely homotopy-invariance of rational Pontryagin classes, (non)existence of positive-scalar-curvature metrics and the zero-in-the-spectrum question, are roughly all on the same footing.

If  $X$  is a closed aspherical Riemannian manifold, one can ask for which  $p$  one has  $0 \in \sigma(\Delta_p)$  on  $\tilde{X}$ . The case of locally symmetric spaces is covered by the second remark after Proposition 7. Another interesting class of aspherical manifolds consists of those with amenable fundamental group. By [5],  $\text{Ker}(\Delta_p) = 0$  for all  $p$ . By Corollary 3,  $0 \in \sigma(\Delta_0)$ .

**PROPOSITION 20.** *If  $X$  is a closed aspherical manifold such that  $\pi_1(X)$  has a nilpotent subgroup of finite index then  $0 \in \sigma(\Delta_p)$  on  $\tilde{X}$  for all  $p \in [0, \dim(X)]$ .*

*Proof.* First,  $X$  is homotopy-equivalent to an infranilmanifold, that is, a quotient of the form  $\Gamma \backslash G / K$  where  $K$  is a finite group,  $G$  is the

semidirect product of  $K$  and a connected simply-connected nilpotent Lie group and  $\Gamma$  is a discrete cocompact subgroup of  $G$  [12, Theorem 6.4]. We may as well assume that  $X = \Gamma \backslash G/K$ . By passing to a finite cover, we may assume that  $K$  is trivial. That is,  $X$  is a nilmanifold. From [27, Corollary 7.28],  $H^p(X; \mathbf{C}) \cong H^p(\mathfrak{g}, \mathbf{C})$ , the Lie algebra cohomology of  $\mathfrak{g}$ . From [7],  $H^p(\mathfrak{g}, \mathbf{C}) \neq 0$  for all  $p \in [0, \dim(X)]$ . Thus for all  $p \in [0, \dim(X)]$ ,  $H^p(X; \mathbf{C}) \neq 0$ .

Now let  $\omega$  be a nonzero harmonic  $p$ -form on  $X$ . Let  $\pi^*\omega$  be its pullback to  $\tilde{X}$ . The idea is to construct low-energy square-integrable  $p$ -forms on  $X$  by multiplying  $\pi^*\omega$  by appropriate functions on  $X$ . We define the functions as in [2, §2]. Take a smooth triangulation of  $X$  and choose a fundamental domain  $F$  for the lifted triangulation of  $\tilde{X}$ . If  $E$  is a finite subset of  $\pi_1(X)$ , let  $\chi_H$  be the characteristic function of  $H = \bigcup_{g \in E} g \cdot F$ . Given numbers  $0 < \epsilon_1 < \epsilon_2 < 1$ , choose a nonincreasing function  $\psi \in C_0^\infty([0, \infty))$  which is identically one on  $[0, \epsilon_1]$  and identically zero on  $[\epsilon_2, \infty)$ . Define a compactly-supported function  $f_E$  on  $\tilde{X}$  by  $f_E(m) = \psi(d(m, H))$ . Then there is a constant  $C_1 > 0$ , independent of  $E$ , such that

$$(5.12) \quad \int_{\tilde{X}} |df_E|^2 \leq C_1 \text{area}(\partial H).$$

Define  $\rho_E \in \Lambda^p(\tilde{X})$  by  $\rho_E = f_E \cdot \pi^*\omega$ . We have  $d\rho_E = df_E \wedge \pi^*\omega$  and  $d^*\rho_E = -i(df_E)\pi^*\omega$ . As  $f_E$  is identically one on  $H$ , it follows that there is a constant  $C > 0$ , independent of  $E$ , such that

$$(5.13) \quad \frac{\int_{\tilde{X}} [ |d\rho_E|^2 + |d^*\rho_E|^2 ]}{\int_{\tilde{X}} |\rho_E|^2} \leq C \frac{\text{area}(\partial H)}{\text{vol}(H)}.$$

As  $\pi_1(X)$  is amenable, by an appropriate choice of  $E$  this ratio can be made arbitrarily small. Thus  $0 \in \sigma(\Delta_p)$ .  $\square$

QUESTION. Does the conclusion of Proposition 20 hold if we only assume that  $\pi_1(X)$  is amenable?

## 6. TOPOLOGICALLY TAME MANIFOLDS

Another class of manifolds for which one can hope to get some nontrivial results about the zero-in-the-spectrum question is given by *topologically tame* manifolds, meaning manifolds  $M$  which are diffeomorphic to the interior of a compact manifold  $N$  with boundary. If  $M$  has finite volume then  $\text{Ker}(\Delta_0) \neq 0$ ,