

6. Further Applications

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The t -sequence t^n is interesting because the adjunction problem is already proved (without the torsion-free hypothesis) for words with this t -shape [L]. However the methods discussed here do not extend this result to t -sequences in normal form based on t^n . An example is $t^3 t^{-1} t^3 t^{-1}$.

Another interesting case is the sequence tt^{-1} which is not amenable. However a simple trick (substitute u^2 for t) makes it suitable. Hence theorem 5.1 implies a solution to the adjunction problem (over torsion-free groups) for words of the form $gtg't^{-1}$. For words of this shape, torsion-free is a necessary condition as the example in the introduction shows!

We do not yet have a simple test for amenability though it is easy from the definition to write down large classes of amenable sequences. However it can be seen that, speaking very roughly, a sequence is amenable unless it has a uniform slope, like $t^5 t^{-3} t^5 t^{-3}$ or $t^3 t^{-3} t^3 t^{-3}$ (slope zero).

6. FURTHER APPLICATIONS

We give here the other applications from [Kl] of the crash theorems, not covered above.

THEOREM 6.1 (Application to free products). *Let A, B be groups and suppose each (cyclic) factor of $u \in A * B - A$ has infinite order. Then the natural homomorphism $A \rightarrow \langle A * B \mid [A, u] = 1 \rangle$ is injective.*

Proof. Suppose not. Then the conditions of the first transversality lemma apply and there is a non trivial element $a \in A$ such that $a \in \langle\langle [A, u] \rangle\rangle$. So we have a cell subdivision K of the 2-sphere such that reading round from the base point $*$ for every 2-cell in K spells out the word

$$w(a) = (c_0^{-1} a c_0) c_1 \cdots c_{n-1} (c_n^{-1} a^{-1} c_n) c_{n-1}^{-1} \cdots c_1^{-1}$$

for some $a \in A$, see figure 8. Note that if this 2-cell has the opposite orientation then the word spelt out is $w(a^{-1})$.




FIGURE 8
The 2-cell labelled by $w(a)$

Now consider the traffic flow defined as follows. The car associated to a 2-cell starts out from the base point $*$ and proceeds in an anticlockwise manner so that it takes a unit amount of time to reach the next corner. It is clear that any crash must take place at a 0-cell. By the crash theorem there are at least two total crashes so we can assume it takes place at a 0-cell where the angle labels multiply to 1. There are two cases to consider.

If the crash occurs at time $0 \bmod n$ then the clockwise labelling around this 0-cell is $c_\alpha^{-1} a_1 c_\alpha, c_\alpha^{-1} a_2 c_\alpha, \dots, c_\alpha^{-1} a_k c_\alpha$ where $a_1 a_2 \cdots a_k = 1$ and $\alpha = 0$ or $\alpha = n$. A simple calculation shows that the anticlockwise product of the remaining angles of these k 2-cells is 1. So we may simplify the situation by collapsing these k 2-cells to a point.

If the crash occurs at a time $\neq 0 \bmod n$ then the clockwise labelling around this 0-cell is $c_i^k = 1$ for some $0 < i < n$ and some $k > 1$ contradicting the torsion free hypothesis. $\square$

Let H, H' be groups and let $\phi: H \rightarrow H'$ be an isomorphism. We shall use the notation h^ϕ to denote the image of $h \in H$ under ϕ . Similarly we shall write $a^b := b^{-1}ab$ for conjugation.

THEOREM 6.2 (Application to HNN extensions). *Let H and H' be two isomorphic subgroups of the group A under the isomorphism $h \rightarrow h^\phi, h \in H$. Let B be a group and let $w \in A * B - A$ have torsion free factors. Then the natural map*

$$A \rightarrow \langle A, B \mid w^{-1}hw = h^\phi, h \in H \rangle$$

is injective.

Proof. Consider the following groups

$$A' = \langle A, t \mid t^{-1}ht = h^\phi, h \in H \rangle,$$

$$A'' = \langle A, t, B \mid t^{-1}ht = h^\phi, [a, t^{-1}w] = 1, [t, w] = 1, h \in H, a \in A \rangle$$

$$= \langle A', B \mid [a, t^{-1}w] = 1, [t, w] = 1, a \in A \rangle,$$

$$A''' = \langle A, B \mid w^{-1}hw = h^\phi, h \in H \rangle.$$

We can construct the following commuting diagram,

$$\begin{array}{ccccc} & & & & A''' \\ & & \delta & & \downarrow \gamma \\ & & \nearrow & & \\ A & \xrightarrow{\alpha} & A' & \xrightarrow{\beta} & A'' \end{array}$$

where the maps α, β, γ and δ are induced by inclusion. In order for γ to be a well defined homomorphism it is necessary to check that the relation $w^{-1}hw = h^\phi$, $h \in H$ is a consequence of the relations $t^{-1}ht = h^\phi$, $[a, t^{-1}w] = 1$, $[t, w] = 1$, $h \in H$, $a \in A$. But this follows because $w^{-1}hw = w^{-1}tt^{-1}htt^{-1}w = w^{-1}th^\phi t^{-1}w = h^\phi$. Now α is injective because A' is an HNN extension of A (see [DD, p. 33] or [Se, p. 9]) and β is injective because of theorem 6.1. So δ is injective and this proves the theorem. $\square$

THEOREM 6.3. *Let*

$$(*) \quad u_i(t) = 1, i \in I$$

*be a set of equations over the group A where the exponent sum of t in each $u_i(t)$ is zero. Suppose $w = w(t) \in A * \langle t \rangle - A$ and the factors of w are all torsion free. Then the set of equations*

$$(**) \quad u_i(w(t)) = 1, i \in I$$

has a solution over A if and only if the set $()$ has a solution over A .*

Proof. Let $w(t) = at$ where $a \in A$ has infinite order. Then a solution x for $u_i(w(t)) = 1$ defines a solution at for $(*)$.

Conversely suppose $x \in A'$ is a solution of the set of equations $\{u_i(t) = 1 \mid i \in I\}$. Let G be the subgroup of A' generated by

$$\{x^{-n}ax^n \mid a \in A, n \in \mathbf{Z}\}.$$

Then A is a subgroup of G and G is a subgroup of

$$H = \langle G, t \mid w^{-1}gw = g^\phi, g \in G \rangle$$

where $g^\phi = x^{-1}gx$ by theorem 6.2. Because of the exponent sum condition $u_i(w) = 1, i \in I$. $\square$

REFERENCES

- [BRS] BUONCHRISTIANO, S., B.J. SANDERSON and C.P. ROURKE. *A geometric approach to homology theory, VII: the geometry of CW complexes*. London Maths. Soc. Lecture Note Series 18, 131-149, C.U.P. (1976).
- [DD] DICKS, W. and M. DUNWOODY. *Groups acting on Graphs*. Cambridge Studies in Advanced Maths. 17, C.U.P. (1983).
- [EH] EDJVET, M. and J. HOWIE. The solution of length four equations over groups. *Trans. Amer. Math. Soc.* 326 (1991), 345-369.
- [F] FENN, R.A. *Techniques of Geometric Topology*. London Maths. Soc. Lecture Note Series 57, C.U.P. (1983).