

2.3 Frequencies of factors of Sturmian sequences

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **44 (1998)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARK. In fact, one can prove the following: the frequencies of factors of length n take at most $p(n+1) - p(n) + r_n + l_n$ values, where r_n (respectively, l_n) denotes the number of factors having more than one right (respectively, left) extension.

We deduce from this theorem that if $p(n+1) - p(n)$ is uniformly bounded with n , the frequencies of factors of given length take a finite number of values. Indeed, using a theorem of Cassaigne quoted below (see [10]), we can easily state the following corollary.

THEOREM 7. *If the complexity $p(n)$ of a sequence with values in a finite alphabet is sub-affine then $p(n+1) - p(n)$ is bounded.*

COROLLARY 1. *If a sequence over a finite alphabet has a sub-affine complexity, then the frequencies of its factors of given length take a finite number of values.*

Examples of sequences with sub-affine complexity function include the fixed point of a uniform substitution (i.e., of a substitution such that the images of the letters have the same length), the coding of a rotation or the coding of the orbit of a point under an interval exchange map with respect to the intervals of the interval exchange map.

2.3 FREQUENCIES OF FACTORS OF STURMIAN SEQUENCES

In particular, in the Sturmian case ($p(n) = n + 1$, for every integer n), Theorem 6 implies the following result (see [3]).

THEOREM 8. *The frequencies of factors of given length of a Sturmian sequence take at most three values.*

Consider a Sturmian sequence of angle α . We have seen in Section 2.1 that the frequency of a factor $w_1 \dots w_n$ of u is equal to the length of the interval

$$I(w_1, \dots, w_n) = \bigcap_{j=0}^{n-1} R^{-j}(I_{w_{j+1}}),$$

and that these sets $I(w_1, \dots, w_n)$ are exactly the intervals bounded by the points

$$0, \{1 - \alpha\}, \dots, \{n(1 - \alpha)\}.$$

We deduce from Theorem 8 that the lengths of the intervals $I(w_1, \dots, w_n)$, and thus the lengths of the intervals obtained by placing the points $0, \{1 - \alpha\}, \dots, \{n(1 - \alpha)\}$ on the unit circle, take at most three values. Hence Theorem 8 is equivalent to the three distance theorem and provides a combinatorial proof of this result.

REMARK. In fact this point of view, and more precisely the study of the evolution of the graphs of words with respect to the length n of the factors, allows us to give a proof of the most complete version of the three distance theorem as given in [53] (for more details, the reader is referred to [3]).

3. THE THREE DISTANCE THEOREM

The three distance theorem was initially conjectured by Steinhaus, first proved by V.T. Sós (see [53] and also [52]), and then by Świerczkowski [56], Surányi [55], Slater [51], Halton [31]. More recent proofs have also been given by van Ravenstein [44] and Langevin [35]. A survey of the different approaches used by these authors is to be found in [44, 51, 35]. In the literature this theorem is called *the Steinhaus theorem*, *the three length*, *three gap* or *the three step theorem*. In order to avoid any ambiguity, we will always call it the three distance theorem, reserving the name *three gap* for the theorem introduced in the next section.

THREE DISTANCE THEOREM. *Let $0 < \alpha < 1$ be an irrational number and n a positive integer. The points $\{i\alpha\}$, for $0 \leq i \leq n$, partition the unit circle into $n+1$ intervals, the lengths of which take at most three values, one being the sum of the other two.*

More precisely, let $(\frac{p_k}{q_k})_{k \in \mathbb{N}}$ and $(c_k)_{k \in \mathbb{N}}$ be the sequences of convergents and partial quotients associated to α in its continued fraction expansion (if $\alpha = [0, c_1, c_2, \dots]$, then $\frac{p_n}{q_n} = [0, c_1, \dots, c_n]$). Let $\eta_k = (-1)^k(q_k\alpha - p_k)$. Let n be a positive integer. There exists a unique expression for n of the form

$$n = mq_k + q_{k-1} + r,$$

with $1 \leq m \leq c_{k+1}$ and $0 \leq r < q_k$. Then the circle is divided by the points $0, \{\alpha\}, \{2\alpha\}, \dots, \{n\alpha\}$ into $n+1$ intervals which satisfy: