

4.1 Connectedness index

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **44 (1998)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **18.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARKS.

• Suppose that α is an irrational number. By density of the sequence $(\{n\alpha\})_{n \in \mathbb{N}}$, this theorem still holds when considering the gaps between the successive integers k such that $\{\alpha k\} \in I$, where I denotes any interval of the unit circle of length β .

• Furthermore, the third gap, which is the largest, can have frequency 0, when $\eta_k = \psi$, with the above notation. This means that this gap does not appear at all, as a consequence of the uniform distribution of the sequence $(\{n\alpha\})_{n \in \mathbb{N}}$ in the circle.

• The other two gaps do always appear (infinitely often, in fact, because of their positive frequencies) and are shown to be equal to the smallest positive integers l_1 and l_2 such that $\{l_1\alpha\} < \beta$ and $\{l_2\alpha\} > 1 - \beta$ (see [51]).

• The study of the rational case proves the equivalence between the three distance and the three gap theorems, as observed by Slater [51] in the case of an open interval and by Langevin, for any interval, in [35].

4.1 CONNECTEDNESS INDEX

Let $u = (u_n)_{n \in \mathbb{N}}$ be a coding of a rotation by irrational angle $0 < \alpha < 1$ with respect to the partition

$$\mathcal{P} = \{[\beta_0, \beta_1[, [\beta_1, \beta_2[, \dots, [\beta_{p-1}, \beta_p[\} .$$

We have seen in Section 2.1 that the sets $I(w_1, \dots, w_n) = \bigcap_{j=0}^{n-1} R^{-j}(I_{w_{j+1}})$, where $I_k = [\beta_k, \beta_{k+1}[$, for $0 \leq j \leq p-1$, are connected except for $w_1 \cdots w_n$ of the form a_K^n , where K denotes the index of the interval of $\mathcal{P}$ (if such an interval exists) of length greater than $\sup(\alpha, 1 - \alpha)$.

Let us suppose that there exists an interval of $\mathcal{P}$ of length L greater than $1 - \alpha$ and index K , say. We deduce from the three gap theorem that the set of integers n such that a_K^n is a factor of the sequence u is bounded. More precisely, let us define $n^{(1)}$ as the largest integer n such that a_K^n is a factor of the sequence u . We will call the integer $n^{(1)}$ the *index of connectedness* of the sequence u . (If every interval of $\mathcal{P}$ has length smaller than or equal to $\sup(\alpha, 1 - \alpha)$ then the connectedness index of u is equal to 1.) The three gap theorem enables us to give an exact expression for the connectedness index. Indeed $n^{(1)} + 1$ is the largest gap between the consecutive values of k for which $0 < \{k\alpha\} < 1 - L$. We thus have the following

THEOREM 9. Let $u = (u_n)_{n \in \mathbb{N}}$ be a coding of the rotation by irrational angle α . Suppose that there exists an interval of $\mathcal{P}$ of length $L > \sup(\alpha, 1 - \alpha)$. Let $(\frac{p_k}{q_k})_{k \in \mathbb{N}}$ and $(c_k)_{k \in \mathbb{N}}$ be the sequences of convergents and partial quotients associated to α in its continued fraction expansion. Let $\eta_k = (-1)^k(q_k\alpha - p_k)$. Write

$$1 - L = m\eta_k + \eta_{k+1} + \psi,$$

with $k \geq 1$, $0 < \psi \leq \eta_k$ and $1 \leq m \leq c_{k+1}$. The connectedness index $n^{(1)}$ of the sequence u satisfies

$$\begin{aligned} n^{(1)} &= q_{k+1} - (m-1)q_k - 1, \text{ if } \psi \neq \eta_k, \\ n^{(1)} &= q_{k+1} - mq_k - 1, \text{ if } \psi = \eta_k \text{ and } m < c_{k+1}, \\ n^{(1)} &= q_k - 1, \text{ if } \psi = \eta_k \text{ and } m = c_{k+1}. \end{aligned}$$

4.2 APPLICATIONS

Precise knowledge of the connectedness index is useful, as shown by the following. Indeed Lemma 1 can be rephrased as follows.

LEMMA 3. Let u be a coding of an irrational rotation on the unit circle with respect to the partition $\{[\beta_0, \beta_1[, [\beta_1, \beta_2[, \dots, [\beta_{p-1}, \beta_p[\}$. The frequencies of factors of u of length $n \geq n^{(1)}$, where $n^{(1)}$ denotes the connectedness index, are equal to the lengths of the intervals bounded by the points

$$\{k(1 - \alpha) + \beta_i\}, \text{ for } 0 \leq k \leq n - 1, \quad 0 \leq i \leq p - 1.$$

The complexity of a coding on p letters of an irrational rotation ultimately has the form $p(n) = an + b$, where $a \leq p$, and depends on the algebraic relations between the angle and the lengths of the intervals of the coding. More precisely, we have the following theorem proved in [1].

THEOREM 10. Let $u = (u_n)_{n \in \mathbb{N}}$ be a coding of the irrational rotation R of irrational angle α with respect to the partition

$$\mathcal{P} = \{[\beta_0, \beta_1[, [\beta_1, \beta_2[, \dots, [\beta_{p-1}, \beta_p[\}.$$

Let $(k_n)_{n \in \mathbb{N}}$ be the sequence defined by

$$k_0 = p = \text{card}(F),$$

$$k_n = \text{card} \{ \beta_i \in F; \forall k \in [1, \dots, n], R^{-k}(\beta_i) \notin F \}.$$