

2.2 Group-theoretic Radon transforms

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2.2 GROUP-THEORETIC RADON TRANSFORMS

Let G be a real Lie group and K a (closed) Lie subgroup, equipped with left-invariant Haar measures dg , dk respectively. If the homogeneous space G/K admits a G -invariant measure $d(gK)$, the measures can then be normalized so that

$$\int_G f(g) dg = \int_{G/K} d(gK) \int_K f(gk) dk,$$

for any $f \in C_c(G)$. This applies in particular if K is compact (on invariant measures, see [9], Chap. I, § 1).

Throughout the paper G will be a Lie group, K a compact subgroup, and H a (closed) Lie subgroup of G . The Haar measure dk of K will be normalized by $\int_K dk = 1$.

Let u be a (complex-valued) function on $X = G/K$. Its Radon transform is the function Ru on $Y = G/H$ defined by

$$Ru(gH) = \int_H u(ghK) dh,$$

for $g \in G$, whenever this makes sense (e.g. if $u \in C_c(X)$). The left invariance of dh implies that the integral only depends on the left coset gH of g . Given $y = gH$ in $Y = G/H$, the value $Ru(y)$ is an integral of u over all x incident to y . A more precise statement can be given in the following important example.

EXAMPLE. Let X be a connected Riemannian manifold, G a transitive Lie group of isometries of X and K the isotropy subgroup of some origin $x_o \in X$; then K is compact ([8], p.204) and $X = G/K$. Let y_o be a given closed submanifold of X , containing x_o , and let Y be the set of all submanifolds $y = g \cdot y_o$ of X , with $g \in G$.

The set H of all $h \in G$ such that $h \cdot y_o = y_o$ (i.e. the submanifold y_o is globally invariant under h) is a closed Lie subgroup of G . Indeed if $h_n \in H$ converges to h in G , for any $x \in y_o$ the point $\lim h_n \cdot x = h \cdot x$ belongs to y_o ; similarly $h^{-1} \cdot x \in y_o$, so that $h \cdot y_o = y_o$. Thus $Y = G/H$ can be endowed with a structure of manifold and we obtain a double fibration of homogeneous spaces.

The following lemma allows one to compute the Radon transform without knowing H explicitly.

LEMMA 1. *Keeping the notation of this example, assume furthermore that $y_o = G' \cdot x_o$ is a closed orbit of the origin $x_o = K$ under some Lie subgroup G' of G .*

Then $G' \subset H \subset G'K$ and $y_o = H \cdot x_o$. The incidence relation between $X = G/K$ and $Y = G/H$ is simply $x \in y$. Besides, the left-invariant Haar measures dh, dg' of the groups H, G' can be normalized so that

$$\begin{aligned} Ru(y) &= \int_H u(gh \cdot x_o) dh = \int_{G'} u(gg' \cdot x_o) dg' \\ &= \int_y u(x) dm_y(x), \end{aligned}$$

where dm_y is the Riemannian measure induced by X on its submanifold $y = g \cdot y_o$.

REMARK. The subgroup H can of course be strictly bigger than G' . This occurs for instance if y_o is a line in $X = \mathbf{R}^n$ and G' is the group of translations along this line, or a horocycle in a Riemannian symmetric space X of the noncompact type (for which $G' = N$ and $H = MN = NM$ in the usual semisimple notations).

Proof of Lemma 1. If $y_o = G' \cdot x_o$, then H obviously contains G' and it follows that

$$y_o = G' \cdot x_o \subset H \cdot x_o \subset y_o,$$

whence $H \cdot x_o = G' \cdot x_o$ and $H \subset G'K$.

A point $x \in X$ is incident to $y = g \cdot y_o \in Y$ if and only if there exists $h \in H$ such that $x = gh \cdot x_o$, i.e. $x \in gH \cdot x_o = g \cdot y_o = y$.

An isometry g transforms the Riemannian measure of y_o into the Riemannian measure of $y = g \cdot y_o$, and it suffices to prove the integral formula for $g = e$. Now $y_o = H \cdot x_o$ can be identified to the homogeneous space $H/(H \cap K)$, and dm_{y_o} (which is invariant under all isometries of y_o) to an H -invariant measure on this space. The Haar measure dh can therefore be normalized so that the corresponding measure on $H/(H \cap K)$ satisfies

$$\begin{aligned} \int_{y_o} u(x) dm_{y_o}(x) &= \int_{H/(H \cap K)} u(h \cdot x_o) d(h(H \cap K)) \\ &= \int_H u(h \cdot x_o) dh = Ru(y_o). \end{aligned}$$

The proof is similar for $\int_{G'}$, whence the lemma. □

Going back to general double fibrations, the *Radon dual transform* of a (continuous, say) function v on $Y = G/H$ is the function on $X = G/K$ defined by

$$R^*v(gK) = \int_K v(gkH) dk,$$

for $g \in G$, an integral of v over all y incident to $x = gK$. The word “dual” is of course motivated by the classical projective duality between points and hyperplanes in the basic example, but it also stems from the following proposition.

PROPOSITION 2. *Let $X = G/K$ with K compact, and assume that $Y = G/H$ has a G -invariant measure. Let $u \in C_c(X)$, $v \in C(Y)$. Then $Ru \in C_c(Y)$, $R^*v \in C(X)$ and*

$$\int_X u(x) R^*v(x) dx = \int_Y Ru(y) v(y) dy = \int_Z u(x) v(y) dz,$$

where dx, dy, dz are the respective G -invariant measures on X, Y and $Z = G/(K \cap H)$.

In the latter integral $u(x)v(y)$ is considered as a function of $z = (x, y)$ on Z (Section 2.1). We omit the proof, a classical exercise on invariant integrals (cf. [9], p. 144 and [11], p. 41); all groups are assumed unimodular there, but the proof only uses the invariant measures on the homogeneous spaces, thus extends to the present situation.

Proposition 2 allows a natural extension of the transforms R and R^* to distributions. Given $u \in \mathcal{E}'(X)$, the distribution $Ru \in \mathcal{E}'(Y)$ is defined by

$$\langle Ru, v \rangle = \langle u, R^*v \rangle,$$

for all test functions $v \in C^\infty(Y)$. Similarly, given $v \in \mathcal{D}'(Y)$, the distribution $R^*v \in \mathcal{D}'(X)$ is defined by

$$\langle R^*v, u \rangle = \langle v, Ru \rangle,$$

for all $u \in \mathcal{D}(X)$. Again we refer to Helgason ([11], p. 42) for details, based on the compactness of K . These definitions do extend the Radon integrals for functions, as Proposition 2 shows, when identifying a function u with the distribution $u(x) dx$, and similarly for v .