

2. REVIEW OF THE L^2 -INDEX THEOREM

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

ATIYAH'S L^2 -INDEX THEOREM

by Indira CHATTERJI and Guido MISLIN

1. INTRODUCTION

The L^2 -Index Theorem of Atiyah [1] expresses the index of an elliptic operator on a closed manifold M in terms of the G -equivariant index of some regular covering $\tilde{M}$ of M , with G the group of covering transformations. Atiyah's proof is analytic in nature. Our proof is algebraic and involves an embedding of a given group into an acyclic one, together with naturality properties of the indices.

2. REVIEW OF THE L^2 -INDEX THEOREM

The main reference for this section is Atiyah's paper [1]. All manifolds considered are smooth Riemannian, without boundary. Covering spaces of manifolds carry the induced smooth and Riemannian structure. Let M be a closed manifold and let E, F denote two complex (Hermitian) vector bundles over M . Consider an elliptic pseudo-differential operator

$$D: C^\infty(M, E) \rightarrow C^\infty(M, F)$$

acting on the smooth sections of the vector bundles. One defines its space of solutions

$$S_D = \{s \in C^\infty(M, E) \mid Ds = 0\}.$$

The complex vector space S_D has finite dimension (see [13]), and so has S_{D^*} the space of solutions of the adjoint D^* of D where

$$D^*: C^\infty(M, F) \rightarrow C^\infty(M, E)$$

is the unique continuous linear map satisfying

$$\langle Ds, s' \rangle = \int_M \langle Ds(m), s'(m) \rangle_F dm = \langle s, D^* s' \rangle = \int_M \langle s(m), D^* s'(m) \rangle_E dm$$

for all $s \in C^\infty(M, E)$, $s' \in C^\infty(M, F)$. One now defines the *index* of D as follows:

$$\text{Index}(D) = \dim_{\mathbf{C}}(S_D) - \dim_{\mathbf{C}}(S_{D^*}) \in \mathbf{Z}.$$

An explicit formula for $\text{Index}(D)$ is given by the famous Atiyah-Singer Theorem (cf. [2]). Consider a not necessarily connected, regular covering $\pi: \tilde{M} \rightarrow M$ with countable covering transformation group G . The projection π can be used to define an elliptic operator

$$\tilde{D} := \pi^*(D): C_c^\infty(\tilde{M}, \pi^*E) \rightarrow C_c^\infty(\tilde{M}, \pi^*F).$$

Denote by $S_{\tilde{D}}$ the closure of $\{s \in C_c^\infty(\tilde{M}, \pi^*E) \mid \tilde{D}s = 0\}$ in $L^2(\tilde{M}, \pi^*E)$. Let $\tilde{D}^*$ denote the adjoint of $\tilde{D}$. The space $S_{\tilde{D}}$ is not necessarily finite dimensional, but being a closed G -invariant subspace of the L^2 -completion $L^2(\tilde{M}, \pi^*E)$ of the space of smooth sections with compact supports $C_c^\infty(\tilde{M}, \pi^*E)$, its von Neumann dimension is therefore defined as follows. Write

$$\mathcal{N}(G) = \{P: \ell^2(G) \rightarrow \ell^2(G) \text{ bounded and } G\text{-invariant}\}$$

for the group von Neumann algebra of G , where G acts on $\ell^2(G)$ via the right regular representation. Then $S_{\tilde{D}}$ is a finitely generated Hilbert G -module and hence can be represented by an idempotent matrix $P = (p_{ij}) \in M_n(\mathcal{N}(G))$ (recall that a finitely generated Hilbert G -module is isometrically G -isomorphic to a Hilbert G -subspace of the Hilbert space $\ell^2(G)^n$ for some $n \geq 1$, see [9]). One then sets

$$\dim_G(S_{\tilde{D}}) = \sum_{i=1}^n \langle p_{ii}(e), e \rangle = \kappa(P) \in \mathbf{R},$$

where by abuse of notation e denotes the element in $\ell^2(G)$ taking value 1 on the neutral element $e \in G$ and 0 elsewhere (see Eckmann's survey [9] on L^2 -cohomology for more on von Neumann dimensions). The map $\kappa: M_n(\mathcal{N}(G)) \rightarrow \mathbf{C}$ is the Kaplansky trace. One defines the L^2 -index of $\tilde{D}$ by

$$\text{Index}_G(\tilde{D}) = \dim_G(S_{\tilde{D}}) - \dim_G(S_{\tilde{D}^*}).$$

We can now state Atiyah's L^2 -Index Theorem.

THEOREM 2.1 (Atiyah [1]). For D an elliptic pseudo-differential operator on a closed Riemannian manifold M

$$\text{Index}(D) = \text{Index}_G(\tilde{D})$$

for any countable group G and any lift $\tilde{D}$ of D to a regular G -cover $\tilde{M}$ of M .

In particular, the L^2 -index of $\tilde{D}$ is always an integer, even though it is a priori given in terms of real numbers. The following serves as an illustration of the L^2 -Index Theorem.

EXAMPLE 2.2 (Atiyah's formula [1]). Let $\Omega^\bullet$ be the de Rham complex of complex valued differential forms on the closed connected manifold M and consider the de Rham differential $D = d + d^*: \Omega^{ev} \rightarrow \Omega^{odd}$. Let $\pi: \tilde{M} \rightarrow M$ be the universal cover of M so that $G = \pi_1(M)$. Then

- $\text{Index}(D) = \chi(M)$, the ordinary Euler characteristic of M .
- $\text{Index}_G(\tilde{D}) = \sum_j (-1)^j \beta^j(M)$, the L^2 -Euler characteristic of M .

The $\beta^j(M)$'s denote the L^2 -Betti numbers of M . Thus the L^2 -Index Theorem translates into Atiyah's formula

$$\chi(M) = \sum_j (-1)^j \beta^j(M).$$

We recall that the L^2 -Betti numbers $\beta^j(M)$ are in general not integers. For instance, if $\pi_1(M)$ is a finite group, one checks that

$$\beta^j(M) = \frac{1}{|\pi_1(M)|} b^j(\tilde{M}),$$

where $b^j(\tilde{M})$ stands for the ordinary j 'th Betti number of the universal cover $\tilde{M}$ of M . In particular, for $1 < |\pi_1(M)| < \infty$, $\beta^0(M) = 1/|\pi_1(M)|$ is not an integer and the L^2 -Index Theorem reduces to the well-known fact that

$$\chi(M) = \frac{\chi(\tilde{M})}{|\pi_1(M)|}.$$

It is a conjecture (Atiyah Conjecture) that for a general closed connected manifold M the L^2 -Betti numbers $\beta^j(M)$ are always rational numbers, and even integers in case that $\pi_1(M)$ is torsion-free. For some interesting examples, which might lead to counterexamples, see Dicks and Schick [8].