

Construction of classifying spaces with isotropy in prescribed families of subgroups

Autor(en): **Lafont, Jean-François**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **22.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109915>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

46

CONSTRUCTION OF CLASSIFYING SPACES WITH ISOTROPY IN PRESCRIBED FAMILIES OF SUBGROUPS

by Jean-François LAFONT

For an infinite group Γ , the Farrell–Jones isomorphism conjecture [3] states that the algebraic K -theory $K_n(\mathbf{Z}\Gamma)$ of the integral group ring of Γ coincides with $H_n^\Gamma(E_{\text{VC}}\Gamma; \mathbf{K}\mathbf{Z}^{-\infty})$, a certain equivariant generalized homology theory of the Γ -space $E_{\text{VC}}\Gamma$. This space is a model for the classifying space for Γ with isotropy in the family of virtually cyclic subgroups, i.e. a contractible Γ -CW-complex with the property that the fixed subset of a subgroup H is contractible if H is a virtually cyclic subgroup, and is empty otherwise. Such a space is unique up to Γ -equivariant homotopy equivalence. From such a classifying space, the homology $H_n^\Gamma(E_{\text{VC}}\Gamma; \mathbf{K}\mathbf{Z}^{-\infty})$ can be computed via an Atiyah–Hirzebruch type spectral sequence discovered by Quinn [9]. The ingredients entering into the E^2 -term of the spectral sequence are the algebraic K -theory of the various cell-stabilizers. In particular, for computational purposes, it is interesting to have a model for $E_{\text{VC}}\Gamma$ that is as “small” as possible. Let us denote by $\text{hdim}^\Gamma(X)$, for a Γ -space X , the minimal dimension of a CW-complex Γ -equivariantly homotopy equivalent to X . The discussion above motivates the first:

PROBLEM 46.1. *Find an efficient algebraic criterion that determines whether a finitely generated group Γ has a finite dimensional model for $E_{\text{VC}}\Gamma$, i.e. whether $\text{hdim}^\Gamma(E_{\text{VC}}\Gamma) < \infty$.*

In particular, one can consider the following problem:

PROBLEM 46.2. *For various classical families of finitely-generated groups appearing in mathematics, either (1) give a construction for a finite dimensional model for $E_{\text{VC}}\Gamma$, or (2) prove that some group within the family satisfies $\text{hdim}^\Gamma(E_{\text{VC}}\Gamma) = \infty$.*

A few classes of groups are known to have finite dimensional models for $E_{\mathcal{VC}}\Gamma$. For instance, it is clear that for virtually cyclic group, one can take a point as a model for $E_{\mathcal{VC}}\Gamma$. Less trivial examples consist of:

- δ -hyperbolic groups (due to Juan-Pineda and Leary [4], and independently to Lück [6]),
- crystallographic groups (different finite dimensional models were given by Alves and Ontaneda [1], and by Connolly, Fehrman and Hartglass [2]),
- groups hyperbolic relative to subgroups which themselves have finite dimensional classifying spaces, for instance non-uniform lattices in $\mathrm{SO}(n, 1)$ (due to Lafont and Ortiz [5]).
- virtually poly- \mathbf{Z} groups, and groups which are countable locally virtually cyclic (due to Lück and Weiermann [7]).

In general, given a family \mathcal{F} of subgroups of Γ , one can define a model for the classifying space $E_{\mathcal{F}}\Gamma$ of Γ with isotropy in the family \mathcal{F} (see the extensive survey in [6]), which will again be unique up to Γ -equivariant homotopy equivalence. For the family \mathcal{FIN} consisting of finite subgroups, the classifying space $E_{\mathcal{FIN}}\Gamma$ has been extensively studied, and explicit finite dimensional models are known for various classes of groups (δ -hyperbolic groups, groups acting by isometries on finite dimensional $\mathrm{CAT}(0)$ spaces, Coxeter groups, etc).

In a paper with I. Ortiz [5], we defined the notion of a collection of subgroups to be *adapted* to a nested pair $\mathcal{F} \subset \overline{\mathcal{F}}$ of families of subgroups (for instance, one could take $\mathcal{FIN} \subset \mathcal{VC}$). This consists of a collection of subgroups $\{H_\alpha\}$ satisfying the following properties: (1) the collection is conjugacy closed, (2) the groups H_α are self-normalizing, (3) distinct groups in the collection intersect in elements of \mathcal{F} , and (4) every group in $\overline{\mathcal{F}} - \mathcal{F}$ is contained in one of the H_α .

When there exists a collection of subgroups adapted to a pair $\mathcal{F} \subset \overline{\mathcal{F}}$, we explain how to modify a model for $E_{\mathcal{F}}\Gamma$ to obtain a model for $E_{\overline{\mathcal{F}}}\Gamma$. The modifications involve the collection of classifying spaces $E_{\overline{\mathcal{F}}(H_\alpha)}H_\alpha$, where $\overline{\mathcal{F}}(H_\alpha)$ is the restriction of the family $\overline{\mathcal{F}}$ to the subgroup H_α . In particular, when both the $E_{\mathcal{F}}\Gamma$ and the $E_{\overline{\mathcal{F}}(H_\alpha)}H_\alpha$ are finite dimensional, the construction yields a finite dimensional $E_{\overline{\mathcal{F}}}\Gamma$. This prompts the following

PROBLEM 46.3. *Try to identify “natural” non-trivial collections of subgroups adapted to the pair $\mathcal{FIN} \subset \mathcal{VC}$ for various classical families of finitely-generated groups.*

In all the examples the author knows of where the minimal dimensions of models for $E_{\mathcal{FIN}}\Gamma$ and $E_{\mathcal{VC}}\Gamma$ are explicitly known, one has that both these numbers are finite. It is known that if $\text{hdim}^\Gamma(E_{\mathcal{FIN}}\Gamma) = \infty$, then $\text{hdim}^\Gamma(E_{\mathcal{VC}}\Gamma) = \infty$ (see [7], Cor. 5.4). The converse is likely to be false, and one can ask:

PROBLEM 46.4. *Find examples of finitely generated groups Γ for which (1) $\text{hdim}^\Gamma(E_{\mathcal{FIN}}\Gamma) < \infty$, but (2) $\text{hdim}^\Gamma(E_{\mathcal{VC}}\Gamma) = \infty$.*

One might think that in general, one can find families of subgroups for which the classifying spaces can be arbitrarily complicated, prompting:

PROBLEM 46.5. *For Γ a (non-abelian) infinite group, does there always exist a family \mathcal{F} of subgroups, with $\mathcal{FIN} \subset \mathcal{F}$, and $\text{hdim}^\Gamma(E_{\mathcal{F}}\Gamma) = \infty$?*

Recently, Quinn has suggested a possible refinement of the Farrell–Jones isomorphism conjecture. In his paper [8], Quinn considers *p-hyper-elementary* groups, defined to be groups G that fit into a short exact sequence:

$$1 \rightarrow C \rightarrow G \rightarrow P \rightarrow 1$$

where P is a finite p -group, and C is cyclic. The family \mathcal{HE} of hyper-elementary subgroups of a finitely generated group Γ gives rise to a classifying space $E_{\mathcal{HE}}\Gamma$, and Quinn suggests that the algebraic K -theory $K_n(\mathbb{Z}\Gamma)$ of the integral group ring of Γ coincides with $H_n^\Gamma(E_{\mathcal{HE}}\Gamma; \mathbf{K}\mathbb{Z}^{-\infty})$. Note that every hyper-elementary group is automatically virtually cyclic, hence we have a containment $\mathcal{HE} \subset \mathcal{VC}$. From the computational viewpoint, this refinement would be particularly useful if these classifying spaces $E_{\mathcal{HE}}\Gamma$ were “smaller” than $E_{\mathcal{VC}}\Gamma$. Hence it would again be of interest to obtain concrete models for the $E_{\mathcal{HE}}\Gamma$:

PROBLEM 46.6. *For various classical families of groups, give a construction for a finite dimensional $E_{\mathcal{HE}}\Gamma$. In particular, find an example of a group Γ for which (1) $\text{hdim}^\Gamma(E_{\mathcal{HE}}\Gamma) < \infty$ but (2) $\text{hdim}^\Gamma(E_{\mathcal{VC}}\Gamma) = \infty$.*

So far we have mostly considered families that are smaller than \mathcal{VC} . But in some cases, it is conceivable that the classifying spaces might be *easier* to construct for a *larger* family than \mathcal{VC} . One natural candidate family to consider is the family \mathcal{VA} of virtually abelian subgroups. In particular, constructing a classifying space with isotropy in \mathcal{VA} would be of interest for

groups where one has a fairly good structure theory for the virtually abelian subgroups, and for which the Farrell–Jones isomorphism conjecture is known to hold. To give a concrete example, one can ask:

PROBLEM 46.7. *Give a procedure to construct a finite-dimensional model for $E_{\text{VA}}\Gamma$, when Γ is either (1) a uniform lattice in a higher rank symmetric space, or (2) an irreducible, non-affine, infinite Coxeter group.*

REFERENCES

- [1] ALVES, A. and P. ONTANEDA. A formula for the Whitehead group of a three-dimensional crystallographic group. *Topology* 45 (2006), 1–25.
- [2] CONNOLLY, F., B. FEHRMAN and M. HARTGLASS. On the dimension of the virtually cyclic classifying space of a crystallographic group. Preprint arXiv: math.AT/0610387 (2006).
- [3] FARRELL, F. T. and L. E. JONES. Isomorphism conjectures in algebraic K -theory. *J. Amer. Math. Soc.* 6 (1993), 249–297.
- [4] JUAN-PINEDA, D. and I. J. LEARY. On classifying spaces for the family of virtually cyclic subgroups. In: *Recent Developments in Algebraic Topology, 2003*, 135–145. *Contemp. Math.* 407. Amer. Math. Soc., 2006.
- [5] LAFONT, J.-F. and I. J. ORTIZ. Relative hyperbolicity, classifying spaces, and lower algebraic K -theory. *Topology* 46 (2007), 527–553.
- [6] LÜCK, W. Survey on classifying spaces for families of subgroups. In: *Infinite Groups: Geometric, Combinatorial and Dynamical Aspects*, 269–322. *Progress in Mathematics* 248. Birkhäuser, Basel, 2005.
- [7] LÜCK, W. and M. WEIERMANN. On the classifying space of the family of virtually cyclic subgroups. Preprint arXiv: math.AT/0702646 (2007).
- [8] QUINN, F. Hyperbolic assembly for K -theory of virtually abelian groups. Preprint arXiv: math.KT/0509294 (2005–2006).
- [9] ———. Ends of maps, II. *Invent. Math.* 68 (1982), 353–424.

J.-F. Lafont

Department of Mathematics
 The Ohio State University
 231 West 18th Avenue
 Columbus, OH 43210-1174
 USA
e-mail: jlafont@math.ohio-state.edu