

The extended quotient

Autor(en): **Baum, Paul**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **17.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109877>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

8

THE EXTENDED QUOTIENT

by Paul BAUM

Let Γ be a finite group acting on a (topological space X or) an affine variety X . The quotient variety (or quotient topological space) X/Γ is obtained by collapsing each orbit to a point. For $x \in X$, Γ_x denotes the stabilizer group of x , that is

$$\Gamma_x = \{\gamma \in \Gamma \mid \gamma x = x\},$$

and $c(\Gamma_x)$ denotes the set of conjugacy classes of Γ_x .

The *extended quotient* is obtained by replacing the orbit of x by $c(\Gamma_x)$. This is done as follows:

Set $\tilde{X} = \{(\gamma, x) \in \Gamma \times X \mid \gamma x = x\}$, this is an affine variety and is a sub-variety of $\Gamma \times X$. Moreover, Γ acts on $\tilde{X}$ as follows

$$\Gamma \times \tilde{X} \longrightarrow \tilde{X}$$

$$(g, (\gamma, x)) \mapsto g(\gamma, x) = (g\gamma g^{-1}, gx) \quad \text{where } g \in \Gamma, (\gamma, x) \in \tilde{X}.$$

The extended quotient, denoted by $X//\Gamma$, is given by $\tilde{X}/\Gamma$, the ordinary quotient of $\tilde{X}$ by the above action of Γ .

The extended quotient is an affine variety (or a topological space). The evident projection $\tilde{X} \rightarrow X$, $(\gamma, x) \mapsto x$ passes to quotient spaces to give a map $\rho: X//\Gamma \rightarrow X/\Gamma$. The map ρ is the projection of the extended quotient onto the ordinary quotient.

Let G be a reductive p -adic group (examples are $\mathrm{GL}(n, F)$ and $\mathrm{SL}(n, F)$, where F is any finite extension of the p -adic numbers $\mathbf{Q}_p$). Let V be a vector space over the complex numbers $\mathbf{C}$.

DEFINITION 8.1. A representation $\phi: G \rightarrow \mathrm{Aut}_{\mathbf{C}}(V)$ of G is *smooth* if for every $v \in V$,

$$G_v = \{g \in G \mid \phi(g)v = v\}$$

is an open subgroup of G .

We will denote by $\widehat{G}$ the set of equivalence classes of smooth irreducible representations of G . One of the main problems in the representation theory of p -adic groups (which is closely related to the local Langlands conjecture) is to describe $\widehat{G}$.

The *Hecke algebra* of G , denoted by $\mathcal{H}G$, is the convolution algebra of all complex-valued locally-constant compactly-supported functions $f: G \rightarrow \mathbf{C}$. Then $\widehat{G}$ is in bijection with $\text{Prim } \mathcal{H}G$, the set of primitive ideals in $\mathcal{H}G$. On $\text{Prim } \mathcal{H}G$ there is the Jacobson topology. Hence we may consider each connected component of the primitive ideal space. Typically there will be countably many of these connected components.

Let $\mathbf{C}^\times$ denote the (complex) affine variety $\mathbf{C} - \{0\}$.

DEFINITION 8.2. A *complex torus* is a (complex) affine variety T such that there exists an isomorphism of affine varieties

$$T \cong \mathbf{C}^\times \times \mathbf{C}^\times \times \cdots \times \mathbf{C}^\times.$$

J. Bernstein assigns to each $\alpha \in \pi_0(\text{Prim } \mathcal{H}G)$ a complex torus T_α and a finite group Γ_α acting on T_α .

He then forms the quotient variety T_α/Γ_α and proves that there is a surjective map (the infinitesimal character)

$$\pi_\alpha: X_\alpha \twoheadrightarrow T_\alpha/\Gamma_\alpha.$$

The set X_α is the connected component of $\text{Prim } \mathcal{H}G$ corresponding to α . In Bernstein's work X_α is a set (i.e. is only a set) so π_α is a map of sets, which is surjective, finite-to-one and generically one-to-one.

CONJECTURE 8.3. *There is a certain resemblance between*

$$\begin{array}{ccc} T_\alpha//\Gamma_\alpha & & X_\alpha \\ \rho_\alpha \downarrow & \text{and} & \downarrow \pi_\alpha \\ T_\alpha/\Gamma_\alpha & & T_\alpha/\Gamma_\alpha \end{array}$$

Here ρ_α is (as above) the projection of the extended quotient onto the ordinary quotient.

For the precise conjecture, see papers of A.-M. Aubert, P. Baum and R. Plymen [1] and [2], but we now explain what is meant by *resemblance*.

For each $\alpha \in \pi_0(\text{Prim } \mathcal{H}G)$ there exists (conjecturally) a bijection

$$\nu_\alpha: T_\alpha // \Gamma_\alpha \longrightarrow X_\alpha$$

such that :

- In the possibly non-commutative diagram

$$\begin{array}{ccc} T_\alpha // \Gamma_\alpha & \xrightarrow{\nu_\alpha} & X_\alpha \\ \rho_\alpha \downarrow & & \downarrow \pi_\alpha \\ T_\alpha / \Gamma_\alpha & \xrightarrow{I} & T_\alpha / \Gamma_\alpha \end{array}$$

the bijection $\nu_\alpha: T_\alpha // \Gamma_\alpha \longrightarrow X_\alpha$ is continuous, where $T_\alpha // \Gamma_\alpha$ has the Zariski topology, X_α the Jacobson topology, and the composition

$$\pi_\alpha \circ \nu_\alpha: T_\alpha // \Gamma_\alpha \longrightarrow T_\alpha / \Gamma_\alpha$$

is a morphism of algebraic varieties.

- For each $\alpha \in \pi_0(\text{Prim } \mathcal{H}G)$, there is an algebraic family

$$\theta_t: T_\alpha // \Gamma_\alpha \longrightarrow T_\alpha / \Gamma_\alpha$$

of morphisms of algebraic varieties, with $t \in \mathbf{C}^\times$, such that $\theta_1 = \rho_\alpha$ and $\theta_{\sqrt{q}} = \pi_\alpha \circ \nu_\alpha$, where q is the order of the residue field of the p -adic field F over which G is defined, and π_α is the infinitesimal character of Bernstein.

This conjecture is true for $\text{GL}(n, F)$, where n is any positive integer and F is any finite extension of the p -adic numbers $\mathbf{Q}_p$, see [3].

REFERENCES

- [1] AUBERT, A.-M., P. BAUM and R. PLYMEN. The Hecke algebra of a reductive p -adic group: a geometric conjecture. In: *Noncommutative Geometry and Number Theory* (K. Consani and M. Marcolli, eds.) Aspects of Math. Vol. 37, Vieweg, Wiesbaden, 2006, 1–34.

- [2] AUBERT, A.-M., P. BAUM and R. PLYMEN. Geometric structure in the representation theory of p -adic groups. *C.R. Acad. Sci. Paris. Sér. I Math.* 345 (2007), 573–578.
- [3] BRODZKI, J. and R. PLYMEN. Geometry of the smooth dual of $GL(n)$. *C.R. Acad. Sci. Paris Sér. I Math.* 331 (2000), 213–218.

Paul Frank Baum

Mathematics Department
206 McAllister Building
The Pennsylvania State University
University Park, PA 16802
USA
e-mail: baum@math.psu.edu